

1
Rank

Cracking the
JEE Main Code
360 out of 360
Kalpita Veerwal



www.mtg.in | June 2017 | Pages 92 | ₹ 30

ACE YOUR WAY CBSE

Class XI | XII

MATHEMATICS

today

India's #1
MATHEMATICS MONTHLY
for JEE (Main & Advanced)

**MONTHLY
PRACTICE
PROBLEMS**
(XI & XII)

**CHALLENGING
PROBLEMS**

OLYMPIAD CORNER

**MATHS
MUSING**
10 GREAT
PROBLEMS

**WB JEE
KERALA PET
KARNATAKA CET**
SOLVED PAPERS 2017

Z

3

D

X

mtg

Trust of more than
1 Crore Readers
Since 1982



2017100006685

**MATH
ARCHIVES**

**You Ask ?
We Answer ✓**

MATHEMATICS today

Vol. XXXV No. 6 June 2017

Corporate Office:

Plot 99, Sector 44 Institutional Area,
Gurgaon -122 003 (HR), Tel : 0124-6601200
e-mail : info@mtg.in website : www.mtg.in

Regd. Office:

406, Taj Apartment, Near Safdarjung Hospital,
Ring Road, New Delhi - 110029.
Managing Editor : Mahabir Singh
Editor : Anil Ahlawat

CONTENTS

Class XI	8	Topper's Talk
	12	Ace Your Way (Series 2)
	18	MPP-2
Class XII	20	Ace Your Way (Series 2)
	28	MPP-2
Competition Edge	31	Solved Paper - WB JEE
	43	Solved Paper - Kerala PET
	63	Solved Paper - Karnataka CET
	72	Maths Musing Problem Set - 174
	73	Challenging Problems
	77	Math Archives
	80	Olympiad Corner
	84	You Ask We Answer
	85	Maths Musing Solutions



Subscribe online at WWW.mtg.in

Individual Subscription Rates

	1 yr.	2 yrs.	3 yrs.
Mathematics Today	330	600	775
Chemistry Today	330	600	775
Physics For You	330	600	775
Biology Today	330	600	775

Combined Subscription Rates

	1 yr.	2 yrs.	3 yrs.
PCM	900	1500	1900
PCB	900	1500	1900
PCMB	1000	1800	2300

Send D.D./M.O in favour of MTG Learning Media (P) Ltd.
Payments should be made directly to : MTG Learning Media (P) Ltd,
Plot 99, Sector 44 Institutional Area, Gurgaon - 122 003, Haryana.
We have not appointed any subscription agent.

Owned, Printed and Published by MTG Learning Media Pvt. Ltd. 406, Taj Apartment, New Delhi - 29 and printed by HT Media Ltd., B-2, Sector-63, Noida, UP-201307. Readers are advised to make appropriate thorough enquiries before acting upon any advertisements published in this magazine. Focus/Infocus features are marketing incentives. MTG does not vouch or subscribe to the claims and representations made by advertisers. All disputes are subject to Delhi jurisdiction only.
Editor : Anil Ahlawat
Copyright © MTG Learning Media (P) Ltd.
All rights reserved. Reproduction in any form is prohibited.

mtg Team Applauds

JEE (Main) 2017 Topper

“It gives us immense pleasure to felicitate the achievement of our reader Kalpit Veerwal (MTG Subscription Code PCM-102332). We feel proud that we could lay a brick for the foundation of his success. We are sharing his success story here so that it can inspire others to ace in exams.”



Kalpit Veerwal

1
Rank

- **MTG : Why did you appear for Engineering Entrance?**

Kalpit : I had interest in studying Science and Mathematics, so wanted to appear for the entrance exams. Now, I want to opt for Computer Science Engineering.

- **MTG : What exams have you appeared for and what are your ranks in these exams?**

Kalpit : I have appeared for JEE Main 2017 in which I got AIR 1 with 360 marks out of 360. I have also achieved ranks in exams like NSO-7th International rank, IMO-17th International rank (conducted by Science Olympiad Foundation, New Delhi), NTSE (Stage I-Rajasthan 1st rank and Stage II-cleared), KVPY, IJSO Stage II (Top 35), IJAO Stage II (Top 20) and IAO Stage II (Top 25).

- **MTG : How many hours in a day did you dedicate for the preparation of the examination?**

Kalpit : Apart from studying in school and coaching, I used to spend 5-6 hours on self study.

- **MTG : Any extra coaching?**

Kalpit : I pursued coaching from Resonance, Udaipur.

- **MTG : Did you appear for the offline JEE Main exam or the online one? What is the reason behind this choice of yours?**

Kalpit : I appeared for the JEE Main offline exam as I believe

that it is the traditional method. I was not very confident with the online option.

Moreover, I was comfortable with the offline pattern as my coaching institute follows it. Therefore, I chose to take my exam in pen and paper pattern.

- **MTG : On which topic and chapters you laid more stress in each subject?**

Kalpit : I focussed on complete syllabus, rather than selective study. Hence, gave equal weightage to all chapters of each subject.

- **MTG : How much time does one require for serious preparation of this exam?**

Kalpit : I studied sincerely from class VIII but any kind of serious preparation from my side for JEE Main started from class XI. Every day after coming back from school, I studied regularly. I never bunked any of my classes and followed my teachers religiously. I used to wake up early in the morning to study.

“I was a 2 year subscriber of the MTG magazines: Physics For You, Chemistry Today and Mathematics Today and was really an avid reader of them. They were highly useful especially because of the good question bank and articles published in every issue. I also studied from NCERT books and solved JEE Main previous years question papers.”

- **MTG : Was there a difference in the preparation strategy during the last months of JEE Main considering that the board exams were also scheduled in this period? How did you manage the preparation for both?**

Kalpit : As you know that board exams were scheduled in March and JEE Main 2017 was scheduled in April so I revised

THANKS FOR MAKING US PROUD

120⁺ Selection in JEE Main-2017

**1st
in C.G.**



Tushar Agrawal

AIR-223



Keshav Gupta

AIR-232



Ashish R. Nair

AIR-323



Ashutosh Chaubey

AIR-575



Ujjawal Sharma

AIR-836



Kunal Das

AIR-844



Gaurav Singhal

AIR-1730



Arooshi Verma

AIR-2463



Vinamra Bathwal

AIR-3070



Jheel Rathoo

AIR-3191



Shrirang Vaidya

AIR-3231



Pravartya Kr. Dewangan

AIR-3496



Harsh Prakash Gupta

AIR-4122



Srijan Kr. Upadhyay

AIR-4849



Anurag Nayak

AIR-5214



Nitin Vinayak Agrawal

AIR-5961



Pranjal Gupta

AIR-6629



Swasti Shreya Mishra

AIR-6978



Tavishi Jatrelle

AIR-8753



Mrudul Kulkarni

AIR-8754

**KCS
Educate**

...revives internal teacher

Knowledge Centre for Success Educate Pvt. Ltd.

Team Avnish

for JEE | Aptitude Test

NTSE | KVPY | Olympiad

BHILAI OFFICE 157, New Civic Centre, Bhilai, Dist. Durg (C.G.)

Telephone : **0788-6454505**

DELHI OFFICE 1B, Block GG1, Vikaspuri, New Delhi - 110018

CIN No. U74140DL2011PTC227887

info@kcseducate.in

www.kcseducate.in

facebook.com/kcseducate

my complete syllabus of class XII boards in November and once again in February. When one month was left I revised the entire syllabus from NCERT.

- **MTG : Which books/magazines you read?**

Kalpita : I read HC Verma, Resnik, Irodov, SL Loney and JD Lee.

- **MTG : In your words what are the components of an ideal preparation plan?**

Kalpita : There are 3 main components of an ideal preparation plan : Focus, Practice and Dedication. Stick to the basics and master the speed with the help of regular tests.

- **MTG : What role did the following play in your success:**

(a) Parents (b) Teachers and School

Kalpita : (a) My parents took care of my health and kept me stress free.

(b) My teachers and school were very supportive as they taught all the concepts and covered the syllabus.

- **MTG : What is your family background?**

Kalpita : My father is a nurse in a government hospital. My mother is a teacher in government school and my brother is pursuing MBBS at AIIMS, Jodhpur.

- **MTG : How have MTG magazines helped you in your preparation?**

Kalpita : I was a 2 year subscriber of MTG magazines : Physics For You, Chemistry Today and Mathematics Today and was really an avid reader of them. These magazines were highly useful especially because of the good question bank and articles published in every issue.

- **MTG : Was this your first attempt?**

Kalpita : Yes, this was my first attempt.

- **MTG : What do you think is the secret of your success?**

Kalpita : Consistency, dedication and hardwork is the secret. Proper preparation requires you to practice questions on regular basis.

- **MTG : How did you de-stress yourself during the preparation? What are your hobbies? How often could you pursue them?**

Kalpita : I used to listen to music, play cricket and badminton to release the stress, in between my studies. I like listening to Coldplay and Linkin Park.

- **MTG : What do you feel is lacking in our education/ examination system? Is the examination system fair to the student?**

Kalpita : Yes, it is fair.

- **MTG : Had you not been selected then what would have been your future plan?**

Kalpita : I have not thought about it.

- **MTG : What advice would you like to give to our readers who are JEE aspirants?**

Kalpita : I advise to all aspirants that continuous hardwork along with dedication and time management would give the desired results.

All the Best! 😊😊

Your favourite MTG Books/Magazines available in **PUNJAB at**

- Sunder Book Depot - Amritsar
Ph: 0183-2544491, 2224884, 5059033; Mob: 9814074241
- Malhotra Book Depot - Amritsar Mob: 8000300086, 9646537157, 9888733844
- Navchattan Book Depot - Barnala
Mob: 97790220692, 9417906880, 9779090135, 9876147263, 9779050692
- Mehta Book Centre - Bathinda Mob: 9876029048, 9464158497
- Goyal Traders Book Sellers - Bathinda
Ph: 0164-2239632; Mob: 9417924911, 9814485520
- Aggarwal Book Centre - Bathinda Ph: 0164-2236042; Mob: 9417816439
- S M Enterprises - Bathinda Ph: 0164-2240450; Mob: 7508622881, 9417363362
- Hans Raj Dogra & Sons - Gurdaspur Mob: 9872032357
- Gupta Technical & Computer Book Shop - Jalandhar
Ph: 0181-2200397; Mob: 9915001916, 9779981081
- Cheap Book Store - Jalandhar Mob: 9872223458
- City Book Shop - Jalandhar Ph: 0181-2620800; Mob: 9417440753
- Deepak Book Distributors - Jalandhar
Ph: 0181-2222131; Mob: 8528391133, 9872328131
- Kiran Book Shop - Jalandhar Mob: 9876631526, 9779223883, 9872377808
- Amit Book Depot - Ludhiana
Ph: 0161-5022930, 5022930; Mob: 9815323429, 9815807871
- Bhatia Book Centre - Ludhiana Ph: 0161-2747713; Mob: 9815277131
- Chabra Book Depot - Ludhiana Ph: 0161-6900900, 2405427; Mob: 9501080070
- Gupta Book World - Ludhiana
Ph: 0161-2446870, 2409097, 3942992; Mob: 9463027555
- Khanna Book Depot - Nabha Ph: 01765-220095; Mob: 9814093193, 9814309320
- Goel & Sons Book Depot - Patiala Ph: 0172-2213643, 2202045; Mob: 9914096991
- Adarsh Pustak Bhandar - Patiala
Ph: 0175-2311430; Mob: 9814347613, 9815651737
- Pepsu Book Depot - Patiala Ph: 0175-2214696; Mob: 9814041623, 9914903305

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call **0124-6601200** for further assistance.

math on **GO**
the online math library



FREE ONLINE MATH CLASSES

Study through video lectures for
JEE Main & Adv | BITSAT | State Entrance Exams

YouTube WWW.YOUTUBE.COM/MATHONGO

f LIVE WWW.FACEBOOK.COM/MATHONGO123

FIND ALL THE VIDEO LECTURES FOR
YOUR REVISION

LIVE CHAT WITH ME FOR ALL YOUR TENSIONS
AND NECESSARY MOTIVATION



SHARE CONTENT AND CHAT WITH
YOUR FELLOW ASPIRANTS

REGISTER YOURSELF NOW

WWW.MATHONGO.COM

FOR ANY QUERY EMAIL AT

mathongochannel@gmail.com

YEARLONG CLASS 12TH COURSE

STARTING THIS SUMMER BREAK

Live Discussion and interaction after every 15 days on various topics like how to prepare, how to reduce anxiety, talks with toppers, how to succeed, which books to follow etc... Platform to openly ask your doubts and reduce your anxiety and tension and learn free.

**MATHONGO HAS NOW 8,00,000 +
MINUTES OF LEARNING CONSUMED.**

CALLING TEACHERS OF THE COUNTRY

IF YOU ARE A TEACHER OR WANNA BE TEACHER & WANT TO HELP IN THIS NOBLE CAUSE,
JUST DROP AN EMAIL. IT'S YOUR CHANCE TO BE A REAL SUPER PROF OUR COUNTRY NEEDS

KEEP SUPPORTING | KEEP LEARNING

I am a Computer Engineer from NSIT and an IIMA Alumni. I have been in the education for almost 8 years now and have produced hundreds of successful ranks at various competitive exams in the country - Highest being AIR 21 at IIT JEE. More than this, I have helped my students even after exams to crack companies such as Microsoft & Google. I hope to reach as many students as possible and help the community by providing high class affordable education.

Anup Gupta



ACE YOUR WAY CBSE



Relations and Functions

IMPORTANT FORMULAE

- ▶ If A and B be two non-empty sets, then cartesian product of sets is


$$A \times B = \{(x_i, y_i) : x_i \in A, y_i \in B\}$$
- ▶ $(x, y) = (p, q) \Leftrightarrow x = p, y = q$
- ▶ $A \times B = B \times A \Rightarrow A = B$
- ▶ If $n(A) = p, n(B) = q$, then $n(A \times B) = pq$
- ▶ $A \times A \times A = \{(a, b, c) : a, b, c \in A\}$ is called ordered triplet.
- ▶ $A \times B = \phi \Leftrightarrow A = \phi$ or $B = \phi$
- ▶ Subset of $X \times Y$ is called a relation from X to Y .
- ▶ If $n(X) = p$ and $n(Y) = q$ then the total number of relations from X to Y is 2^{pq} .
- ▶ The set of all first elements of the ordered pairs in relation R from set X to set Y is called the domain of the relation R .
- ▶ The set of all second elements of the ordered pairs in relation R from set X to Y is called the range of the relation R .
- ▶ The whole set B is called the co-domain of the relation R .
 - $\text{Range} \subseteq \text{Co-domain}$
- ▶ If $R = \{(a, b) : a, b \in R\}$, then

$$R^{-1} = \{(b, a) : b, a \in R\}$$
- ▶ A subset f of $X \times Y$ is called a function (or map or mapping) from X to Y iff for each $x \in X$, there exists a unique $y \in Y$ such that $(x, y) \in f$. It is written as $f: X \rightarrow Y$.
 - Set X is called domain and Set Y is called co-domain of the function f .
 - The set of elements of Y , which are assigned to the elements of X is called range of f .
- ▶ Algebra of real functions


If $f: X \rightarrow R$ and $g: X \rightarrow R$, then

 - $(f + g)(x) = f(x) + g(x), \forall x \in X$
 - $(f - g)(x) = f(x) - g(x), \forall x \in X$
 - $(\alpha f)(x) = \alpha f(x), \forall x \in X$
 - $(fg)(x) = f(x)g(x), \forall x \in X$
 - $\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}, g(x) \neq 0, x \in X$
- ▶ $\text{Function} \subseteq \text{Relation} \subseteq \text{Cartesian Product}$
- ▶ If A and B be two non-empty sets having n elements in common, then $A \times B$ and $B \times A$ have n^2 elements in common.


Concerned about your performance in **Class XII** Boards?




₹ 495



₹ 495



₹ 575



₹ 450

Well, fear no more, help is at hand.....

To excel, studying in right direction is more important than studying hard. Which is why we created the Excel Series. These books – for Physics, Chemistry, Biology & Mathematics – have been put together totally keeping in mind the prescribed syllabus and the pattern of CBSE's Board examinations, so that students prepare and practice with just the right study material to excel in board exams.

Did you know nearly all questions in CBSE's 2017 Board Examination were a part of our Excel books? That too fully solved !

HIGHLIGHTS:

- Comprehensive theory strictly based on NCERT, complemented with illustrations, activities and solutions of NCERT questions
- Practice questions & Model Test Papers for Board Exams
- Value based questions
- Previous years' CBSE Board Examination Papers (Solved)
- CBSE Board Papers 2017 Included



Scan now with your smartphone or tablet*

Visit
www.mtg.in
for latest offers
and to buy
online!



Available at all leading book shops throughout the country.

For more information or for help in placing your order:

Call 0124-6601200 or email: info@mtg.in

*Application to read QR codes required

WORK IT OUT**VERY SHORT ANSWER TYPE**

- Let $A = \{1, 2\}$ and $B = \{3, 4, 5\}$. Find
(i) $B \times A$ (ii) $A \times A \times A$
- Let $P = \{x, y, z\}$ and $Q = \{3, 4\}$. Find the number of relations from P to Q .
- Let f be the exponential function and g be the logarithmic function. Find $(fg)(1)$.
- Let $f(x) = x^2$ and $g(x) = (3x + 2)$ be two real functions. Then, find $(f + g)(x)$.
- Let $g = \{(1, 2), (2, 5), (3, 8), (4, 10), (5, 12), (6, 12)\}$. Is g a function? If yes, find its domain and range. If no, give reason.

SHORT ANSWER TYPE

- Let $f: Z \rightarrow Z, g: Z \rightarrow Z$ be functions defined by $f = \{n, n^2\} : n \in Z$ and $g = \{(n, |n|^2) : n \in Z\}$. Show that $f = g$.
- The function $F(x) = \frac{9x}{5} + 32$ is the formula to convert $x^\circ C$ to Fahrenheit units. Find
(i) $F(0)$ (ii) $F(-10)$
(iii) the value of x when $F(x) = 212$.
- If $f(x) = \log\left(\frac{1+x}{1-x}\right)$, prove that $f\left(\frac{2x}{1+x^2}\right) = 2f(x)$.
- Let R be a relation on the set of natural numbers N defined by $xRY \Leftrightarrow x + 2y = 41; \forall x, y \in N$. Find the domain and range of R .
- If $P = \{a, b\}$ and $Q = \{x, y, z\}$, then show that $P \times Q \neq Q \times P$.

LONG ANSWER TYPE - I

- Given $f(x) = \frac{1}{(1-x)}$, $g(x) = f\{f(x)\}$ and $h(x) = f\{f\{f(x)\}\}$. Then find the value of $f(x) \cdot g(x) \cdot h(x)$.
- Find the domain of the function
$$y = \frac{1}{\log_{10}(1-x)} + \sqrt{x+2}$$
- If $A \subseteq B$ and $C \subseteq D$, prove that $A \times C \subseteq B \times D$.
- Let f be a real valued function defined by
$$f(x) = \frac{e^x - e^{-|x|}}{e^x + e^{|x|}}$$
, then find the range of f .

- Consider the following :

(i) $f: R \rightarrow R : f(x) = \log_e x$

(ii) $g: R \rightarrow R : g(x) = \sqrt{x}$

(iii) $h: A \rightarrow R : h(x) = \frac{1}{x^2 - 4}$, where $A = R - \{-2, 2\}$

Which of them are functions? Also find their range, if they are functions.

LONG ANSWER TYPE - II

- Find the domain and range of the real valued function $f(x)$ given by $f(x) = \frac{4-x}{x-4}$.
- If $f: R \rightarrow R$ is defined by $f(x) = x^3 + 1$ and $g: R \rightarrow R$ is defined by $g(x) = x + 1$, then find
(i) $f + g$ (ii) $f - g$ (iii) $f \cdot g$
(iv) $\frac{f}{g}$ (v) $\alpha f (\alpha \in R)$
- Find the domain and range of the function
$$f(x) = \frac{1}{2 - \sin 3x}$$
- Let $A = \{x \in N : x^2 - 5x + 6 = 0\}$, $B = \{x \in Z : 0 \leq x < 2\}$ and $C = \{x \in N : x < 3\}$, then verify that:
(i) $A \times (B \cup C) = (A \times B) \cup (A \times C)$
(ii) $A \times (B \cap C) = (A \times B) \cap (A \times C)$
- Find the domain of the function
$$f(x) = \sqrt{(\log_2(x))} + \sqrt{7x - x^2 - 6}$$

SOLUTIONS

- (i) $B \times A = \{3, 4, 5\} \times \{1, 2\}$
 $= \{(3, 1), (3, 2), (4, 1), (4, 2), (5, 1), (5, 2)\}$
(ii) $A \times A \times A = \{1, 2\} \times \{1, 2\} \times \{1, 2\}$
 $= \{(1, 1, 1), (1, 1, 2), (1, 2, 1), (1, 2, 2), (2, 1, 1), (2, 1, 2), (2, 2, 1), (2, 2, 2)\}$
- Given, $P = \{x, y, z\}$ and $Q = \{3, 4\}$
 $\therefore n(P) = 3$ and $n(Q) = 2$
 $\therefore n(P \times Q) = 3 \cdot 2 = 6$
Total number of relations from P to Q = number of subsets of $P \times Q = 2^6 = 64$.
- We have, $f: R \rightarrow R$ given by $f(x) = e^x$ and $g: R^+ \rightarrow R$ given by $g(x) = \log_e x$
Domain $(f) \cap \text{Domain } (g) = R \cap R^+ = R^+$.
 $\therefore fg: R^+ \rightarrow R$ is given by $(fg)(x) = f(x)g(x) = e^x \cdot \log_e x$.
Now, $(fg)(1) = f(1)g(1) = e^1 \times \log_e 1 = e \times 0 = 0$.
- We have $(f + g)(x) = f(x) + g(x) = x^2 + (3x + 2)$
- Yes, $\text{dom } (g) = \{1, 2, 3, 4, 5, 6\}$,
 $\text{range } (g) = \{2, 5, 8, 10, 12\}$

6. Given, domain of $f = Z$ and domain of $g = Z$

Hence, domain $f = \text{domain } g = Z$

Also, $f(n) = n^2$, for all $n \in Z$ and $g(n) = |n|^2 = n^2$ for all $n \in Z$

Hence, $f(n) = g(n)$ for all $n \in Z$

From (1) and (2), we have $f = g$.

7. $F(x) = \frac{9x}{5} + 32$ (given)

$$(i) F(0) = \left(\frac{9 \times 0}{5} + 32 \right) = 32 \Rightarrow F(0) = 32$$

$$(ii) F(-10) = \left\{ \frac{9 \times (-10)}{5} + 32 \right\} = 14 \Rightarrow F(-10) = 14$$

$$(iii) F(x) = 212 \Leftrightarrow \frac{9x}{5} + 32 = 212$$

$$\Leftrightarrow 9x = (5 \times 180) \Leftrightarrow x = 100$$

8. Given, $f(x) = \log \left(\frac{1+x}{1-x} \right)$

$$\begin{aligned} \therefore f \left(\frac{2x}{1+x^2} \right) &= \log \left(\frac{1 + \frac{2x}{1+x^2}}{1 - \frac{2x}{1+x^2}} \right) = \log \left(\frac{1+x^2+2x}{1+x^2-2x} \right) \\ &= \log \left(\frac{1+x}{1-x} \right)^2 = 2 \log \left(\frac{1+x}{1-x} \right) = 2f(x). \end{aligned}$$

9. We have, $y = \frac{41-x}{2} \in N$

Clearly $x = 1, 3, 5, 7, \dots, 39$

$$\therefore \text{Domain } R = \{x : (x, y) \in R; x + 2y = 41\}$$

$$= \{1, 3, 5, 7, \dots, 39\}$$

= set of odd natural numbers less than 40.

Now, y can be only those natural numbers for which $x \in N$ i.e., $x = 41 - 2y \in N$.

Clearly, $y = 1, 2, 3, \dots, 20$.

$$\therefore \text{Range of } R = \{y : x + 2y = 41\} = \{1, 2, 3, \dots, 20\}$$

= set of natural numbers less than 21.

10. We have, $P = \{a, b\}$ and $Q = \{x, y, z\}$

Now, $P \times Q = \{(a, x), (a, y), (a, z), (b, x), (b, y), (b, z)\}$

$Q \times P = \{(x, a), (x, b), (y, a), (y, b), (z, a), (z, b)\}$

$\therefore P \times Q \neq Q \times P$.

11. Given, $g(x) = f\{f(x)\}$

$$= f \left(\frac{1}{1-x} \right) = \frac{1}{1 - \frac{1}{1-x}} = - \left(\frac{1-x}{x} \right)$$

$$\text{and } h(x) = f[f\{f(x)\}] = f\{g(x)\} = f \left(- \left(\frac{1-x}{x} \right) \right)$$

$$= \frac{1}{1 - \frac{1}{1-x}} = x$$

$$\therefore f(x) \cdot g(x) \cdot h(x) = \frac{1}{1-x} \left\{ - \left(\frac{1-x}{x} \right) \right\} \cdot x = -1$$

12. For y to be defined

(i) $\log_{10}(1-x)$ must be defined $\Rightarrow 1-x > 0 \Rightarrow x < 1$

(ii) $\log_{10}(1-x) \neq 0 \Rightarrow 1-x \neq 10^0 \Rightarrow 1-x \neq 1 \Rightarrow x \neq 0$

(iii) $x+2 \geq 0 \Rightarrow x \geq -2$

From (i), (ii) and (iii), we get $-2 \leq x < 1$ and $x \neq 0$

$\therefore -2 \leq x < 0$ or $0 < x < 1$

Hence domain = $[-2, 0) \cup (0, 1)$.

13. Let (a, b) be an arbitrary element of $A \times C$. Then,

$(a, b) \in A \times C$

$\Rightarrow a \in A$ and $b \in C$

$\Rightarrow a \in B$ and $b \in D$ [$\because A \subseteq B$ and $C \subseteq D$]

$\Rightarrow (a, b) \in B \times D$

Thus, $(a, b) \in A \times C$

$\Rightarrow (a, b) \in B \times D$ for all $(a, b) \in (A \times C)$.

$\therefore A \times C \subseteq B \times D$

$$14. \text{ Let } y = f(x) = \frac{e^x - e^{-|x|}}{e^x + e^{|x|}}$$

$$\text{If } x \geq 0 \text{ then } y = \frac{e^{2x} - 1}{2e^{2x}}$$

$$\Rightarrow e^{2x} = \frac{1}{1-2y} \geq 1 \quad (\because x \geq 0)$$

$$\Rightarrow \frac{1}{1-2y} - 1 \geq 0 \Rightarrow \frac{y}{1-2y} \geq 0 \quad \text{or} \quad \frac{y}{2y-1} \leq 0$$

$$\therefore 0 \leq y < \frac{1}{2} \Rightarrow y \in \left[0, \frac{1}{2} \right)$$

15. f and g are not functions as they are not defined for negative values of x . But h is a function.

$$\therefore \text{For range of } h, \text{ Let } y = h(x) = \frac{1}{x^2 - 4}$$

$$\Rightarrow x^2 - 4 = \frac{1}{y} \Rightarrow x^2 = 4 + \frac{1}{y} \Rightarrow x = \sqrt{\frac{4y+1}{y}}$$

$$\text{Hence, range of } h = \left(-\infty, -\frac{1}{4} \right] \cup (0, \infty)$$

MPP-2 CLASS XII

ANSWER KEY

1. (c) 2. (b) 3. (b) 4. (d) 5. (c)
6. (c) 7. (a,b) 8. (b,c) 9. (b,c) 10. (b)
11. (a,b) 12. (a) 13. (a,b,d) 14. (d) 15. (b)
16. (d) 17. (2) 18. (3) 19. (2) 20. (7)

16. We have, $f(x) = \frac{4-x}{x-4}$.

Domain of f : We observe that $f(x)$ is defined for all x except at $x = 4$. At $x = 4$, $f(x)$ takes the indeterminate form $\frac{0}{0}$. Therefore, Domain (f) = $R - \{4\}$.

Range of f : For any $x \in \text{Domain}(f)$ i.e. for any $x \neq 4$, we have

$$f(x) = \frac{4-x}{x-4} = \frac{-(x-4)}{x-4} = -1.$$

$$\therefore \text{Range}(f) = \{-1\}.$$

17. (i) $f+g: R \rightarrow R$ is defined by

$$(f+g)(x) = f(x) + g(x) = x^3 + 1 + x + 1 = x^3 + x + 2$$

(ii) $f-g: R \rightarrow R$ is defined by

$$(f-g)(x) = f(x) - g(x) = x^3 + 1 - x - 1 = x^3 - x$$

(iii) $f \cdot g: R \rightarrow R$ is defined by

$$(fg)(x) = f(x)g(x) = (x^3 + 1)(x + 1) = x^4 + x^3 + x + 1$$

(iv) $\frac{f}{g}: R - \{-1\} \rightarrow R$ is defined by

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} = \frac{x^3 + 1}{x + 1} = \frac{(x+1)(x^2 - x + 1)}{x + 1} = x^2 - x + 1$$

(v) $\alpha f: R \rightarrow R$ is defined by

$$(\alpha f)(x) = \alpha f(x) = \alpha(x^3 + 1) = \alpha x^3 + \alpha$$

18. We have, $f(x) = \frac{1}{2 - \sin 3x}$

Domain of f : We know that $-1 \leq \sin 3x \leq 1$ for all $x \in R$

$$\Rightarrow -1 \leq -\sin 3x \leq 1 \text{ for all } x \in R$$

$$\Rightarrow 1 \leq 2 - \sin 3x \leq 3 \text{ for all } x \in R$$

$$\Rightarrow 2 - \sin 3x \neq 0 \text{ for any } x \in R$$

$$\Rightarrow f(x) = \frac{1}{2 - \sin 3x} \text{ is defined for all } x \in R$$

Hence, domain (f) = R .

Range of f : $\because 1 \leq 2 - \sin 3x \leq 3$ for all $x \in R$

$$\Rightarrow \frac{1}{3} \leq \frac{1}{2 - \sin 3x} \leq 1 \text{ for all } x \in R$$

$$\Rightarrow \frac{1}{3} \leq f(x) \leq 1 \text{ for all } x \in R$$

$$\Rightarrow f(x) \in [1/3, 1]$$

Hence, range (f) = $[1/3, 1]$

19. We have

$$A = \{x \in N : x^2 - 5x + 6 = 0\} = \{2, 3\};$$

$$B = \{x \in Z : 0 \leq x < 2\} = \{0, 1\} \text{ and}$$

$$C = \{x \in N : x < 3\} = \{1, 2\}$$

$$\therefore A = \{2, 3\}, B = \{0, 1\} \text{ and } C = \{1, 2\}$$

$$(i) (B \cup C) = \{0, 1, 2\}$$

$$\therefore A \times (B \cup C) = \{2, 3\} \times \{0, 1, 2\}$$

$$= \{(2, 0), (2, 1), (2, 2), (3, 0), (3, 1), (3, 2)\}$$

$$(A \times B) = \{2, 3\} \times \{0, 1\} = \{(2, 0), (2, 1), (3, 0), (3, 1)\}$$

$$(A \times C) = \{2, 3\} \times \{1, 2\} = \{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$\therefore (A \times B) \cup (A \times C) = \{(2, 0), (2, 1), (2, 2), (3, 0), (3, 1), (3, 2)\}$$

$$\text{Hence, } A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$(ii) (B \cap C) = \{0, 1\} = \{1\}$$

$$\therefore A \times (B \cap C) = \{2, 3\} \times \{1\} = \{(2, 1), (3, 1)\}$$

And,

$$(A \times B) \cap (A \times C) = \{(2, 1), (3, 1)\}$$

$$\text{Hence, } A \times (B \cap C) = (A \times B) \cap (A \times C)$$

20. We have, $f(x) = \sqrt{(\log_2(x))} + \sqrt{7x - x^2 - 6}$

$$= \sqrt{(\log_2(x))} + \sqrt{(1-x)(x-6)}$$

For $f(x)$ to be defined.

$$(i) (\log_2(x)) \geq 0 \Rightarrow x \geq 2^0 \Rightarrow x \geq 1$$

$$(ii) (1-x)(x-6) \geq 0 \Rightarrow 1 \leq x \leq 6$$

Therefore domain of $f = [1, 6]$.



Your favourite MTG Books/Magazines available in RAJASTHAN at

- Competition Book House - Alwar Ph: 0144-2338391; Mob: 9460607836
- Nakoda Book Depot - Bhilwara Ph: 01482-239653; Mob: 9214983594
- Alankar Book Depot - Bhiwadi Ph: 01493-222294; Mob: 9414707462
- Uttam Pustak Kendra - Bikaner Mob: 8955543195, 9414572625
- Yadu Books & Stationers - Bikaner Mob: 9251653481
- Goyal Books & Stationers - Jaipur
Ph: 0141-2742454; Mob: 9414326406, 9929638435
- India Book House - Jaipur
Ph: 0141-2314983, 2311191, 2651784; Mob: 9829014143, 9414079983
- Ravi Enterprises - Jaipur
Ph: 0141-2602517, 2619958, 2606998; Mob: 9829060694
- Shri Shyam Pustak Mandir - Jaipur Ph: 0141-2317972; Mob: 9928450717
- Sarvodaya Book Stall - Jodhpur Ph: 0291-2653734, 35; Mob: 8107589141
- Bhandari Stationers - Kota
Ph: 0744-2327576, 2391678; Mob: 9001094271, 9829038758
- Raj Traders - Kota Ph: 0744-2429090; Mob: 9309232829, 9214335300
- Vardhman Sports & Stationers - Kota
Mob: 9461051901, 9351581238, 9828139717
- Jhuria Book Shop - Sikar Mob: 9784246419, 9460838235, 8432943550
- Popular Book Depot - Udaipur
Ph: 2442881, 0487-2329436, 2421467; Mob: 9388513335, 9847922545

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call **0124-6601200** for further assistance.

The only thing you NEED for excellence in Class -12 Boards



CBSE CHAMPION Chapterwise -Topicwise Solved Papers Series

CBSE CHAMPION Chapterwise-Topicwise Solved Papers series contains topicwise questions and solutions asked over last 11 Years in CBSE-Board Class-12 examination. Questions are supported with topicwise graphical analysis of previous 11 years CBSE Board questions as well as comprehensive and lucid theory. The questions in each topic have been arranged in descending order (2017-2007) as per their marking scheme. Questions from Delhi, All India, Foreign and Compartment papers are included. This ensures that all types of questions that are necessary for Board exam preparation have been covered.

Important feature of these books is that the solutions to all the questions have been given according to CBSE marking scheme. Latest CBSE sample paper and practice papers are also supplemented.

Examination papers for Class-12 Boards are based on a certain pattern. To excel, studying right is therefore more important than studying hard, which is why we created this series.

MPP-2 MONTHLY Practice Problems

Class XI

This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.



Complex Numbers and Quadratic Equations

Total Marks : 80

Time Taken : 60 Min.

Only One Option Correct Type

- $$\left(\frac{1 + i \sin \frac{\pi}{8} + \cos \frac{\pi}{8}}{1 - i \sin \frac{\pi}{8} + \cos \frac{\pi}{8}} \right)^8$$
 equals
 - 2^8
 - 0
 - 1
 - 1
- If a, b, c are in G.P., then the equations $ax^2 + 2bx + c = 0$, $dx^2 + 2ex + f = 0$ have a common root if $\frac{d}{a}, \frac{e}{b}, \frac{f}{c}$ are in
 - A.P.
 - G.P.
 - H.P.
 - none of these
- For positive integers n_1 and n_2 the value of the expression $(1+i)^{n_1} + (1+i^3)^{n_1} + (1+i^5)^{n_2} + (1+i^7)^{n_2}$ where $i = \sqrt{-1}$ is a real number iff
 - $n_1 = n_2$
 - $n_2 = n_1 - 1$
 - $n_1 = n_2 + 1$
 - $\forall n_1$ and n_2
- If x, y, z are distinct positive reals such that $\frac{\log x}{y-z} = \frac{\log y}{z-x} = \frac{\log z}{x-y}$, then value of $x^x y^y z^z$ is
 - 1
 - 0
 - 1
 - none of these
- If the roots of the equation $bx^2 + cx + a = 0$ be imaginary, then for all real values of x , the expression $3b^2x^2 + 6bcx + 2c^2$ is
 - less than $4ab$
 - greater than $-4ab$
 - less than $-4ab$
 - greater than $4ab$

- Solve for x : $\log_{2x+3} (6x^2 + 23x + 21) + \log_{3x+7} (4x^2 + 12x + 9) = 4$
 - 4
 - 2
 - $-\frac{1}{4}$
 - All of these

One or More Than One Option(s) Correct Type

- ABCD is a rhombus, its diagonals AC and BD intersect at the point R where $BD = 2AC$. Its points D and R represent the complex numbers $1 + i$ and $2 - i$ respectively, then the complex number represented by A is
 - $(3, -1/2)$ or $(1, -1/2)$
 - $(3, -1/2)$ or $(1, -3/2)$
 - $(-1/2, -3/2)$ or $(-3/2, -1/2)$
 - None of these
- $(1+i)^5 + (1-i)^5 =$
 - 8
 - 8
 - $2^{7/2} \cos \frac{5\pi}{4}$
 - $-2^{7/2} \cos \frac{5\pi}{4}$
- If α and β are non-real cube roots of unity and $x = a + b, y = a\alpha + b\beta, z = a\beta + b\alpha$, then
 - $x + y + z = 1$
 - $x + y + z = 0$
 - $x^3 + y^3 + z^3 = 3(a^3 + b^3)$
 - none of these
- The equation whose roots are $\alpha\gamma + \beta\delta$ and $\alpha\delta + \gamma\beta$, if α, β are the roots of the equation $ax^2 + bx + c = 0$ and γ, δ are roots of the equation $a'x^2 + b'x + c' = 0$, is

- (a) $aa'x^2 + bb'x + cc' = 0$
 (b) $ax^2 + (a + a' + b')x + bc' = 0$
 (c) $aa'x^2 - bb'x + cc' = 0$
 (d) none of these

11. If A and B are the points (3, -1) and (2, 1) respectively, then the locus of the points $P(z)$, $z = x + yi$, $x, y \in R$, such that $|z - 3 + i| = |z - 2 - i|$ is
 (a) a circle containing A and B
 (b) P is equidistant from A and B
 (c) right bisector of segment joining A and B
 (d) none of these

12. The value of x : $|x^2 + 2x - 8| + x - 2 = 0$ is
 (a) -5 (b) -2
 (c) 2 (d) -3

13. If $a \in C$ be such that $|a| = 1$, then the equation

$$\left(\frac{1+iz}{1-iz}\right)^4 = a \text{ has all the roots}$$

- (a) real and distinct
 (b) non-real
 (c) two real and two non-real
 (d) none of these

Comprehension Type

Let α, β be the roots of the equation $6x^2 + 6px + p^2 = 0$, where p is a real number.

14. If both α and β are greater than 2, then
 (a) $p < -4$
 (b) $p < -2\sqrt{6}$
 (c) $p < -6 - 2\sqrt{6}$
 (d) none of these
15. The equation whose roots are $(\alpha + \beta)^2$ and $(\alpha - \beta)^2$ is
 (a) $3x^2 + 4p^2x + p^4 = 0$
 (b) $3x^2 - 4p^2x + p^4 = 0$
 (c) $3x^2 - 4p^2x - p^4 = 0$
 (d) none of these

Matrix Match Type

16. Match the following :

Column I		Column II	
P.	If a and b are positive numbers and $\log \frac{a+b}{2} = \frac{1}{2}(\log a + \log b)$ then $\frac{a}{b}$ is equal to	1.	$3\sqrt{3}$
Q.	Let $A(2 + 0i)$, $B(-1 + \sqrt{3}i)$ and $C(-1 - \sqrt{3}i)$ be the vertices of $\triangle ABC$. Then, $6 \sin A$ is equal to	2.	1
R.	If one root of the equation $(x - 1)(7 - x) = \lambda$ is three times the other, then $\lambda =$	3.	5
S.	Conjugate of the complex number $-\frac{7}{2} - \frac{3\sqrt{3}i}{2}$ is	4.	$-2 + 3\omega$

	P	Q	R	S
(a)	4	2	3	1
(b)	3	4	1	2
(c)	1	4	3	2
(d)	2	1	3	4

Integer Answer Type

17. The number of values of x in the interval $[0, 3\pi]$ satisfying the equation $2\sin^2 x + 5\sin x - 3 = 0$ is
18. The least integral value of k for which $(k - 2)x^2 + 8x + k + 4 > 0$ for all $x \in R$, is
19. If α, β are non-real cube roots of unity then $(1 + \alpha)(1 + \beta)(1 + \alpha^2)(1 + \beta^2)(1 + \alpha^4)(1 + \beta^4) \dots$ upto $2n$ factors is equal to
20. If α be non real cube root of unity, then $|\sqrt{\alpha}|$ equals



Keys are published in this issue. Search now! ☺

SELF CHECK

No. of questions attempted
 No. of questions correct
 Marks scored in percentage

Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY !	Revise thoroughly and strengthen your concepts.

ACE YOUR WAY CBSE

Inverse Trigonometric Functions



IMPORTANT FORMULAE

Functions	Domain	Range	Functions	Domain	Range
$\sin^{-1}x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	$\cot^{-1}x$	$(-\infty, \infty)$	$(0, \pi)$
$\cos^{-1}x$	$[-1, 1]$	$[0, \pi]$	$\operatorname{cosec}^{-1}x$	$(-\infty, -1] \cup [1, \infty)$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
$\tan^{-1}x$	$(-\infty, \infty)$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$	$\sec^{-1}x$	$(-\infty, -1] \cup [1, \infty)$	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$

PROPERTIES OF INVERSE TRIGONOMETRIC FUNCTIONS

1.	<ul style="list-style-type: none"> $\sin^{-1}(\sin x) = x, \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ $\cos^{-1}(\cos x) = x, \forall x \in [0, \pi]$ $\tan^{-1}(\tan x) = x, \forall x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ $\cot^{-1}(\cot x) = x, \forall x \in (0, \pi)$ $\sec^{-1}(\sec x) = x, \forall x \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$ $\operatorname{cosec}^{-1}(\operatorname{cosec} x) = x, \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$ 	2.	<ul style="list-style-type: none"> $\sin(\sin^{-1}x) = x, \forall x \in [-1, 1]$ $\cos(\cos^{-1}x) = x, \forall x \in [-1, 1]$ $\tan(\tan^{-1}x) = x, \forall x \in R$ $\cot(\cot^{-1}x) = x, \forall x \in R$ $\sec(\sec^{-1}x) = x, \forall x \in R - (-1, 1)$ $\operatorname{cosec}(\operatorname{cosec}^{-1}x) = x, \forall x \in R - (-1, 1)$
3.	<ul style="list-style-type: none"> $\sin^{-1}(-x) = -\sin^{-1}x, \forall x \in [-1, 1]$ $\cos^{-1}(-x) = \pi - \cos^{-1}x, \forall x \in [-1, 1]$ $\tan^{-1}(-x) = -\tan^{-1}x, \forall x \in R$ 		<ul style="list-style-type: none"> $\cot^{-1}(-x) = \pi - \cot^{-1}x, \forall x \in R$ $\sec^{-1}(-x) = \pi - \sec^{-1}x, \forall x \in (-\infty, -1] \cup [1, \infty)$ $\operatorname{cosec}^{-1}(-x) = -\operatorname{cosec}^{-1}x, \forall x \in (-\infty, -1] \cup [1, \infty)$

4.	<ul style="list-style-type: none"> $\sin^{-1}(1/x) = \operatorname{cosec}^{-1}x, \forall x \in (-\infty, -1] \cup [1, \infty)$ $\cos^{-1}(1/x) = \sec^{-1}x, \forall x \in (-\infty, -1] \cup [1, \infty)$ $\tan^{-1}(1/x) = \begin{cases} \cot^{-1}x & , \text{ for } x > 0 \\ -\pi + \cot^{-1}x & , \text{ for } x < 0 \end{cases}$ 	5.	<ul style="list-style-type: none"> $\sin^{-1}x + \cos^{-1}x = \pi/2, \forall x \in [-1, 1]$ $\tan^{-1}x + \cot^{-1}x = \pi/2, \forall x \in \mathbb{R}$ $\sec^{-1}x + \operatorname{cosec}^{-1}x = \pi/2, \forall x \in (-\infty, -1] \cup [1, \infty)$
6.	<ul style="list-style-type: none"> $\tan^{-1}x + \tan^{-1}y = \begin{cases} \tan^{-1}\left(\frac{x+y}{1-xy}\right) & , \text{ if } x > 0, y > 0, xy < 1 \\ \pi + \tan^{-1}\left(\frac{x+y}{1-xy}\right) & , \text{ if } x > 0, y > 0 \text{ and } xy > 1 \\ -\pi + \tan^{-1}\left(\frac{x+y}{1-xy}\right) & , \text{ if } x < 0, y < 0 \text{ and } xy > 1 \end{cases}$ $\tan^{-1}x - \tan^{-1}y = \begin{cases} \tan^{-1}\left(\frac{x-y}{1+xy}\right) & , \text{ if } x > 0, y < 0, xy > -1 \\ \pi + \tan^{-1}\left(\frac{x-y}{1+xy}\right) & , \text{ if } x > 0, y < 0 \text{ and } xy < -1 \\ -\pi + \tan^{-1}\left(\frac{x-y}{1+xy}\right) & , \text{ if } x < 0, y > 0 \text{ and } xy < -1 \end{cases}$ 		
7.	<ul style="list-style-type: none"> $\sin^{-1}x + \sin^{-1}y = \begin{cases} \sin^{-1}\{x\sqrt{1-y^2} + y\sqrt{1-x^2}\} & , \text{ if } -1 \leq x, y \leq 1 \text{ and } x^2 + y^2 \leq 1 \\ \text{or} \\ \pi - \sin^{-1}\{x\sqrt{1-y^2} + y\sqrt{1-x^2}\} & , \text{ if } 0 < x, y \leq 1 \text{ and } x^2 + y^2 > 1 \\ -\pi - \sin^{-1}\{x\sqrt{1-y^2} + y\sqrt{1-x^2}\} & , \text{ if } -1 \leq x, y < 0 \text{ and } x^2 + y^2 \geq 1 \end{cases}$ $\sin^{-1}x - \sin^{-1}y = \begin{cases} \sin^{-1}\{x\sqrt{1-y^2} - y\sqrt{1-x^2}\} & , \text{ if } -1 \leq x, y \leq 1 \text{ and } x^2 + y^2 \leq 1 \\ \text{or} \\ \pi - \sin^{-1}\{x\sqrt{1-y^2} - y\sqrt{1-x^2}\} & , \text{ if } 0 < x \leq 1, -1 \leq y \leq 0 \text{ and } x^2 + y^2 > 1 \\ -\pi - \sin^{-1}\{x\sqrt{1-y^2} - y\sqrt{1-x^2}\} & , \text{ if } -1 \leq x < 0, 0 < y \leq 1 \text{ and } x^2 + y^2 > 1 \end{cases}$ 		
8.	<ul style="list-style-type: none"> $\cos^{-1}x + \cos^{-1}y = \begin{cases} \cos^{-1}\{xy - \sqrt{1-x^2}\sqrt{1-y^2}\} & , \text{ if } -1 \leq x, y \leq 1 \text{ and } x+y \geq 0 \\ 2\pi - \cos^{-1}\{xy - \sqrt{1-x^2}\sqrt{1-y^2}\} & , \text{ if } -1 \leq x, y \leq 1 \text{ and } x+y \leq 0 \end{cases}$ $\cos^{-1}x - \cos^{-1}y = \begin{cases} \cos^{-1}\{xy + \sqrt{1-x^2}\sqrt{1-y^2}\} & , \text{ if } -1 \leq x, y \leq 1 \text{ and } x \leq y \\ -\cos^{-1}\{xy + \sqrt{1-x^2}\sqrt{1-y^2}\} & , \text{ if } -1 \leq y \leq 0, 0 < x \leq 1 \text{ and } x \geq y \end{cases}$ 		

9.	$2 \sin^{-1} x = \begin{cases} \sin^{-1}(2x\sqrt{1-x^2}) & , \text{ if } -\frac{1}{\sqrt{2}} \leq x \leq \frac{1}{\sqrt{2}} \\ \pi - \sin^{-1}(2x\sqrt{1-x^2}) & , \text{ if } \frac{1}{\sqrt{2}} \leq x \leq 1 \\ -\pi - \sin^{-1}(2x\sqrt{1-x^2}) & , \text{ if } -1 \leq x \leq -\frac{1}{\sqrt{2}} \end{cases}$	
	$3 \sin^{-1} x = \begin{cases} \sin^{-1}(3x-4x^3) & , \text{ if } -\frac{1}{2} \leq x \leq \frac{1}{2} \\ \pi - \sin^{-1}(3x-4x^3) & , \text{ if } \frac{1}{2} < x \leq 1 \\ -\pi - \sin^{-1}(3x-4x^3) & , \text{ if } -1 \leq x < -\frac{1}{2} \end{cases}$	
10.	$2 \cos^{-1} x = \begin{cases} \cos^{-1}(2x^2-1), & \text{if } 0 \leq x \leq 1 \\ 2\pi - \cos^{-1}(2x^2-1), & \text{if } -1 \leq x \leq 0 \end{cases}$	$3 \cos^{-1} x = \begin{cases} \cos^{-1}(4x^3-3x) & , \text{ if } \frac{1}{2} \leq x \leq 1 \\ 2\pi - \cos^{-1}(4x^3-3x), & \text{if } -\frac{1}{2} \leq x \leq \frac{1}{2} \\ 2\pi + \cos^{-1}(4x^3-3x), & \text{if } -1 \leq x \leq -\frac{1}{2} \end{cases}$
11.	$2 \tan^{-1} x = \begin{cases} \tan^{-1}\left(\frac{2x}{1-x^2}\right), & \text{if } -1 < x < 1 \\ \pi + \tan^{-1}\left(\frac{2x}{1-x^2}\right), & \text{if } x > 1 \\ -\pi + \tan^{-1}\left(\frac{2x}{1-x^2}\right), & \text{if } x < -1 \end{cases}$	$3 \tan^{-1} x = \begin{cases} \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right) & , \text{ if } -\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}} \\ \pi + \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right), & \text{if } x > \frac{1}{\sqrt{3}} \\ -\pi + \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right), & \text{if } x < -\frac{1}{\sqrt{3}} \end{cases}$
12.	$2 \tan^{-1} x = \begin{cases} \sin^{-1}\left(\frac{2x}{1+x^2}\right), & \text{if } -1 \leq x \leq 1 \\ \pi - \sin^{-1}\left(\frac{2x}{1+x^2}\right), & \text{if } x > 1 \\ -\pi - \sin^{-1}\left(\frac{2x}{1+x^2}\right), & \text{if } x < -1 \end{cases}$	$2 \tan^{-1} x = \begin{cases} \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right), & \text{if } 0 \leq x < \infty \\ -\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right), & \text{if } -\infty < x \leq 0 \end{cases}$
13.	$\sin^{-1} x = \cos^{-1}(\sqrt{1-x^2}) = \tan^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right) = \cot^{-1}\left(\frac{\sqrt{1-x^2}}{x}\right) = \sec^{-1}\left(\frac{1}{\sqrt{1-x^2}}\right) = \operatorname{cosec}^{-1}\left(\frac{1}{x}\right)$	
	$\cos^{-1} x = \sin^{-1}(\sqrt{1-x^2}) = \tan^{-1}\left(\frac{\sqrt{1-x^2}}{x}\right) = \cot^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right) = \sec^{-1}\left(\frac{1}{x}\right) = \operatorname{cosec}^{-1}\left(\frac{1}{\sqrt{1-x^2}}\right)$	
	$\tan^{-1} x = \sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right) = \cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right) = \cot^{-1}\left(\frac{1}{x}\right) = \sec^{-1}(\sqrt{1+x^2}) = \operatorname{cosec}^{-1}\left(\frac{\sqrt{1+x^2}}{x}\right)$	
14.	<p>If $x_1, x_2, \dots, x_n \in R$, then $\tan^{-1} x_1 + \tan^{-1} x_2 + \dots + \tan^{-1} x_n = \tan^{-1} \left(\frac{S_1 - S_3 + S_5 - S_7 + \dots}{1 - S_2 + S_4 - S_6 + \dots} \right)$ where S_k = Sum of the products of x_1, x_2, \dots, x_n taken k at a time.</p>	

WORK IT OUT**VERY SHORT ANSWER TYPE**

- Find the value of $\cot\left(\frac{\pi}{4} - 2 \cot^{-1} 3\right)$.
- If $\tan^{-1} \frac{4}{3} = \theta$, find the value of $\cos \theta$.
- Find the principal value of $\sin^{-1}\left(\sin \frac{2\pi}{3}\right)$.
- For the principal values, evaluate the following:
 $\cot^{-1}(-\sqrt{3}) + \tan^{-1}(1) + \sec^{-1}\left(\frac{2}{\sqrt{3}}\right)$
- Evaluate : $\cos^{-1}(\cos(-680^\circ))$

SHORT ANSWER TYPE

- Evaluate : (i) $\sin(\cot^{-1} x)$ (ii) $\cos(\tan^{-1} x)$
- Prove that $\sin^{-1} \frac{8}{17} + \sin^{-1} \frac{3}{5} = \cos^{-1} \frac{36}{85}$.
- Solve for x : $\cos(\tan^{-1} x) = \sin\left(\sec^{-1} \frac{13}{12}\right)$.
- Solve : $\sin^{-1} x = \frac{\pi}{6} + \cos^{-1} x$
- If in a $\triangle ABC$, $\angle A = \tan^{-1} 2$ and $\angle B = \tan^{-1} 3$, then show that $\angle C$ is equal to $\frac{\pi}{4}$.

LONG ANSWER TYPE-I

- Write each of the following in the simplest form:
 (i) $\tan^{-1}(\sec x + \tan x)$ (ii) $\sin\left(2 \tan^{-1} \sqrt{\frac{1-x}{1+x}}\right)$
- Show that $2 \tan^{-1}\left(\frac{1+x}{1-x}\right) + \sin^{-1}\left(\frac{1-x^2}{1+x^2}\right) = \pi$
- Evaluate the following :
 (i) $\sin(2 \sin^{-1} 0.8)$ (ii) $\tan\left[2 \tan^{-1}\left(\frac{1}{5}\right) - \frac{\pi}{4}\right]$
- Find the value of
 $\tan^{-1}\left(\tan \frac{5\pi}{6}\right) + \cos^{-1}\left(\cos \frac{13\pi}{6}\right) + \sec^{-1}\left(\sec \frac{9\pi}{5}\right)$
- If $\tan^{-1} \left\{ \frac{\sqrt{1+x^2} - \sqrt{1-x^2}}{\sqrt{1+x^2} + \sqrt{1-x^2}} \right\} = \alpha$, then prove that
 $x^2 = \sin 2\alpha$.

LONG ANSWER TYPE-II

- If $a, b, c > 0$ such that $a + b + c = abc$, find the value of $\tan^{-1} a + \tan^{-1} b + \tan^{-1} c$.
- If $\cos^{-1} \frac{x}{a} + \cos^{-1} \frac{y}{b} = \alpha$, prove that

$$\frac{x^2}{a^2} - \frac{2xy}{ab} \cos \alpha + \frac{y^2}{b^2} = \sin^2 \alpha$$
- If $a_1, a_2, a_3, \dots, a_n$ is an arithmetic progression with common difference d , then show that

$$\tan\left[\tan^{-1}\left(\frac{d}{1+a_1a_2}\right) + \tan^{-1}\left(\frac{d}{1+a_2a_3}\right) + \tan^{-1}\left(\frac{d}{1+a_3a_4}\right) + \dots + \tan^{-1}\left(\frac{d}{1+a_{n-1}a_n}\right)\right]$$
 is equal to $\frac{a_n - a_1}{1 + a_1a_n}$.
- Solve :
 (i) $\cos\left(\sin^{-1} \frac{1}{3} + \cos^{-1} x\right) = 0$
 (ii) $\tan^{-1}\left(\frac{x+1}{x-1}\right) + \tan^{-1}\left(\frac{x-1}{x}\right) = \pi + \tan^{-1}(-7)$
- Prove that
 (i) $\sin^{-1} \frac{12}{13} + \cos^{-1} \frac{4}{5} + \tan^{-1} \frac{63}{16} = \pi$
 (ii) $\cos^{-1} \frac{4}{5} + \tan^{-1} \frac{3}{5} = \tan^{-1} \frac{27}{11}$

SOLUTIONS

- $$\cot\left(\frac{\pi}{4} - 2 \cot^{-1} 3\right) = \cot\left[\frac{\pi}{4} - \cot^{-1}\left(\frac{3^2 - 1}{2 \times 3}\right)\right]$$

$$= \cot\left(\frac{\pi}{4} - \cot^{-1} \frac{4}{3}\right) = \frac{\cot \frac{\pi}{4} \cdot \frac{4}{3} + 1}{\frac{4}{3} - \cot \frac{\pi}{4}} = \frac{\frac{4}{3} + 1}{\frac{4}{3} - 1} = 7$$
- Given, $\tan^{-1} \frac{4}{3} = \theta$, where $\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
 $\therefore \tan \theta = \frac{4}{3}$
 We know that $\cos \theta > 0$, when $\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.

$$\therefore \cos \theta = \frac{1}{\sec \theta} = \frac{1}{\sqrt{1 + \tan^2 \theta}} = \frac{1}{\sqrt{1 + \frac{16}{9}}} = \frac{3}{5}$$

3. Since, $\sin \frac{2\pi}{3} = \sin \left(\pi - \frac{\pi}{3} \right) = \sin \frac{\pi}{3}$
 $\therefore \sin^{-1} \left(\sin \frac{2\pi}{3} \right) = \sin^{-1} \left(\sin \frac{\pi}{3} \right) = \frac{\pi}{3}$
Hence, the principal value of $\sin^{-1} \left(\sin \frac{2\pi}{3} \right)$ is $\frac{\pi}{3}$.

4. $\cot^{-1}(-\sqrt{3}) + \tan^{-1}(1) + \sec^{-1} \left(\frac{2}{\sqrt{3}} \right)$
 $= \cot^{-1} \cot \left(\frac{5\pi}{6} \right) + \tan^{-1} \left(\tan \frac{\pi}{4} \right) + \sec^{-1} \left(\sec \frac{\pi}{6} \right)$
 $= \frac{5\pi}{6} + \frac{\pi}{4} + \frac{\pi}{6} = \frac{5\pi}{4}$

5. $\cos^{-1} \{ \cos(-680^\circ) \} = \cos^{-1}(\cos 680^\circ)$
 $= \cos^{-1} \left(\cos \frac{34\pi}{9} \right) = \cos^{-1} \left\{ \cos \left(4\pi - \frac{2\pi}{9} \right) \right\}$
 $= \cos^{-1} \left(\cos \frac{2\pi}{9} \right) = \frac{2\pi}{9} = 40^\circ$

6. (i) We have, $\sin(\cot^{-1} x) = \sin \left(\cot^{-1} \frac{x}{1} \right)$
Now, $\cot^{-1} x = \sin^{-1} \left(\frac{1}{\sqrt{1+x^2}} \right)$
Hence, $\sin(\cot^{-1} x) = \sin \left(\sin^{-1} \left(\frac{1}{\sqrt{1+x^2}} \right) \right) = \frac{1}{\sqrt{1+x^2}}$

(ii) $\cos(\tan^{-1} x) = \cos \left(\tan^{-1} \frac{x}{1} \right)$
Now, $\tan^{-1} x = \cos^{-1} \left(\frac{1}{\sqrt{1+x^2}} \right)$
Hence, $\cos(\tan^{-1} x) = \cos \left(\cos^{-1} \left(\frac{1}{\sqrt{1+x^2}} \right) \right) = \frac{1}{\sqrt{1+x^2}}$

7. L.H.S. = $\sin^{-1} \frac{8}{17} + \sin^{-1} \frac{3}{5} = \cos^{-1} \frac{15}{17} + \cos^{-1} \frac{4}{5}$
 $= \cos^{-1} \left\{ \left(\frac{15}{17} \times \frac{4}{5} \right) - \sqrt{1 - \left(\frac{15}{17} \right)^2} \cdot \sqrt{1 - \left(\frac{4}{5} \right)^2} \right\}$
 $= \cos^{-1} \left\{ \frac{12}{17} - \sqrt{\frac{64}{289}} \cdot \sqrt{\frac{9}{25}} \right\} = \cos^{-1} \left\{ \frac{12}{17} - \frac{8}{17} \times \frac{3}{5} \right\}$
 $= \cos^{-1} \left\{ \frac{12}{17} - \frac{24}{85} \right\} = \cos^{-1} \left(\frac{36}{85} \right) = \text{R.H.S.}$

Hence, $\sin^{-1} \frac{8}{17} + \sin^{-1} \frac{3}{5} = \cos^{-1} \frac{36}{85}$.

8. Let $\tan^{-1} x = \phi$. Then, $\tan \phi = \frac{x}{1}$
 $\therefore \cos \phi = \frac{1}{\sqrt{1+x^2}} \Rightarrow \phi = \cos^{-1} \left(\frac{1}{\sqrt{1+x^2}} \right)$
So, $\cos(\tan^{-1} x) = \cos \left(\cos^{-1} \left(\frac{1}{\sqrt{1+x^2}} \right) \right) = \frac{1}{\sqrt{1+x^2}}$

Also, let $\sec^{-1} \frac{13}{12} = \theta$. Then, $\sec \theta = \frac{13}{12}$
 $\therefore \sin \theta = \frac{5}{13} \Rightarrow \theta = \sin^{-1} \frac{5}{13}$

So, $\sin \left(\sec^{-1} \frac{13}{12} \right) = \sin \left(\sin^{-1} \frac{5}{13} \right) = \frac{5}{13}$
Thus, $\frac{1}{\sqrt{1+x^2}} = \frac{5}{13} \Rightarrow \frac{1}{(1+x^2)} = \frac{25}{169}$
 $\Rightarrow 1+x^2 = \frac{169}{25} \Rightarrow x^2 = \frac{169}{25} - 1 \Rightarrow x = \pm \frac{144}{25}$


Hence, $x = \pm \frac{12}{5}$.

9. We have, $\sin^{-1} x = \frac{\pi}{6} + \cos^{-1} x$
 $\Rightarrow \sin^{-1} x - \cos^{-1} x = \frac{\pi}{6}$
 $\Rightarrow \sin^{-1} x - \left(\frac{\pi}{2} - \sin^{-1} x \right) = \frac{\pi}{6}$
 $\Rightarrow 2\sin^{-1} x = \frac{2\pi}{3} \Rightarrow \sin^{-1} x = \frac{\pi}{3} \Rightarrow x = \frac{\sqrt{3}}{2}$

10. We have, $\angle A = \tan^{-1} 2$, $\angle B = \tan^{-1} 3$
We know that, $\angle A + \angle B + \angle C = \pi$
 $\Rightarrow \tan^{-1} 2 + \tan^{-1} 3 + \angle C = \pi$
 $\Rightarrow \pi + \tan^{-1} \left(\frac{2+3}{1-2 \times 3} \right) + \angle C = \pi$
 $\Rightarrow \pi + \tan^{-1}(-1) + \angle C = \pi \Rightarrow \pi - \frac{\pi}{4} + \angle C = \pi$
 $\Rightarrow \frac{3\pi}{4} + \angle C = \pi \Rightarrow \angle C = \frac{\pi}{4}$

11. (i) $\sec x + \tan x = \frac{1 + \sin x}{\cos x} = \frac{1 - \cos \left(\frac{\pi}{2} + x \right)}{\sin \left(\frac{\pi}{2} + x \right)}$
 $= \frac{2 \sin^2 \left(\frac{\pi}{4} + \frac{x}{2} \right)}{2 \sin \left(\frac{\pi}{4} + \frac{x}{2} \right) \cos \left(\frac{\pi}{4} + \frac{x}{2} \right)} = \tan \left(\frac{\pi}{4} + \frac{x}{2} \right)$

Concerned about your performance in Class XII Boards?



Well, fear no more, help is at hand.....

To excel, studying in right direction is more important than studying hard. Which is why we created the Excel Series. These books – for Physics, Chemistry, Biology & Mathematics – have been put together totally keeping in mind the prescribed syllabus and the pattern of CBSE's Board examinations, so that students prepare and practice with just the right study material to excel in board exams.

Did you know nearly all questions in CBSE's 2017 Board Examination were a part of our Excel books? That too fully solved !

HIGHLIGHTS:

- Comprehensive theory strictly based on NCERT, complemented with illustrations, activities and solutions of NCERT questions
- Practice questions & Model Test Papers for Board Exams
- Value based questions
- Previous years' CBSE Board Examination Papers (Solved)
- CBSE Board Papers 2017 Included



Scan now with your smartphone or tablet*



Available at all leading book shops throughout the country.

For more information or for help in placing your order:

Call 0124-6601200 or email: info@mtg.in

*Application to read QR codes required

Visit
www.mtg.in
for latest offers
and to buy
online!

$$\therefore \tan^{-1}\left(\tan\left(\frac{\pi}{4} + \frac{x}{2}\right)\right) = \frac{\pi}{4} + \frac{x}{2}$$

(ii) Put $x = \cos \theta$

$$\begin{aligned}\therefore \sin\left(2\tan^{-1}\sqrt{\frac{1-x}{1+x}}\right) &= \sin\left(2\tan^{-1}\sqrt{\frac{1-\cos\theta}{1+\cos\theta}}\right) \\ &= \sin\left(2\tan^{-1}\sqrt{\frac{2\sin^2\theta/2}{2\cos^2\theta/2}}\right) \\ &= \sin\left[2\tan^{-1}\left(\tan\frac{\theta}{2}\right)\right] = \sin\theta = \sqrt{1-x^2}\end{aligned}$$

12. Given expression is $2\tan^{-1}\left(\frac{1+x}{1-x}\right) + \sin^{-1}\left(\frac{1-x^2}{1+x^2}\right)$

Putting $x = \tan \theta$, we get

$$\begin{aligned}&2\tan^{-1}\left(\frac{1+\tan\theta}{1-\tan\theta}\right) + \sin^{-1}\left(\frac{1-\tan^2\theta}{1+\tan^2\theta}\right) \\ &= 2\tan^{-1}\left(\frac{\tan\frac{\pi}{4} + \tan\theta}{1 - \tan\frac{\pi}{4} \cdot \tan\theta}\right) + \sin^{-1}(\cos 2\theta) \\ &= 2\tan^{-1}\left[\tan\left(\frac{\pi}{4} + \theta\right)\right] + \sin^{-1}\left[\sin\left(\frac{\pi}{2} - 2\theta\right)\right] \\ &= 2\left(\frac{\pi}{4} + \theta\right) + \frac{\pi}{2} - 2\theta = \frac{\pi}{2} + 2\theta + \frac{\pi}{2} - 2\theta = \pi\end{aligned}$$

13. (i) $\sin(2\sin^{-1}0.8) = \sin\left[\sin^{-1}\left(2\left(0.8\sqrt{1-(0.8)^2}\right)\right)\right]$
 $= \sin[\sin^{-1}(2 \times 0.8 \times 0.6)]$
 $= \sin[\sin^{-1}(0.96)] = 0.96$

$$\begin{aligned}\text{(ii)} \quad \tan\left[2\tan^{-1}\frac{1}{5} - \frac{\pi}{4}\right] &= \tan\left[\tan^{-1}\left(\frac{2\left(\frac{1}{5}\right)}{1 - \left(\frac{1}{5}\right)^2}\right) - \frac{\pi}{4}\right] \\ &= \tan\left[\tan^{-1}\frac{5}{12} - \frac{\pi}{4}\right] = \tan\left[\tan^{-1}\frac{5}{12} - \tan^{-1}1\right] \\ &= \tan\left[\tan^{-1}\left(\frac{\frac{5}{12} - 1}{1 + \frac{5}{12} \times 1}\right)\right] = \tan\left[\tan^{-1}\left(\frac{-7}{17}\right)\right] = -\frac{7}{17}\end{aligned}$$

$$14. \tan^{-1}\left(\tan\frac{5\pi}{6}\right) + \cos^{-1}\left(\cos\frac{13\pi}{6}\right) + \sec^{-1}\left(\sec\frac{9\pi}{5}\right)$$

$$\begin{aligned}&\Rightarrow \tan^{-1}\left[\tan\left(\pi - \frac{\pi}{6}\right)\right] + \cos^{-1}\left[\cos\left(2\pi + \frac{\pi}{6}\right)\right] \\ &\quad + \sec^{-1}\left[\sec\left(2\pi - \frac{\pi}{5}\right)\right] \\ &\Rightarrow \tan^{-1}\left(-\tan\left(\frac{\pi}{6}\right)\right) + \cos^{-1}\left[\cos\left(\frac{\pi}{6}\right)\right] \\ &\quad + \sec^{-1}\left[\sec\left(\frac{\pi}{5}\right)\right] \\ &= -\frac{\pi}{6} + \frac{\pi}{6} + \frac{\pi}{5} = \frac{\pi}{5}\end{aligned}$$

15. We have, $\tan^{-1}\left\{\frac{\sqrt{1+x^2} - \sqrt{1-x^2}}{\sqrt{1+x^2} + \sqrt{1-x^2}}\right\} = \alpha$

$$\begin{aligned}&\Rightarrow \frac{\sqrt{1+x^2} - \sqrt{1-x^2}}{\sqrt{1+x^2} + \sqrt{1-x^2}} = \tan \alpha \\ &\Rightarrow \frac{(\sqrt{1+x^2} - \sqrt{1-x^2}) + (\sqrt{1+x^2} + \sqrt{1-x^2})}{(\sqrt{1+x^2} - \sqrt{1-x^2}) - (\sqrt{1+x^2} + \sqrt{1-x^2})} = \frac{\tan \alpha + 1}{\tan \alpha - 1}\end{aligned}$$

$$\Rightarrow \frac{2\sqrt{1+x^2}}{-2\sqrt{1-x^2}} = \frac{\tan \alpha + 1}{\tan \alpha - 1} \Rightarrow \frac{\sqrt{1-x^2}}{\sqrt{1+x^2}} = \frac{1 - \tan \alpha}{1 + \tan \alpha}$$

$$\Rightarrow \frac{\sqrt{1-x^2}}{\sqrt{1+x^2}} = \frac{\cos \alpha - \sin \alpha}{\cos \alpha + \sin \alpha}$$

$$\Rightarrow \frac{1-x^2}{1+x^2} = \frac{1 - \sin 2\alpha}{1 + \sin 2\alpha} \Rightarrow x^2 = \sin 2\alpha$$

16. It is given that $a + b + c = abc$

$$\therefore \frac{abc}{c} = \frac{a}{c} + \frac{b}{c} + 1$$

$$\begin{aligned}&\Rightarrow ab = 1 + \left(\frac{a}{c} + \frac{b}{c}\right) \Rightarrow ab - 1 = \frac{a+b}{c} \\ &\Rightarrow ab - 1 > 0 \quad \left[\because a, b, c > 0 \therefore \frac{a+b}{c} > 0\right] \\ &\Rightarrow ab > 1\end{aligned}$$

Now, $\tan^{-1}a + \tan^{-1}b + \tan^{-1}c$

$$= \pi + \tan^{-1}\left(\frac{a+b}{1-ab}\right) + \tan^{-1}c \quad [\because ab > 1]$$

$$= \pi + \tan^{-1}\left(\frac{abc-c}{1-ab}\right) + \tan^{-1}c$$

$$= \pi + \tan^{-1}\left(\frac{-c(1-ab)}{1-ab}\right) + \tan^{-1}c$$

$$= \pi + \tan^{-1}(-c) + \tan^{-1}c = \pi - \tan^{-1}c + \tan^{-1}c = \pi$$

17. We have $\cos^{-1} \frac{x}{a} + \cos^{-1} \frac{y}{b} = \alpha$

$$\Rightarrow \cos^{-1} \left[\frac{x}{a} \cdot \frac{y}{b} - \sqrt{\left(1 - \frac{x^2}{a^2}\right) \left(1 - \frac{y^2}{b^2}\right)} \right] = \alpha$$

$$\Rightarrow \frac{xy}{ab} - \sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{x^2 y^2}{a^2 b^2}} = \cos \alpha$$

$$\Rightarrow \frac{xy}{ab} - \cos \alpha = \sqrt{1 - \frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{x^2 y^2}{a^2 b^2}}$$

Squaring both sides, we have

$$\frac{x^2 y^2}{a^2 b^2} - \frac{2xy}{ab} \cos \alpha + \cos^2 \alpha = 1 - \frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{x^2 y^2}{a^2 b^2}$$

$$\Rightarrow \frac{x^2}{a^2} - \frac{2xy}{ab} \cos \alpha + \frac{y^2}{b^2} = 1 - \cos^2 \alpha$$

$$\therefore \frac{x^2}{a^2} - \frac{2xy}{ab} \cos \alpha + \frac{y^2}{b^2} = \sin^2 \alpha$$

18. Given, $a_1, a_2, a_3, a_4, \dots, a_n$ is an arithmetic progression, then $d = a_2 - a_1 = a_3 - a_2 = \dots = a_n - a_{n-1}$

$$\therefore \tan \left[\tan^{-1} \left(\frac{d}{1+a_1 a_2} \right) + \tan^{-1} \left(\frac{d}{1+a_2 a_3} \right) \right. \\ \left. + \tan^{-1} \left(\frac{d}{1+a_3 a_4} \right) + \dots + \tan^{-1} \left(\frac{d}{1+a_{n-1} a_n} \right) \right]$$

$$= \tan \left[\tan^{-1} \left(\frac{a_2 - a_1}{1+a_1 a_2} \right) + \tan^{-1} \left(\frac{a_3 - a_2}{1+a_2 a_3} \right) \right. \\ \left. + \tan^{-1} \left(\frac{a_4 - a_3}{1+a_3 a_4} \right) + \dots + \tan^{-1} \left(\frac{a_n - a_{n-1}}{1+a_{n-1} a_n} \right) \right]$$

$$= \tan [\tan^{-1} a_2 - \tan^{-1} a_1 + \tan^{-1} a_3 - \tan^{-1} a_2 + \dots + \tan^{-1} a_n - \tan^{-1} a_{n-1}]$$

$$= \tan [\tan^{-1} a_n - \tan^{-1} a_1]$$

$$= \tan \left[\tan^{-1} \left(\frac{a_n - a_1}{1+a_n a_1} \right) \right] = \frac{a_n - a_1}{1+a_n a_1}$$

19. (i) Here, $\cos \left(\sin^{-1} \frac{1}{3} + \cos^{-1} x \right) = 0 = \cos \left(\pm \frac{\pi}{2} \right)$

$$\therefore \sin^{-1} \frac{1}{3} + \cos^{-1} x = \pm \frac{\pi}{2}$$

$$\text{i.e., } \cos^{-1} x = \pm \frac{\pi}{2} - \sin^{-1} \frac{1}{3}$$

$$\therefore x = \cos \left(\pm \frac{\pi}{2} - \sin^{-1} \frac{1}{3} \right)$$

$$x = \cos \left(\frac{\pi}{2} - \sin^{-1} \frac{1}{3} \right) \quad \left| \quad x = \cos \left(-\frac{\pi}{2} - \sin^{-1} \frac{1}{3} \right) \right.$$

$$= \sin \left(\sin^{-1} \frac{1}{3} \right) = \frac{1}{3} \quad \left| \quad = \cos \left(\frac{\pi}{2} + \sin^{-1} \frac{1}{3} \right) \right.$$

$$\therefore x = \pm \frac{1}{3} \quad \left| \quad = -\sin \left(\sin^{-1} \frac{1}{3} \right) = -\frac{1}{3} \right.$$

(ii) We have,

$$\tan^{-1} \left(\frac{x+1}{x-1} \right) + \tan^{-1} \left(\frac{x-1}{x} \right) = \pi + \tan^{-1}(-7)$$

$$\Rightarrow \tan^{-1} \left(\frac{\frac{x+1}{x-1} + \frac{x-1}{x}}{1 - \frac{x+1}{x-1} \cdot \frac{x-1}{x}} \right) = \pi + \tan^{-1}(-7)$$

$$\therefore \frac{(x+1)x + (x-1)^2}{(x-1)x - (x+1)(x-1)} = \tan[\pi + \tan^{-1}(-7)]$$

$$\Rightarrow x^2 + x + x^2 - 2x + 1 = -7(x^2 - x - x^2 + 1)$$

$$\Rightarrow 2x^2 - 8x + 8 = 0 \text{ i.e., } x^2 - 4x + 4 = 0$$

$$\Rightarrow (x-2)^2 = 0$$

$$\therefore x = 2$$

20. (i) L.H.S. = $\sin^{-1} \frac{12}{13} + \cos^{-1} \frac{4}{5} + \tan^{-1} \frac{63}{16}$

$$= \tan^{-1} \frac{12}{5} + \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{63}{16}$$

$$= \pi + \tan^{-1} \left(\frac{\frac{12}{5} + \frac{3}{4}}{1 - \frac{12}{5} \times \frac{3}{4}} \right) + \tan^{-1} \frac{63}{16}$$

$$= \pi + \tan^{-1} \left(-\frac{63}{16} \right) + \tan^{-1} \frac{63}{16}$$

$$= \pi - \tan^{-1} \frac{63}{16} + \tan^{-1} \frac{63}{16} = \pi$$

(ii) Let $\cos^{-1} \frac{4}{5} = \theta$. Then $\cos \theta = \frac{4}{5}$.

$$\therefore \tan \theta = \frac{\sqrt{1 - \cos^2 \theta}}{\cos \theta} = \frac{3}{4} \Rightarrow \theta = \tan^{-1} \frac{3}{4}$$

Consequently, $\cos^{-1} \frac{4}{5} = \tan^{-1} \frac{3}{4}$

$$\therefore \cos^{-1} \frac{4}{5} + \tan^{-1} \frac{3}{5} = \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{3}{5}$$


$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{3}{5}}{1 - \frac{3}{4} \cdot \frac{3}{5}} \right) = \tan^{-1} \frac{27}{11}$$



MPP-2 MONTHLY Practice Problems

Class XII

This specially designed column enables students to self analyse their extent of understanding of specified chapters. Give yourself four marks for correct answer and deduct one mark for wrong answer. Self check table given at the end will help you to check your readiness.



Inverse Trigonometric Functions

Total Marks : 80

Time Taken : 60 Min.

Only One Option Correct Type


- The number of positive integral solutions of $\tan^{-1} x + \cos^{-1} \frac{y}{\sqrt{1+y^2}} = \sin^{-1} \frac{3}{\sqrt{10}}$ is
(a) 0 (b) 1 (c) 2 (d) > 2
- If $\left| \cos^{-1} \left(\frac{1-x^2}{1+x^2} \right) \right| < \frac{\pi}{3}$, then x belongs to the interval
(a) $\left[-\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right]$ (b) $\left(-\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right)$
(c) $\left(0, \frac{1}{\sqrt{3}} \right)$ (d) none of these
- $\cos^{-1} \left[\cos \left(-\frac{17\pi}{15} \right) \right]$ is equal to
(a) $\frac{17\pi}{15}$ (b) $\frac{13\pi}{15}$
(c) $\frac{3\pi}{15}$ (d) $-\frac{17\pi}{15}$
- The domain of the function $f(x) = \cos^{-1} \left(\frac{2-|x|}{4} \right) + [\log(3-x)]^{-1}$ is
(a) $[-6, 6]$ (b) $[-5, 2) \cup (2, 3)$
(c) $(2, 3)$ (d) $[-6, 2) \cup (2, 3)$
- If θ and ϕ are the roots of the equation $8x^2 + 22x + 5 = 0$, then
(a) both $\sin^{-1} \theta$ and $\sin^{-1} \phi$ are real
(b) both $\sec^{-1} \theta$ and $\sec^{-1} \phi$ are real
(c) both $\tan^{-1} \theta$ and $\tan^{-1} \phi$ are real
(d) none of these

- If $\sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \frac{3\pi}{2}$, then the value of $25(x+y+z) - \frac{216}{(x^3+y^3+z^3)}$ must be
(a) 1 (b) 2
(c) 3 (d) none of these

One or More Than One Option(s) Correct Type

- If $\cos^{-1} x + (\sin^{-1} y)^2 = \frac{p\pi^2}{4}$ and $(\cos^{-1} x)(\sin^{-1} y)^2 = \frac{\pi^2}{16}$, then
(a) $0 \leq p \leq \frac{4}{\pi} + 1$
(b) $p = 2$ is the integral value of p
(c) $p = 0, 1, 2$ (integral values)
(d) none of these
- For $0 < \phi < \frac{\pi}{2}$, if $x = \sum_{n=0}^{\infty} \cos^{2n} \phi$, $y = \sum_{n=0}^{\infty} \sin^{2n} \phi$, $z = \sum_{n=0}^{\infty} \cos^{2n} \phi \sin^{2n} \phi$, then
(a) $xyz = xz + y$ (b) $xyz = xy + z$
(c) $xyz = x + y + z$ (d) $xyz = yz + x$
- For the equation $2x = \tan(2 \tan^{-1} a) + 2 \tan(\tan^{-1} a + \tan^{-1} a^3)$, which of the following is invalid?
(a) $a^2 x + 2a = x$ (b) $a^2 + 2ax + 1 = 0$
(c) $a \neq 0$ (d) $a \neq -1, 1$
- $\tan^{-1} \left(\frac{a_1 x - y}{a_1 y + x} \right) + \tan^{-1} \left(\frac{a_2 - a_1}{a_1 a_2 + 1} \right) + \tan^{-1} \left(\frac{a_3 - a_2}{a_2 a_3 + 1} \right) + \dots + \tan^{-1} \left(\frac{a_n - a_{n-1}}{a_{n-1} a_n + 1} \right) + \tan^{-1} \frac{1}{a_n}$ is equal to

BEST TOOLS FOR SUCCESS IN JEE Main



10 Very Similar Practice Test Papers

16 Years JEE MAIN 2017-2015(Offline & Online)-2013 & AIEEE (2012-2002)



Available at all leading book shops throughout India.
For more information or for help in placing your order:
Call 0124-6601200 or email: info@mtg.in

Visit
www.mtg.in
for latest offers
and to buy
online!

- (a) $\tan^{-1} xy$ (b) $\tan^{-1} \frac{x}{y}$
 (c) $\tan^{-1} \frac{y}{x}$ (d) none of these

11. The value of θ for which

$$\theta = \tan^{-1}(2 \tan^2 \theta) - \frac{1}{2} \sin^{-1} \left(\frac{3 \sin 2\theta}{5 + 4 \cos 2\theta} \right), \text{ is/are:}$$

- (a) $n\pi + \tan^{-1}(-2)$ (b) $n\pi, n\pi + \frac{\pi}{4}$
 (c) $n\pi + \cot^{-1}(-2)$ (d) none of these

12. If x, y and z are in A.P. and $\tan^{-1} x, \tan^{-1} y$ and $\tan^{-1} z$ are also in A.P. then

- (a) $x = y = z$ (b) $2x = 3y = 6z$
 (c) $6x = 3y = 2z$ (d) $6x = 4y = 3z$

13. In $\triangle ABC$, $\angle C = \frac{\pi}{2}$ and

$$\sin^{-1} x = \sin^{-1} \left(\frac{ax}{c} \right) + \sin^{-1} \left(\frac{bx}{c} \right)$$

where a, b and c are the sides of triangle, then the values of x is/are

- (a) 0 (b) 1 (c) 2 (d) -1

Comprehension Type

$$\sum_{r=1}^n \tan^{-1} \left(\frac{x_r - x_{r-1}}{1 + x_{r-1} x_r} \right) = \sum_{r=1}^n \left(\tan^{-1} x_r - \tan^{-1} x_{r-1} \right)$$

$$= \tan^{-1} x_n - \tan^{-1} x_0, \forall n \in N$$

On the basis of above information, answer the following questions:

14. The value of $\operatorname{cosec}^{-1} \sqrt{5} + \operatorname{cosec}^{-1} \sqrt{65} +$

$\operatorname{cosec}^{-1} \sqrt{(325)} + \dots$ to ∞ is

- (a) π (b) $\frac{3\pi}{4}$ (c) $\frac{\pi}{2}$ (d) $\frac{\pi}{4}$

15. The sum to infinite terms of the series

$$\tan^{-1} \left(\frac{2}{1-1^2+1^4} \right) + \tan^{-1} \left(\frac{4}{1-2^2+2^4} \right)$$

$$+ \tan^{-1} \left(\frac{6}{1-3^2+3^4} \right) + \dots \text{ is}$$

- (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{2}$
 (c) $\frac{3\pi}{4}$ (d) none of these

Matrix Match Type

16. Match the following :

Column I		Column II	
P.	If $2 \tan^{-1}(2x+1) = \cos^{-1}(-x)$, then x is	1.	$-\frac{1}{\sqrt{2}}$
Q.	If $2 \cos^{-1} x = \sin^{-1} \left(2x\sqrt{1-x^2} \right)$, then x is	2.	$\frac{\sqrt{3}}{2}$
R.	If $\tan^{-1} \left(\frac{x-1}{x-2} \right) + \tan^{-1} \left(\frac{x+1}{x+2} \right) = \frac{\pi}{4}$, then x is	3.	-1
		4.	0

	P	Q	R
(a)	2	1	3
(b)	3	2	1
(c)	3	1	2
(d)	4	2	1

Integer Answer Type

17. The number of solutions for the equation

$$2 \sin^{-1} \sqrt{(x^2 - x + 1)} + \cos^{-1} \sqrt{(x^2 - x)} = \frac{3\pi}{2} \text{ is}$$

18. If $\tan \theta + \tan \left(\frac{\pi}{3} + \theta \right) + \tan \left(-\frac{\pi}{3} + \theta \right) = a \tan 3\theta$,

then a is equal to

19. If $\sin^{-1} x + \sin^{-1} y + \sin^{-1} z = \frac{3\pi}{2}$ and $f(1) = 2$,

$f(p+q) = f(p) \cdot f(q), \forall p, q \in R$, then

$x^{f(1)} + y^{f(2)} + z^{f(3)} - \frac{(x+y+z)}{x^{f(1)} + y^{f(2)} + z^{f(3)}}$ is equal to

20. If $\lambda = \tan \left(2 \tan^{-1} \frac{1}{5} - \frac{\pi}{4} \right)$, then the value of $-17 \lambda^2$ must be



Keys are published in this issue. Search now! ☺

SELF CHECK

No. of questions attempted

No. of questions correct

Marks scored in percentage

Check your score! If your score is

> 90%	EXCELLENT WORK !	You are well prepared to take the challenge of final exam.
90-75%	GOOD WORK !	You can score good in the final exam.
74-60%	SATISFACTORY !	You need to score more next time.
< 60%	NOT SATISFACTORY!	Revise thoroughly and strengthen your concepts.



WB JEE

SOLVED PAPER 2017

CATEGORY-I (Q. 1 to Q. 50)

Only one answer is correct. Correct answer will fetch full marks 1. Incorrect answer or any combination of more than one answer will fetch $-\frac{1}{4}$ marks. No answer will fetch 0 marks.

- The number of all numbers having 5 digits, with distinct digits is
(a) 99999 (b) $9 \times {}^9P_4$ (c) ${}^{10}P_5$ (d) 9P_4
- The greatest integer which divides $(p+1)(p+2)(p+3)\dots(p+q)$ for all $p \in N$ and fixed $q \in N$ is
(a) $p!$ (b) $q!$ (c) p (d) q
- Let $(1+x+x^2)^9 = a_0 + a_1x + a_2x^2 + \dots + a_{18}x^{18}$. Then
(a) $a_0 + a_2 + \dots + a_{18} = a_1 + a_3 + \dots + a_{17}$
(b) $a_0 + a_2 + \dots + a_{18}$ is even
(c) $a_0 + a_2 + \dots + a_{18}$ is divisible by 9
(d) $a_0 + a_2 + \dots + a_{18}$ is divisible by 3 but not by 9
- The linear system of equations
$$\begin{cases} 8x - 3y - 5z = 0 \\ 5x - 8y + 3z = 0 \\ 3x + 5y - 8z = 0 \end{cases}$$
 has
(a) only zero solution
(b) only finite number of non-zero solutions
(c) no non-zero solution
(d) infinitely many non-zero solutions
- Let P be the set of all non-singular matrices of order 3 over R and Q be the set of all orthogonal matrices of order 3 over R . Then
(a) P is proper subset of Q
(b) Q is proper subset of P
(c) Neither P is proper subset of Q nor Q is proper subset of P
(d) $P \cap Q = \phi$, the void set
- Let $A = \begin{pmatrix} x+2 & 3x \\ 3 & x+2 \end{pmatrix}$, $B = \begin{pmatrix} x & 0 \\ 5 & x+2 \end{pmatrix}$. Then all solutions of the equation $\det(AB) = 0$ is
(a) 1, -1, 0, 2 (b) 1, 4, 0, -2
(c) 1, -1, 4, 3 (d) -1, 4, 0, 3
- The value of $\det A$, where $A = \begin{pmatrix} 1 & \cos\theta & 0 \\ -\cos\theta & 1 & \cos\theta \\ -1 & -\cos\theta & 1 \end{pmatrix}$ lies
(a) in the closed interval $[1, 2]$
(b) in the closed interval $[0, 1]$
(c) in the open interval $(0, 1)$
(d) in the open interval $(1, 2)$
- Let $f: R \rightarrow R$ be such that f is injective and $f(x)f(y) = f(x+y)$, $\forall x, y \in R$. If $f(x), f(y), f(z)$ are in G.P., then x, y, z are in
(a) A.P. always
(b) G.P. always
(c) A.P. depending on the value of x, y, z
(d) G.P. depending on the value of x, y, z
- On the set R of real numbers we define xPy if and only if $xy \geq 0$. Then the relation P is
(a) reflexive but not symmetric
(b) symmetric but not reflexive
(c) transitive but not reflexive
(d) reflexive and symmetric but not transitive
- On R , the relation ρ be defined by ' $x\rho y$ holds if and only if $x - y$ is zero or irrational'. Then
(a) ρ is reflexive and transitive but not symmetric.
(b) ρ is reflexive and symmetric but not transitive.
(c) ρ is symmetric and transitive but not reflexive.
(d) ρ is equivalence relation

11. Mean of n observations x_1, x_2, \dots, x_n is \bar{x} . If an observation x_q is replaced by x'_q then the new mean is
- (a) $\bar{x} - x_q + x'_q$ (b) $\frac{(n-1)\bar{x} + x'_q}{n}$
 (c) $\frac{(n-1)\bar{x} - x_q + x'_q}{n}$ (d) $\frac{n\bar{x} - x_q + x'_q}{n}$
12. The probability that a non leap year selected at random will have 53 Sundays is
 (a) 0 (b) $1/7$ (c) $2/7$ (d) $3/7$
13. The equation $\sin x(\sin x + \cos x) = k$ has real solutions, where k is a real number. Then
 (a) $0 \leq k \leq \frac{1+\sqrt{2}}{2}$ (b) $2 - \sqrt{3} \leq k \leq 2 + \sqrt{3}$
 (c) $0 \leq k \leq 2 - \sqrt{3}$ (d) $\frac{1-\sqrt{2}}{2} \leq k \leq \frac{1+\sqrt{2}}{2}$
14. The possible values of x , which satisfy the trigonometric equation
 $\tan^{-1}\left(\frac{x-1}{x-2}\right) + \tan^{-1}\left(\frac{x+1}{x+2}\right) = \frac{\pi}{4}$ are
 (a) $\pm \frac{1}{\sqrt{2}}$ (b) $\pm \sqrt{2}$ (c) $\pm \frac{1}{2}$ (d) ± 2
15. Transforming to parallel axes through a point (p, q) , the equation $2x^2 + 3xy + 4y^2 + x + 18y + 25 = 0$ becomes $2x^2 + 3xy + 4y^2 = 1$. Then
 (a) $p = -2, q = 3$ (b) $p = 2, q = -3$
 (c) $p = 3, q = -4$ (d) $p = -4, q = 3$
16. Let $A(2, -3)$ and $B(-2, 1)$ be two angular points of $\triangle ABC$. If the centroid of the triangle moves on the line $2x + 3y = 1$, then the locus of the angular point C is given by
 (a) $2x + 3y = 9$ (b) $2x - 3y = 9$
 (c) $3x + 2y = 5$ (d) $3x - 2y = 3$
17. The point $P(3, 6)$ is first reflected on the line $y = x$ and then the image point Q is again reflected on the line $y = -x$ to get the image point Q' . Then the circumcentre of the $\triangle PQQ'$ is
 (a) $(6, 3)$ (b) $(6, -3)$ (c) $(3, -6)$ (d) $(0, 0)$
18. Let d_1 and d_2 be the lengths of the perpendiculars drawn from any point of the line $7x - 9y + 10 = 0$ upon the lines $3x + 4y = 5$ and $12x + 5y = 7$ respectively. Then
 (a) $d_1 > d_2$ (b) $d_1 = d_2$ (c) $d_1 < d_2$ (d) $d_1 = 2d_2$
19. The common chord of the circles $x^2 + y^2 - 4x - 4y = 0$ and $2x^2 + 2y^2 = 32$ subtends at the origin an angle equal to
 (a) $\frac{\pi}{3}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{6}$ (d) $\frac{\pi}{2}$
20. The locus of the mid-points of the chords of the circle $x^2 + y^2 + 2x - 2y - 2 = 0$ which make an angle of 90° at the centre is
 (a) $x^2 + y^2 - 2x - 2y = 0$
 (b) $x^2 + y^2 - 2x + 2y = 0$
 (c) $x^2 + y^2 + 2x - 2y = 0$
 (d) $x^2 + y^2 + 2x - 2y - 1 = 0$
21. Let P be the foot of the perpendicular from focus S of hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ on the line $bx - ay = 0$ and let C be the centre of the hyperbola. Then the area of the rectangle whose sides are equal to that of SP and CP is
 (a) $2ab$ (b) ab
 (c) $\frac{(a^2 + b^2)}{2}$ (d) $\frac{a}{b}$
22. B is an extremity of the minor axis of an ellipse whose foci are S and S' . If $\angle SBS'$ is a right angle, then the eccentricity of the ellipse is
 (a) $\frac{1}{2}$ (b) $\frac{1}{\sqrt{2}}$ (c) $\frac{2}{3}$ (d) $\frac{1}{3}$
23. The axis of the parabola $x^2 + 2xy + y^2 - 5x + 5y - 5 = 0$ is
 (a) $x + y = 0$ (b) $x + y - 1 = 0$
 (c) $x - y + 1 = 0$ (d) $x - y = \frac{1}{\sqrt{2}}$
24. The line segment joining the foci of the hyperbola $x^2 - y^2 + 1 = 0$ is one of the diameters of a circle. The equation of the circle is
 (a) $x^2 + y^2 = 4$ (b) $x^2 + y^2 = \sqrt{2}$
 (c) $x^2 + y^2 = 2$ (d) $x^2 + y^2 = 2\sqrt{2}$
25. The equation of the plane through $(1, 2, -3)$ and $(2, -2, 1)$ and parallel to X -axis is
 (a) $y - z + 1 = 0$ (b) $y - z - 1 = 0$
 (c) $y + z - 1 = 0$ (d) $y + z + 1 = 0$
26. Three lines are drawn from the origin O with direction cosines proportional to $(1, -1, 1)$, $(2, -3, 0)$ and $(1, 0, 3)$. The three lines are
 (a) not coplanar (b) coplanar

- (c) perpendicular to each other
(d) coincident
27. Consider the non-constant differentiable function f of one variable which obeys the relation $\frac{f(x)}{f(y)} = f(x-y)$. If $f'(0) = p$ and $f'(5) = q$, then $f'(-5)$ is
(a) $\frac{p^2}{q}$ (b) $\frac{q}{p}$ (c) $\frac{p}{q}$ (d) q
28. If $f(x) = \log_5 \log_3 x$, then $f'(e)$ is equal to
(a) $e \log_e 5$ (b) $e \log_e 3$
(c) $\frac{1}{e \log_e 5}$ (d) $\frac{1}{e \log_e 3}$
29. Let $F(x) = e^x$, $G(x) = e^{-x}$ and $H(x) = G(F(x))$, where x is a real variable. Then $\frac{dH}{dx}$ at $x = 0$ is
(a) 1 (b) -1 (c) $-\frac{1}{e}$ (d) $-e$
30. If $f''(0) = k$, $k \neq 0$, then the value of $\lim_{x \rightarrow 0} \frac{2f(x) - 3f(2x) + f(4x)}{x^2}$ is
(a) k (b) $2k$ (c) $3k$ (d) $4k$
31. If $y = e^{m \sin^{-1} x}$, then $(1 - x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} - ky = 0$, where k is equal to
(a) m^2 (b) 2 (c) -1 (d) $-m^2$
32. The chord of the curve $y = x^2 + 2ax + b$, joining the points where $x = \alpha$ and $x = \beta$, is parallel to the tangent to the curve at abscissa $x =$
(a) $\frac{a+b}{2}$ (b) $\frac{2a+b}{3}$ (c) $\frac{2\alpha+\beta}{3}$ (d) $\frac{\alpha+\beta}{2}$
33. Let $f(x) = x^{13} + x^{11} + x^9 + x^7 + x^5 + x^3 + x + 19$. Then $f(x) = 0$ has
(a) 13 real roots
(b) only one positive and only two negative real roots
(c) not more than one real root
(d) has two positive and one negative real root
34. Let $f(x) = \begin{cases} \frac{x^p}{(\sin x)^q}, & \text{if } 0 < x \leq \frac{\pi}{2} \\ 0, & \text{if } x = 0 \end{cases}$, $(p, q \in \mathbb{R})$. Then Lagrange's mean value theorem is applicable to $f(x)$ in closed interval $[0, x]$,
(a) for all p, q (b) only when $p > q$
(c) only when $p < q$ (d) for no value of p, q
35. $\lim_{x \rightarrow 0} (\sin x)^{2 \tan x}$
(a) is 2 (b) is 1
(c) is 0 (d) does not exist
36. $\int \cos(\log x) dx = F(x) + c$, where c is an arbitrary constant. Here $F(x) =$
(a) $x[\cos(\log x) + \sin(\log x)]$
(b) $x[\cos(\log x) - \sin(\log x)]$
(c) $\frac{x}{2} [\cos(\log x) + \sin(\log x)]$
(d) $\frac{x}{2} [\cos(\log x) - \sin(\log x)]$
37. $\frac{x^2 - 1}{x^4 + 3x^2 + 1} dx$ ($x > 0$) is
(a) $\tan^{-1}\left(x + \frac{1}{x}\right) + c$ (b) $\tan^{-1}\left(x - \frac{1}{x}\right) + c$
(c) $\log_e \left| \frac{x + \frac{1}{x} - 1}{x + \frac{1}{x} + 1} \right| + c$ (d) $\log_e \left| \frac{x - \frac{1}{x} - 1}{x - \frac{1}{x} + 1} \right| + c$
38. Let $I = \int_{10}^{19} \frac{\sin x}{1 + x^8} dx$. Then,
(a) $|I| < 10^{-9}$ (b) $|I| < 10^{-7}$
(c) $|I| < 10^{-5}$ (d) $|I| > 10^{-7}$
39. Let $I_1 = \int_0^n [x] dx$ and $I_2 = \int_0^n \{x\} dx$, where $[x]$ and $\{x\}$ are integral and fractional parts of x and $n \in \mathbb{N} - \{1\}$. Then I_1/I_2 is equal to
(a) $\frac{1}{n-1}$ (b) $\frac{1}{n}$ (c) n (d) $n-1$
40. The value of $\lim_{n \rightarrow \infty} \left[\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \dots + \frac{1}{2n} \right]$ is
(a) $\frac{n\pi}{4}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{4n}$ (d) $\frac{\pi}{2n}$
41. The value of the integral $\int_0^1 e^{x^2} dx$
(a) is less than 1
(b) is greater than 1
(c) is less than or equal to 1
(d) lies in the closed interval $[1, e]$

42. $\int_0^{100} e^{x-[x]} dx =$

- (a) $\frac{e^{100}-1}{100}$ (b) $\frac{e^{100}-1}{e-1}$
(c) $100(e-1)$ (d) $\frac{e-1}{100}$

43. Solution of $(x+y)^2 \frac{dy}{dx} = a^2$ ('a' being a constant) is

- (a) $\frac{(x+y)}{a} = \tan \frac{y+c}{a}$, c is an arbitrary constant
(b) $xy = a \tan cx$, c is an arbitrary constant
(c) $\frac{x}{a} = \tan \frac{y}{c}$, c is an arbitrary constant
(d) $xy = \tan(x+c)$, c is an arbitrary constant

44. The integrating factor of the first order differential equation

$$x^2(x^2-1) \frac{dy}{dx} + x(x^2+1)y = x^2-1$$

- (a) e^x (b) $x - \frac{1}{x}$ (c) $x + \frac{1}{x}$ (d) $\frac{1}{x^2}$

45. In a G.P. series consisting of positive terms, each term is equal to the sum of next two terms. Then the common ratio of this G.P. series is

- (a) $\sqrt{5}$ (b) $\frac{\sqrt{5}-1}{2}$ (c) $\frac{\sqrt{5}}{2}$ (d) $\frac{\sqrt{5}+1}{2}$

46. If $(\log_5 x)(\log_x 3x)(\log_{3x} y) = \log_x x^3$, then y equals

- (a) 125 (b) 25 (c) 5/3 (d) 243

47. The expression $\frac{(1+i)^n}{(1-i)^{n-2}}$ equals

- (a) $-i^{n+1}$ (b) i^{n+1} (c) $-2i^{n+1}$ (d) 1

48. Let $z = x + iy$, where x and y are real. The points

(x, y) in the X-Y plane for which $\frac{z+i}{z-i}$ purely imaginary lie on

- (a) a straight line (b) an ellipse
(c) a hyperbola (d) a circle

49. If p, q are odd integers, then the roots of the equation $2px^2 + (2p+q)x + q = 0$ are

- (a) rational (b) irrational
(c) non-real (d) equal

50. Out of 7 consonants and 4 vowels, words are formed each having 3 consonants and 2 vowels. The number of such words that can be formed is

- (a) 210 (b) 25200 (c) 2520 (d) 302400

CATEGORY-II (Q. 51 to Q. 65)

Only one answer is correct. Correct answer will fetch full marks 2. Incorrect answer or any combination of more than one answer will fetch $-\frac{1}{2}$ marks. No answer will fetch 0 marks.

51. Let $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ Then for positive integer n, A^n is

- (a) $\begin{pmatrix} 1 & n & n^2 \\ 0 & n^2 & n \\ 0 & 0 & n \end{pmatrix}$ (b) $\begin{pmatrix} 1 & n & n\left(\frac{n+1}{2}\right) \\ 0 & 1 & n \\ 0 & 0 & 1 \end{pmatrix}$
(c) $\begin{pmatrix} 1 & n^2 & n \\ 0 & n & n^2 \\ 0 & 0 & n^2 \end{pmatrix}$ (d) $\begin{pmatrix} 1 & n & 2n-1 \\ 0 & \frac{n+1}{2} & n^2 \\ 0 & 0 & \frac{n+1}{2} \end{pmatrix}$

52. Let a, b, c be such that $b(a+c) \neq 0$.

$$\text{If } \begin{vmatrix} a & a+1 & a-1 \\ -b & b+1 & b-1 \\ c & c-1 & c+1 \end{vmatrix} + \begin{vmatrix} a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \\ (-1)^{n+2}a & (-1)^{n+1}b & (-1)^n c \end{vmatrix} = 0$$

then the value of n is

- (a) any integer (b) zero
(c) any even integer (d) any odd integer

53. On set $A = \{1, 2, 3\}$, relations R and S are given by

$$R = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1)\}$$

$$S = \{(1, 1), (2, 2), (3, 3), (1, 3), (3, 1)\}.$$

- Then
(a) $R \cup S$ is an equivalence relation
(b) $R \cup S$ is reflexive and transitive but not symmetric
(c) $R \cup S$ is reflexive and symmetric but not transitive
(d) $R \cup S$ is symmetric and transitive but not reflexive

54. If one of the diameters of the curve $x^2 + y^2 - 4x - 6y + 9 = 0$ is a chord of a circle with centre (1, 1), the radius of the circle is

- (a) 3 (b) 2 (c) $\sqrt{2}$ (d) 1

55. Let A(-1, 0) and B(2, 0) be two points. A point M moves in the plane in such a way that $\angle MBA = 2\angle MAB$. Then the point M moves along

- (a) a straight line (b) a parabola
(c) an ellipse (d) a hyperbola

56. If $f(x) = \int_{-1}^x |t| dt$, then for any $x \geq 0$, $f(x)$ is equal to

- (a) $\frac{1}{2}(1-x^2)$ (b) $1-x^2$
(c) $\frac{1}{2}(1+x^2)$ (d) $1+x^2$

57. Let for all $x > 0$, $f(x) = \lim_{n \rightarrow \infty} n \left(x^{\frac{1}{n}} - 1 \right)$, then
- (a) $f(x) + f\left(\frac{1}{x}\right) = 1$ (b) $f(xy) = f(x) + f(y)$
 (c) $f(xy) = xf(y) + yf(x)$ (d) $f(xy) = xf(x) + yf(x)$

58. Let $I = \int_0^{100\pi} \sqrt{1 - \cos 2x} \, dx$, then

- (a) $I = 0$ (b) $I = 200\sqrt{2}$
 (c) $I = \pi\sqrt{2}$ (d) $I = 100$

59. The area of the figure bounded by the parabolas $x = -2y^2$ and $x = 1 - 3y^2$ is

- (a) $\frac{4}{3}$ square units (b) $\frac{2}{3}$ square units
 (c) $\frac{3}{7}$ square units (d) $\frac{6}{7}$ square units

60. Tangents are drawn to the ellipse $\frac{x^2}{9} + \frac{y^2}{5} = 1$ at the ends of both latusrectum. The area of the quadrilateral so formed is

- (a) 27 sq. units (b) $\frac{13}{2}$ sq. units
 (c) $\frac{15}{4}$ sq. units (d) 45 sq. units

61. The value of K in order that $f(x) = \sin x - \cos x - Kx + 5$ decreases for all positive real values of x is given by

- (a) $K < 1$ (b) $K \geq 1$ (c) $K > \sqrt{2}$ (d) $K < \sqrt{2}$

62. For any vector \vec{x} , the value of $(\vec{x} \times \hat{i})^2 + (\vec{x} \times \hat{j})^2 + (\vec{x} \times \hat{k})^2$ is equal to

- (a) $|\vec{x}|^2$ (b) $2|\vec{x}|^2$ (c) $3|\vec{x}|^2$ (d) $4|\vec{x}|^2$

63. If the sum of two unit vectors is a unit vector, then the magnitude of their difference is

- (a) $\sqrt{2}$ units (b) 2 units
 (c) $\sqrt{3}$ units (d) $\sqrt{5}$ units

64. Let α and β be the roots of $x^2 + x + 1 = 0$. If n be positive integer, then $\alpha^n + \beta^n$ is

- (a) $2\cos\frac{2n\pi}{3}$ (b) $2\sin\frac{2n\pi}{3}$
 (c) $2\cos\frac{n\pi}{3}$ (d) $2\sin\frac{n\pi}{3}$

65. For real x , the greatest value of $\frac{x^2 + 2x + 4}{2x^2 + 4x + 9}$ is

- (a) 1 (b) -1 (c) $\frac{1}{2}$ (d) $\frac{1}{4}$

CATEGORY-III (Q. 66 to Q. 75)

One or more answer(s) is (are) correct. Correct answer(s) will fetch full marks 2. Any combination containing one or more incorrect answer will fetch 0 marks. Also no answer will fetch 0 marks. If all correct answers are not marked and also no incorrect answer is marked then score = $2 \times \text{number of correct answers marked} \div \text{actual number of correct answers}$.

66. If $a, b \in \{1, 2, 3\}$ and the equation $ax^2 + bx + 1 = 0$ has real roots, then

- (a) $a > b$
 (b) $a \leq b$
 (c) number of possible ordered pairs (a, b) is 3
 (d) $a < b$

67. If the tangent to $y^2 = 4ax$ at the point $(at^2, 2at)$ where $|t| > 1$ is a normal to $x^2 - y^2 = a^2$ at the point $(a \sec \theta, a \tan \theta)$, then

- (a) $t = -\operatorname{cosec} \theta$ (b) $t = -\sec \theta$
 (c) $t = 2 \tan \theta$ (d) $t = 2 \cot \theta$

68. The focus of the conic $x^2 - 6x + 4y + 1 = 0$ is

- (a) (2, 3) (b) (3, 2) (c) (3, 1) (d) (1, 4)

69. Let $f: R \rightarrow R$ be twice continuously. Let $f(0) = f(1) = f'(0) = 0$. Then

- (a) $f''(x) \neq 0$ for all x
 (b) $f''(c) \neq 0$ for some $c \in R$
 (c) $f''(x) \neq 0$ if $x \neq 0$
 (d) $f'(x) > 0$ for all x

70. If $f(x) = x^n$, n being a non-negative integer, then the values of n for which $f'(\alpha + \beta) = f'(\alpha) + f'(\beta)$ for all $\alpha, \beta > 0$ is

- (a) 1 (b) 2 (c) 0 (d) 5

71. Let f be a non-constant continuous function for all $x \geq 0$. Let f satisfy the relation $f(x)f(a-x) = 1$ for

some $a \in R^+$. Then $I = \int_0^a \frac{dx}{1+f(x)}$ is equal to

- (a) a (b) $\frac{a}{4}$ (c) $\frac{a}{2}$ (d) $f(a)$

72. If the line $ax + by + c = 0$, $ab \neq 0$, is a tangent to the curve $xy = 1 - 2x$, then

- (a) $a > 0, b < 0$ (b) $a > 0, b > 0$
 (c) $a < 0, b > 0$ (d) $a < 0, b < 0$

73. Two particles move in the same straight line starting at the same moment from the same point in the same direction. The first moves with constant velocity u and the second starts from rest with constant acceleration f . Then

- (a) they will be at the greatest distance at the end of time $\frac{u}{2f}$ from the start
- (b) they will be at the greatest distance at the end of time $\frac{u}{f}$ from the start
- (c) their greatest distance is $\frac{u^2}{2f}$
- (d) their greatest distance is $\frac{u^2}{f}$

74. The complex number z satisfying the equation $|z - i| = |z + 1| = 1$ is

- (a) 0 (b) $1 + i$ (c) $-1 + i$ (d) $1 - i$

75. On R , the set of real numbers, a relation ρ is defined as ' apb ' if and only if $1 + ab > 0$. Then

- (a) ρ is an equivalence relation
 (b) ρ is reflexive and transitive but not symmetric
 (c) ρ is reflexive and symmetric but not transitive
 (d) ρ is only symmetric

SOLUTIONS

1. (b): 1st digit must be other than 0 (zero) i.e. 9 ways.
 \therefore Other 4 digits can be filled in 9P_4 ways
 \therefore Total no. of numbers having 5 (distinct) digits = $9 \times {}^9P_4$

2. (b): $\therefore (p+1)(p+2)(p+3)\dots(p+q)$ is a product of q consecutive positive integers
 \therefore It must be always divisible by $q!$

3. (b): $(1+x+x^2)^9 = a_0 + a_1x + a_2x^2 + \dots + a_{18}x^{18}$
 Putting $x = 1$ and -1 , we get

$$3^9 = a_0 + a_1 + a_2 + \dots + a_{18} \quad \dots(i)$$

$$1 = a_0 - a_1 + a_2 - \dots + a_{18} \quad \dots(ii)$$

Adding (i) & (ii), we get

$$\frac{3^9 + 1}{2} = a_0 + a_2 + a_4 + \dots + a_{18}$$

$\Rightarrow a_0 + a_2 + a_4 + \dots + a_{18} = 9842$, which is even but not divisible by 3 or 9.

$$4. (d): \therefore \begin{vmatrix} 8 & -3 & -5 \\ 5 & -8 & 3 \\ 3 & 5 & -8 \end{vmatrix} = \begin{vmatrix} 0 & -3 & -5 \\ 0 & -8 & 3 \\ 0 & 5 & -8 \end{vmatrix}$$

$$[C_1 \rightarrow C_1 + C_2 + C_3]$$

$$= 0$$

\therefore Given system of equation has infinitely many non-zero solutions.

5. (b): \therefore Every orthogonal matrix is non-singular but every non-singular matrix may or may not be orthogonal.

$\therefore Q$ is proper subset of P .

$$6. (b): AB = \begin{pmatrix} x+2 & 3x \\ 3 & x+2 \end{pmatrix} \begin{pmatrix} x & 0 \\ 5 & x+2 \end{pmatrix}$$

$$= \begin{pmatrix} x^2 + 17x & 3x^2 + 6x \\ 8x + 10 & x^2 + 4x + 4 \end{pmatrix}$$

$$\det(AB) = x(x+2) \begin{vmatrix} x+17 & 3 \\ 8x+10 & x+2 \end{vmatrix}$$

$$= x(x+2)(x^2 - 5x + 4) = x(x+2)(x-1)(x-4)$$

$$\therefore \det(AB) = 0 \Rightarrow x = 1, 4, 0, -2$$

$$7. (a): |A| = \begin{vmatrix} 1 & \cos \theta & 0 \\ -\cos \theta & 1 & \cos \theta \\ -1 & -\cos \theta & 1 \end{vmatrix} = (1 + \cos^2 \theta) - \cos \theta(0) + 0$$

$$= 1 + \cos^2 \theta \in [1, 2].$$

$$8. (c): \therefore f(x) \cdot f(y) = f(x+y) \quad \dots(i)$$

$$\therefore f(y)f(y) = f(y+y) \Rightarrow \{f(y)\}^2 = f(2y)$$

$$\Rightarrow f(x)f(z) = f(2y) \quad [\because f(x), f(y), f(z) \text{ are in G.P.}]$$

$$\Rightarrow f(x+z) = f(2y) \quad [\text{from (i)}]$$

$\Rightarrow x, y, z$ are in A.P. but will depend on the value of x, y, z .

$$9. (d): \therefore x^2 \geq 0 \therefore x \cdot x \geq 0 \Rightarrow xPx \Rightarrow \text{Reflexive}$$

$$\therefore xy \geq 0 \Rightarrow yx \geq 0 \Rightarrow \text{Symmetric}$$

$$\therefore (-5)(0) \geq 0 \text{ \& } (0)(7) \geq 0$$

$$\text{i.e., } (-5, 0) \in P \text{ \& } (0, 7) \in P$$

$$\text{But, } (-5)(7) < 0 \Rightarrow (-5, 7) \notin P$$

$\therefore P$ is not transitive.

10. (b): Here $(x, y) \in \rho$ if $x - y$ is zero or irrational.

$$\therefore x - x = 0 \text{ for all } x \in R$$

$\Rightarrow \rho$ is reflexive

If $x - y$ is zero or irrational then $y - x$ is also zero or irrational.

$\Rightarrow \rho$ is symmetric.

Let $(x, y) \in \rho$ & $(y, z) \in \rho$

$$\therefore x - y = 0 \text{ or irrational \& } y - z = 0 \text{ or irrational}$$

But, their sum $x - z$ may or may not be 0 or irrational

e.g., $2 - \sqrt{3}$ is irrational & $\sqrt{3} - 5$ both are irrational

but their sum $2 - 5 = -3$ is neither zero nor irrational

$\Rightarrow \rho$ is not transitive.

11. (d): \therefore Mean of n observations x_1, x_2, \dots, x_n is \bar{x} .

$$\therefore \text{Sum of } n \text{ observations} = n\bar{x}$$

If x_q is replaced by x'_q then sum = $n\bar{x} - x_q + x'_q$

$$\therefore \text{New mean} = \frac{n\bar{x} - x_q + x'_q}{n}$$

12. (b): A non-leap year has 52 weeks & 1 extra day

$$\therefore \text{Prob. of 53 sundays} = \frac{1}{7}$$

13. (d) : We have, $k = \sin^2 x + \sin x \cos x = \frac{1}{2} [(1 - \cos 2x) + \sin 2x]$

$$= \frac{1}{2} + \frac{1}{\sqrt{2}} \left(\sin 2x \cdot \frac{1}{\sqrt{2}} - \cos 2x \cdot \frac{1}{\sqrt{2}} \right)$$

$$= \frac{1}{2} + \frac{1}{\sqrt{2}} \sin(2x - 45^\circ)$$

$$\therefore -1 \leq \sin(2x - 45^\circ) \leq 1$$

$$\therefore \frac{1}{2} - \frac{1}{\sqrt{2}} \leq k \leq \frac{1}{2} + \frac{1}{\sqrt{2}}$$

$$\Rightarrow \frac{1 - \sqrt{2}}{2} \leq k \leq \frac{1 + \sqrt{2}}{2}$$

14. (a) : $\tan^{-1} \left(\frac{x+1}{x+2} \right) = \tan^{-1} 1 - \tan^{-1} \left(\frac{x-1}{x-2} \right)$

$$\Rightarrow \frac{x+1}{x+2} = \frac{1 - \frac{x-1}{x-2}}{1 + \frac{x-1}{x-2}} \Rightarrow \frac{x+1}{x+2} = \frac{-1}{2x-3}$$

$$\Rightarrow 2x^2 = 1 \Rightarrow x = \pm \frac{1}{\sqrt{2}}$$

15. (b) : Putting $x = x' + p$ and $y = y' + q$ in given equation, it becomes

$$2(x' + p)^2 + 3(x' + p)(y' + q) + 4(y' + q)^2 + (x' + p) + 18(y' + q) + 25 = 0$$

On comparing with $2x^2 + 3xy + 4y^2 = 1$, we get

$$4p + 3q + 1 = 0 \quad \dots(i)$$

$$3p + 8q + 18 = 0 \quad \dots(ii)$$

$$\text{and } 2p^2 + 3pq + 4q^2 + p + 18q + 25 = -1 \quad \dots(iii)$$

On comparing (i) & (ii), $p = 2, q = -3$ by which (iii) is also satisfied.

16. (a) : Centroid $\equiv \left(\frac{2-2+h}{3}, \frac{-3+1+k}{3} \right)$

i.e., $\left(\frac{h}{3}, \frac{k-2}{3} \right)$

lies on $2x + 3y = 1$

$$\therefore 2 \left(\frac{h}{3} \right) + 3 \left(\frac{k-2}{3} \right) = 1$$

$$\Rightarrow 2h + 3k - 6 = 3$$

$$\Rightarrow 2x + 3y = 9 \text{ is the reqd. locus.}$$

17. (d) : $P(3, 6)$ has reflection on $y = x$ as $Q(6, 3)$

Again reflection of $Q(6, 3)$ on $y = -x$ will be $Q'(-3, -6)$

$$\therefore \text{Slope of } PQ \times \text{slope of } QQ' = (-1)(1) = -1$$

$$\therefore PQQ' \text{ is a right angled } \Delta \text{ with } \angle PQQ' = 90^\circ$$

\therefore Circumcentre will be mid pt. of hypotenuse PQ'
i.e. $(0, 0)$

18. (b) : \therefore Bisectors of $3x + 4y = 5$ & $12x + 5y = 7$

$$\text{are } \frac{3x+4y-5}{\sqrt{3^2+4^2}} = \pm \frac{12x+5y-7}{\sqrt{12^2+5^2}}$$

$$\Rightarrow 13(3x + 4y - 5) = \pm 5(12x + 5y - 7)$$

$$\Rightarrow 21x - 27y + 30 = 0 \text{ \& } 99x + 77y = 100$$

$$\Rightarrow 7x - 9y + 10 = 0 \text{ \& } 99x + 77y = 100$$

$$\therefore 7x - 9y + 10 = 0 \text{ is one of the bisector.}$$

\therefore Perp. from any pt. on it will be equal to both given lines $\therefore d_1 = d_2$.

19. (d) : Equation of common chord i.e., $S_1 - S_2 = 0$ will be

$$\Rightarrow (x^2 + y^2 - 16) - (x^2 + y^2 - 4x - 4y) = 0$$

$$\Rightarrow x + y = 4, \text{ which subtends } 90^\circ \text{ at } (0, 0)$$

20. (c) : Let $M(h, k)$ be the mid point of chord


$$(x+1)^2 + (y-1)^2 = (2)^2 \quad \dots(i)$$

subtending 90° at centre $C(-1, 1)$

$$\therefore 2CM^2 = CB^2$$

$$\Rightarrow (h+1)^2 + (k-1)^2 = (2)^2$$

$$\Rightarrow x^2 + y^2 + 2x - 2y = 0 \text{ is the reqd. locus.}$$



21. (b) : $CS = ae, SP = \frac{b \cdot ae - 0}{\sqrt{b^2 + a^2}} = \frac{abe}{\sqrt{b^2 + a^2}}$

$$\therefore CP = \sqrt{CS^2 - SP^2}$$


$$= \sqrt{a^2 e^2 - \frac{b^2 a^2 e^2}{a^2 + b^2}}$$

$$= \frac{a^2 e}{\sqrt{a^2 + b^2}}$$

\therefore Area of rectangle with sides SP and $CP = SP \cdot CP$

$$= \frac{ab \cdot a^2 e^2}{a^2 + b^2} = ab$$

$$[\because b^2 = a^2(e^2 - 1)]$$



22. (b) : $CB = b, CS = ae$


$$\therefore \angle SBS' = 90^\circ$$

$$\therefore CB = CS$$

$$\Rightarrow b = ae \Rightarrow b^2 = a^2 e^2$$

$$\Rightarrow a^2(1 - e^2) = a^2 e^2$$

$$\Rightarrow 2e^2 = 1 \therefore e = \frac{1}{\sqrt{2}}$$



23. (a) : Parabola is $(x + y)^2 = 5(x - y + 1)$ whose axis is $x + y = 0$.

24. (c) : Foci of hyperbola $y^2 - x^2 = 1$ are

$$(0, \pm be) \text{ i.e., } (0, \pm \sqrt{2})$$

\therefore End points of a diameter of reqd. circle are $(0, \sqrt{2})$ and $(0, -\sqrt{2})$.

\therefore Eqn. of reqd. circle is

$$(x-0)(x-0) + (y-\sqrt{2})(y+\sqrt{2}) = 0$$

$$\text{or } x^2 + y^2 = 2$$

25. (d) : Any plane through $(1, 2, -3)$ is given by

$$a(x-1) + b(y-2) + c(z+3) = 0 \quad \dots(i)$$

If (i) is || to x-axis then $a = 0$

If (i) passes through $(2, -2, 1)$ then

$$b(-2 - 2) + c(1 + 3) = 0 \Rightarrow b = c$$

\therefore (i) becomes $b(y - 2) + b(z + 3) = 0$

$$\Rightarrow y + z + 1 = 0$$

26. (b) : Here, $\begin{vmatrix} 1 & -1 & 1 \\ 2 & -3 & 0 \\ 1 & 0 & 3 \end{vmatrix} = 1(-9 - 0) + 1(6 - 0) + 1(0 + 3) = -9 + 6 + 3 = 0$

\therefore 3 lines with d.r.s $(1, -1, 1)$, $(2, -3, 0)$ & $(1, 0, 3)$ are coplanar.

27. (a) : Given, $f(x)$ is non-constant & differentiable

s.t. $\frac{f(x)}{f(y)} = f(x - y) \quad \dots(i)$

Let $f(x) = a^{mx}$ satisfying (i). Now, $f'(x) = ma^{mx}$

Given $f'(0) = p \Rightarrow ma^0 = p \Rightarrow m = p \quad \dots(ii)$

Also, $f'(5) = q \Rightarrow ma^{5m} = q$

$$\Rightarrow a^{5m} = \frac{q}{m} = \frac{q}{p} \quad \dots(iii)$$

$$\therefore f'(-5) = ma^{m(-5)} = p \cdot a^{-5m} = p \cdot \frac{p}{q} \quad [\text{using (iii)}]$$

$$= \frac{p^2}{q}$$

28. (c) : $f(x) = \log_5 \log_3 x = \log_e (\log_3 x) \log_5 e$
 $= \log_5 e \log_e (\log_3 x)$

$$f'(x) = \log_5 e \cdot \frac{1}{\log_3 x} \cdot \frac{d}{dx} (\log_3 x)$$

$$= \log_5 e \cdot \frac{1}{\log_3 x} \cdot \frac{d}{dx} (\log_e x \cdot \log_3 e)$$

$$= \log_5 e \cdot \frac{1}{\log_3 x} \cdot \log_3 e \cdot \frac{1}{x} = \log_5 e \log_x e \cdot \frac{1}{x}$$

$$\therefore f'(e) = \log_5 e \log_e e \cdot \frac{1}{e} = \frac{1}{e \log_e 5}$$

29. (c) : $H(x) = G(F(x)) = G(e^x) = e^{-e^x}$

$$\therefore \frac{dH}{dx} = e^{-e^x} \cdot \frac{d}{dx} (-e^x) = e^{-e^x} \cdot (-e^x)$$

$$\therefore \left[\frac{dH}{dx} \right]_{x=0} = e^{-e^0} (-e^0) = -e^{-1} = -\frac{1}{e}$$

30. (c) : $\lim_{x \rightarrow 0} \frac{2f(x) - 3f(2x) + f(4x)}{x^2} \quad \left(\frac{0}{0} \text{ form} \right)$

$$= \lim_{x \rightarrow 0} \frac{2f'(x) - 6f'(2x) + 4f'(4x)}{2x} \quad \left(\frac{0}{0} \text{ form} \right)$$

$$= \lim_{x \rightarrow 0} \frac{2f''(x) - 12f''(2x) + 16f''(4x)}{2}$$

$$= f''(0) - 6f''(0) + 8f''(0)$$

$$= 3f''(0) = 3k$$

31. (a) : We have, $y = e^{m \sin^{-1} x}$

$$\Rightarrow \frac{dy}{dx} = e^{m \sin^{-1} x} \cdot m \cdot \frac{1}{\sqrt{1-x^2}} = \frac{my}{\sqrt{1-x^2}}$$

$$\Rightarrow (1-x^2) \left(\frac{dy}{dx} \right)^2 = m^2 y^2$$

$$\Rightarrow (1-x^2) \cdot 2 \frac{dy}{dx} \cdot \frac{d^2 y}{dx^2} - 2x \cdot \left(\frac{dy}{dx} \right)^2 = m^2 \cdot 2y \cdot \frac{dy}{dx}$$

$$\Rightarrow (1-x^2) \left(\frac{d^2 y}{dx^2} \right) - x \cdot \frac{dy}{dx} = m^2 y \Rightarrow k = m^2$$

32. (d) : Using mean value theorem, $f'(c) = \frac{f(b) - f(a)}{b - a}$

$$\Rightarrow 2c + 2a = \frac{f(\beta) - f(\alpha)}{\beta - \alpha}$$

$$= \frac{(\beta^2 + 2a\beta + b) - (\alpha^2 + 2a\alpha + b)}{\beta - \alpha} = \beta + \alpha + 2a$$

$$\therefore c = \frac{\alpha + \beta}{2}$$

33. (c) : $f(x) = x^{13} + x^{11} + x^9 + x^7 + x^5 + x^3 + x + 19$
 $f'(x) = 13x^{12} + 11x^{10} + 9x^8 + 7x^6 + 5x^4 + 3x^2 + 1 > 0 \quad \forall x \in \mathbb{R}$

$\therefore f(x)$ is a strictly increasing function.

$$f(-\infty) = -\infty, f(\infty) = \infty, f(0) = 19$$

$\therefore f(x) = 0$, will have only one real root

34. (b) : \therefore LMVT is applicable to $f(x)$ in $[0, x]$

$\therefore f(x)$ must be continuous in $[0, x]$

$$\therefore \lim_{x \rightarrow 0^+} f(x) = f(0) \Rightarrow \lim_{x \rightarrow 0^+} \frac{x^p}{(\sin x)^q} = 0$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{(0+h)^p}{\{\sin(0+h)\}^q} = 0 \Rightarrow \lim_{h \rightarrow 0} \frac{h^{p-q}}{\left(\frac{\sin h}{h} \right)^q} = 0$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{h^{p-q}}{1} = 0 \quad \therefore \text{L.H.L exists only if } p > q.$$

35. (b) : Let $A = \lim_{x \rightarrow 0} (\sin x)^{2 \tan x}$

$$\log A = \lim_{x \rightarrow 0} 2 \tan x \log(\sin x)$$

$$= 2 \lim_{x \rightarrow 0} \frac{\log(\sin x)}{\cot x} \quad \left(\frac{\infty}{\infty} \text{ form} \right)$$

$$= 2 \lim_{x \rightarrow 0} \frac{\frac{1}{\sin x} \cdot \cos x}{-\csc^2 x} = - \lim_{x \rightarrow 0} \sin 2x = 0 \Rightarrow A = 1$$

36. (c) : Let $I = \int \cos(\log x) dx$

$$= \cos(\log x) \cdot (x) - \int -\sin(\log x) \cdot \frac{1}{x} \cdot (x) dx$$

$$\begin{aligned}
&= x \cos(\log x) + \int \sin(\log x) dx \\
&= x \cos(\log x) + \sin(\log x)(x) - \int \cos(\log x) \cdot \frac{1}{x}(x) dx \\
\Rightarrow 2I &= x[\cos(\log x) + \sin(\log x)] \\
\Rightarrow I &= \frac{x}{2} [\cos(\log x) + \sin(\log x)] + c
\end{aligned}$$

$$\begin{aligned}
37. (a) : \int \frac{x^2 \left(1 - \frac{1}{x^2}\right) dx}{x^2 \left(x^2 + \frac{1}{x^2} + 3\right)} &= \int \frac{\left(1 - \frac{1}{x^2}\right) dx}{\left(x + \frac{1}{x}\right)^2 + 1} \\
&= \int \frac{dz}{z^2 + 1}, \text{ where } z = x + \frac{1}{x} \\
&= \tan^{-1} z + c = \tan^{-1} \left(x + \frac{1}{x}\right) + c
\end{aligned}$$

$$\begin{aligned}
38. (b) : \because I &= \int_{10}^{19} \frac{\sin x}{1+x^8} dx \\
\therefore |I| &= \left| \int_{10}^{19} \frac{\sin x}{1+x^8} dx \right| \leq \int_{10}^{19} \left| \frac{\sin x}{1+x^8} \right| dx \\
\Rightarrow |I| &\leq \int_{10}^{19} \frac{|\sin x|}{1+x^8} dx \leq \int_{10}^{19} \frac{dx}{1+x^8} \quad [\because |\sin x| \leq 1] \\
\Rightarrow |I| &\leq \int_{10}^{19} \frac{dx}{x^8} = \left[\frac{x^{-7}}{-7} \right]_{10}^{19} \\
&= \frac{1}{7} (10^{-7} - 19^{-7}) < \frac{10^{-7}}{7} < 10^{-7}
\end{aligned}$$

$$\begin{aligned}
39. (d) : I_2 &= \int_0^n \{x\} dx = \int_0^{n-1} \{x\} dx + \int_{n-1}^n \{x\} dx \\
&[\because \{x\} \text{ is periodic with period } = 1] \\
&= n \int_0^1 x dx = \frac{n}{2}
\end{aligned}$$

$$I_1 + I_2 = \int_0^n ([x] + \{x\}) dx = \int_0^n x dx = \left[\frac{x^2}{2} \right]_0^n = \frac{n^2}{2}$$

$$\therefore I_1 = \frac{n^2}{2} - \frac{n}{2} \quad \therefore \frac{I_1}{I_2} = \frac{n^2 - n}{n} = n - 1$$

$$\begin{aligned}
40. (b) : \lim_{n \rightarrow \infty} \left[\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \dots + \frac{n}{n^2 + n^2} \right] \\
= \lim_{n \rightarrow \infty} \frac{1}{n} \left[\frac{1}{1 + \left(\frac{1}{n}\right)^2} + \frac{1}{1 + \left(\frac{2}{n}\right)^2} + \dots + \frac{1}{1 + \left(\frac{n}{n}\right)^2} \right] \\
= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^n \frac{1}{1 + \left(\frac{r}{n}\right)^2} = \int_0^1 \frac{dx}{1+x^2} = [\tan^{-1} x]_0^1 = \frac{\pi}{4}
\end{aligned}$$

$$41. (d) : \because 0 \leq x \leq 1 \quad \therefore 1 \leq e^{x^2} \leq e$$

$$\Rightarrow \int_0^1 dx \leq \int_0^1 e^{x^2} dx \leq \int_0^1 e \cdot dx$$

$$\Rightarrow 1 \leq \int_0^1 e^{x^2} dx \leq e$$

$$\begin{aligned}
42. (c) : \int_0^{100(1)} e^{x-[x]} dx &= \int_0^{100(1)} e^{\{x\}} dx = 100 \int_0^1 e^{\{x\}} dx \\
&[\because \{x\} \text{ is periodic with period } 1] \\
&= 100 \int_0^1 e^x dx = 100(e - 1)
\end{aligned}$$

$$43. (a) : \text{Putting } x + y = z \Rightarrow 1 + \frac{dy}{dx} = \frac{dz}{dx}$$

$$\text{Given eq. becomes } z^2 \cdot \left(\frac{dz}{dx} - 1 \right) = a^2$$

$$\Rightarrow \frac{dz}{dx} = 1 + \frac{a^2}{z^2}$$

$$\Rightarrow \int \frac{z^2}{z^2 + a^2} dz = \int dx \Rightarrow \int \frac{(z^2 + a^2) - a^2}{z^2 + a^2} dz = x$$

$$\Rightarrow z - a^2 \cdot \frac{1}{a} \tan^{-1} \frac{z}{a} = x - c$$

$$\Rightarrow x + y - a \tan^{-1} \left(\frac{x+y}{a} \right) = x - c$$

$$\Rightarrow \frac{y+c}{a} = \tan^{-1} \left(\frac{x+y}{a} \right) \Rightarrow \frac{x+y}{a} = \tan \left(\frac{y+c}{a} \right),$$

c is an arbitrary constant.

$$44. (b) : \text{We have, } \frac{dy}{dx} + \frac{x(x^2+1)}{x^2(x^2-1)} \cdot y = \frac{x^2-1}{x^2(x^2-1)}$$

$$\Rightarrow \frac{dy}{dx} + \frac{x^2+1}{x(x^2-1)} \cdot y = \frac{1}{x^2}$$

$$\therefore \text{I.F.} = e^{\int \frac{x^2+1}{x(x^2-1)} dx} = e^{\int \frac{\left(1 + \frac{1}{x^2}\right)}{x-1} dx} = e^{\log_e \left(x - \frac{1}{x}\right)} = x - \frac{1}{x}$$

$$45. (b) : \text{Let G.P. be } a, ar, ar^2, \dots$$

$$\text{A.T.Q., } a = ar + ar^2 \Rightarrow r^2 + r - 1 = 0$$

$$\Rightarrow r = \frac{-1 \pm \sqrt{1+4}}{2} = \frac{-1 \pm \sqrt{5}}{2}$$

$$\because \text{Terms are positive } \therefore r > 0 \quad \therefore r = \frac{-1 + \sqrt{5}}{2}$$

$$46. (a) : \text{Using property of logarithm, } \log_3 y = 3 \log_x x = 3$$

$$\therefore y = 5^3 = 125$$

$$47. (c) : \left(\frac{1+i}{1-i} \right)^n (1-i)^2 = \left\{ \frac{(1+i)^2}{1-i^2} \right\}^n (1+i^2-2i) \\ = \left\{ \frac{1-1+2i}{2} \right\}^n (-2i) = -2i^{n+1}$$

$$48. (d) : \text{Let } \frac{z+i}{z-i} = ki \quad (k \in R)$$

By componendo & dividendo, we have $\frac{2z}{2i} = \frac{ki+1}{ki-1}$

$$\Rightarrow z = \frac{ki+1}{ki-1}, i \Rightarrow |z| = \frac{\sqrt{k^2+1^2}}{\sqrt{k^2+(-1)^2}} \cdot |i| = 1$$

$$\Rightarrow x^2 + y^2 = 1 \text{ which represents a circle}$$

$$49. (a) : 2px^2 + 2px + qx + q = 0$$

$$\Rightarrow 2px(x+1) + q(x+1) = 0$$

$\therefore x = -1, \frac{-q}{2p}$, which are rational as p & q are odd integers.

50. (b) : 3 out of 7 consonants can be chosen in 7C_3 ways and 12 out of 4 vowels can be chosen in 4C_2 ways
 \therefore Total no. of words that can be formed = ${}^7C_3 \times {}^4C_2 \times \underline{5}$
 $= 25200$

$$51. (b) : A^2 = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}$$

Only (b) is satisfied by putting $n = 2$

$$52. (d) : D_1 = \begin{vmatrix} a & a+1 & a-1 \\ -b & b+1 & b-1 \\ c & c-1 & c+1 \end{vmatrix} = \begin{vmatrix} a & -b & c \\ a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \end{vmatrix}$$

[Interchanging rows & columns]

$$= (-1)^2 \begin{vmatrix} a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \\ a & -b & c \end{vmatrix} = \begin{vmatrix} a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \\ a & -b & c \end{vmatrix}$$

$$D_2 = \begin{vmatrix} a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \\ (-1)^{n+2}a & (-1)^{n+1}b & (-1)^n c \end{vmatrix}$$

$$= (-1)^n \begin{vmatrix} a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \\ a & -b & c \end{vmatrix}$$

$\therefore D_1 + D_2 = 0$ is possible only when n is any odd integer.

53. (c) : $R \cup S = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1), (1, 3), (3, 1)\}$
 $A = \{1, 2, 3\}$

(i) $\therefore (1, 1), (2, 2), (3, 3) \in R \cup S \Rightarrow$ Reflexive

(ii) $(a, b) \in R \cup S$

$\Rightarrow (b, a) \in R \cup S \forall a, b \in \{1, 2, 3\} \Rightarrow$ Symmetric

(iii) $\therefore (2, 1) \& (1, 3) \in R \cup S$ but $(2, 3) \notin R \cup S$

\Rightarrow Not transitive

54. (a) : Let radius of circle with centre $(1, 1)$ be a

Its eqn. is $(x-1)^2 + (y-1)^2 = a^2$... (i)

Given circle is $x^2 + y^2 - 4x - 6y + 9 = 0$... (ii)

Eqn. of common chord ($S_1 - S_2 = 0$) is

$$2x + 4y = a^2 + 7 \quad \dots (iii)$$

If (iii) be a diameter of (ii) then centre $(2, 3)$ will lie on (iii)

$$\Rightarrow 4 + 12 = a^2 + 7 \Rightarrow a^2 = 9$$

\Rightarrow Radius = 3 units

$$55. (d) : \tan \theta = \frac{k}{h+1} \& \tan 2\theta = \frac{k}{2-h}$$

$$\Rightarrow \frac{k}{2-h} = \frac{2 \tan \theta}{1 - \tan^2 \theta} = \frac{2 \left(\frac{k}{h+1} \right)}{1 - \left(\frac{k}{h+1} \right)^2}$$

$$= \frac{2k(h+1)}{(h+1)^2 - k^2}$$

$$\Rightarrow (h+1)^2 - k^2 = 2(2-h)(h+1)$$

$$\Rightarrow h^2 + 2h + 1 - k^2 = 4h + 4 - 2h^2 - 2h$$

$$\Rightarrow 3h^2 - k^2 = 3 \Rightarrow \frac{h^2}{1} - \frac{k^2}{3} = 1$$

$\therefore M$ moves on a hyperbola


$$56. (c) : f(x) = \int_{-1}^x |t| dt = \int_{-1}^0 |t| dt + \int_0^x |t| dt \\ = \int_{-1}^0 |x| dx + \int_0^x |x| dx = \int_{-1}^0 (-x) dx + \int_0^x x dx \\ = \left[-\frac{x^2}{2} \right]_{-1}^0 + \left[\frac{x^2}{2} \right]_0^x = -\frac{1}{2}(0-1) + \left(\frac{x^2}{2} - 0 \right) = \frac{x^2+1}{2}$$

$$57. (b) : f(x) = \lim_{n \rightarrow \infty} n \left(x^{\frac{1}{n}} - 1 \right) = \lim_{k \rightarrow 0} \frac{x^k - 1}{k}, \text{ where } n = \frac{1}{k} \\ = \log_e x$$

$$\text{Now, } f(x) + f\left(\frac{1}{x}\right) = \log_e x + \log_e 1/x = \log_e 1 = 0$$


$$\therefore f(xy) = \log_e xy = \log_e x + \log_e y = f(x) + f(y)$$

$$58. (b) : I = \int_0^{100\pi} \sqrt{2 \sin^2 x} dx$$



$$\begin{aligned}
 &= \sqrt{2} \int_0^{100\pi} |\sin x| dx = \sqrt{2} \cdot 100 \int_0^{\pi} |\sin x| dx \\
 &\quad [\because \text{Period of } |\sin x| \text{ is } \pi] \\
 &= 100\sqrt{2} \int_0^{\pi} \sin x dx = 100\sqrt{2} [-\cos x]_0^{\pi} \\
 &= 100\sqrt{2}(1+1) = 200\sqrt{2}
 \end{aligned}$$

59. (a) :



Parabolas are $y^2 = -\frac{x}{2}$... (i) & $y^2 = -\frac{1}{3}(x-1)$... (ii)

On solving, $-\frac{x}{2} = -\frac{(x-1)}{3}$

$\Rightarrow -2x + 2 = -3x \Rightarrow x = -2$

$\therefore \text{Reqd. area} = 2 \left[\int_{-2}^0 y_2 dx - \int_{-2}^0 y_1 dx \right]$

$[y_1, y_2 \text{ are values of } y \text{ from (i) \& (ii) resp.}]$

$$= 2 \left[\int_{-2}^0 \sqrt{\frac{1-x}{3}} dx - \int_{-2}^0 \sqrt{-\frac{x}{2}} dx \right]$$

$$= \frac{2}{\sqrt{3}} \left[\frac{(1-x)^{3/2}}{\left(-\frac{3}{2}\right)} \right]_{-2}^0 - \frac{2}{\sqrt{2}} \left[\frac{(-x)^{3/2}}{\left(-\frac{3}{2}\right)} \right]_{-2}^0$$

$$= -\frac{4}{3\sqrt{3}} [0 - 3^{3/2}] + \frac{4}{3\sqrt{2}} [0 - 2^{3/2}]$$

$$= 4 - \frac{8}{3} = \frac{4}{3} \text{ square units}$$

60. (a) : We have,


$$\frac{x^2}{9} + \frac{y^2}{5} = 1 \quad \dots (i)$$

$$e = \sqrt{1 - \frac{5}{9}} = \frac{2}{3}$$

$$\therefore L \left(ae, \frac{b^2}{a} \right) \equiv \left(2, \frac{5}{3} \right)$$

Eqn. of tangent to (i) at $L \left(2, \frac{5}{3} \right)$ is

$$\frac{x}{9} \cdot 2 + \frac{y}{5} \cdot \frac{5}{3} = 1 \Rightarrow \frac{x}{9/2} + \frac{y}{3} = 1$$



$$\therefore CA = \frac{9}{2}, CB = 3$$

$\therefore \text{Reqd. area of quad.} = 4 \times \text{Area of } \Delta CAB$

$$= 4 \times \frac{1}{2} \cdot \frac{9}{2} \cdot 3 = 27 \text{ sq. units.}$$

61. (c) : $f(x) = \sin x - \cos x - Kx + 5$

$$\Rightarrow f'(x) = \cos x + \sin x - K = \sqrt{2} \sin \left(x + \frac{\pi}{4} \right) - K$$

$$\therefore \sqrt{2} \sin \left(x + \frac{\pi}{4} \right) \leq \sqrt{2}$$

$\therefore f(x)$ will be decreasing for all +ve real x if $f'(x) < 0$

$$\sqrt{2} \sin \left(x + \frac{\pi}{4} \right) - K < 0$$

$$\Rightarrow K > \sqrt{2} \sin \left(x + \frac{\pi}{4} \right) \Rightarrow K > \sqrt{2}$$

62. (b) : Let $\vec{x} = a\hat{i} + b\hat{j} + c\hat{k}$

$$\therefore \vec{x} \times \hat{i} = -b\hat{k} + c\hat{j}$$

$$(\vec{x} \times \hat{i})^2 = |\vec{x} \times \hat{i}|^2 = b^2 + c^2$$

Similarly, $(\vec{x} \times \hat{j})^2 = c^2 + a^2$ & $(\vec{x} \times \hat{k})^2 = a^2 + b^2$

$$\therefore (\vec{x} \times \hat{i})^2 + (\vec{x} \times \hat{j})^2 + (\vec{x} \times \hat{k})^2 = 2(a^2 + b^2 + c^2) = 2|\vec{x}|^2$$

63. (c) : Let $\vec{a}, \vec{b}, \vec{c}$ be three unit vectors such that

$$\vec{c} = \vec{a} + \vec{b}$$

$$\therefore \vec{c}^2 = (\vec{a} + \vec{b})^2 \Rightarrow 1 = 1 + 1 + 2\vec{a} \cdot \vec{b}$$

$$\Rightarrow 2\vec{a} \cdot \vec{b} = -1$$

Now, $|\vec{a} - \vec{b}|^2 = 1 + 1 - 2\vec{a} \cdot \vec{b} = 3 \Rightarrow |\vec{a} - \vec{b}| = \sqrt{3}$ units.

64. (a) : \because Roots of $x^2 + x + 1 = 0$ are ω & ω^2

$$\text{Let } \alpha = \omega = \frac{-1 + \sqrt{3}i}{2} = \cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}$$

$$\text{and } \beta = \omega^2 = \frac{-1 - \sqrt{3}i}{2} = \cos \left(-\frac{2\pi}{3} \right) + i \sin \left(-\frac{2\pi}{3} \right)$$

$$= \cos \frac{2\pi}{3} - i \sin \frac{2\pi}{3} \quad \therefore \alpha^n + \beta^n = 2 \cos \frac{2n\pi}{3}$$

65. (c) : Let $y = \frac{x^2 + 2x + 4}{2x^2 + 4x + 9}$

$$\Rightarrow (2y - 1)x^2 + 2(2y - 1)x + (9y - 4) = 0$$

$$\text{For real } x, D \geq 0 \Rightarrow 4(2y - 1)^2 - 4(2y - 1)(9y - 4) \geq 0$$

$$\Rightarrow 4y^2 - 4y + 1 - 18y^2 + 17y - 4 \geq 0 \Rightarrow 14y^2 - 13y + 3 \leq 0$$

$$\Rightarrow (7y - 3)(2y - 1) \leq 0 \Rightarrow \frac{3}{7} \leq y \leq \frac{1}{2}$$

$$\Rightarrow \text{Greatest value of } y \text{ i.e. } \frac{x^2 + 2x + 4}{2x^2 + 4x + 9} \text{ is } \frac{1}{2}$$

66. (c, d) : $\because a, b \in \{1, 2, 3\}$ & $ax^2 + bx + 1 = 0$ has real roots

$$\Rightarrow D \geq 0 \text{ i.e., } b^2 \geq 4a \quad \dots (i)$$

∴ Possible ordered pair (a, b) are $(1, 2)$, $(1, 3)$ & $(2, 3)$ only

67. (a, c) : Equation of tangent to $y^2 = 4ax$ at $(at^2, 2at)$ is $y \cdot 2at = 2a(x + at^2)$

$$\Rightarrow ty = x + at^2 \quad \dots(i)$$

$$\text{For } x^2 - y^2 = a^2, \frac{dy}{dx} = \frac{x}{y}$$

$$\Rightarrow \text{At } (a \sec \theta, a \tan \theta), \frac{dy}{dx} = \frac{a \sec \theta}{a \tan \theta} = \frac{1}{\sin \theta}$$

∴ Eqn. of normal to $x^2 - y^2 = a^2$ at $(a \sec \theta, a \tan \theta)$ is $y - a \tan \theta = -\sin \theta (x - a \sec \theta)$

$$\Rightarrow y = -x \sin \theta + 2a \tan \theta \quad \dots(ii)$$

∴ (i) & (ii) are identical

$$\therefore \frac{t}{1} = \frac{1}{-\sin \theta} = \frac{at^2}{2a \tan \theta} = \frac{t^2}{2 \tan \theta}$$

$$\Rightarrow t = -\operatorname{cosec} \theta \quad \& \quad t = 2 \tan \theta$$

68. (c) : Conic may be written as $(x - 3)^2 = -4(y - 2)$
 $X^2 = 4AY$...(i)

Here, $X = x - 3$, $Y = y - 2$ & $4A = -4$...(ii)

For focus, $X = 0$, $Y = A \Rightarrow x - 3 = 0$, $y - 2 = -1$

$\Rightarrow x = 3$, $y = 1$ ∴ Focus is $(3, 1)$

69. (b)

70. (b, c) : $f(x) = x^n$...(i) $f'(\alpha + \beta) = f'(\alpha) + f'(\beta)$...(ii)

When $n = 2$, $f(x) = x^2$, $f'(x) = 2x$

$$\therefore f'(\alpha + \beta) = 2(\alpha + \beta) = 2\alpha + 2\beta = f'(\alpha) + f'(\beta)$$

\Rightarrow (ii) is satisfied

When $n = 0$, $f(x) = 1 \Rightarrow f'(x) = 0 \Rightarrow$ (ii) is satisfied.

(ii) is not satisfied if $n = 1$ or 5

$$\mathbf{71. (c) : I = \int_0^a \frac{dx}{1 + f(x)} \quad \dots(i)$$

$$= \int_0^a \frac{dx}{1 + f(a-x)} = \int_0^a \frac{f(x)f(a-x)}{f(x)f(a-x) + f(a-x)} dx$$

$$I = \int_0^a \frac{f(x)}{f(x)+1} dx \quad \dots(ii)$$

$$\text{Adding (i) \& (ii), } 2I = \int_0^a dx = a \Rightarrow I = \frac{a}{2}$$

72. (b, d) : $ax + by + c = 0$...(i)

$$\text{Its slope} = \frac{-a}{b} \quad (ab \neq 0)$$

$$\text{Now, } xy = 1 - 2x \Rightarrow y + 2 = \frac{1}{x} \quad \dots(ii)$$

Differentiating (ii), we get, $\frac{dy}{dx} = -\frac{1}{x^2}$

$$\therefore \frac{dy}{dx} < 0$$

If (i) be a tangent to (ii) then $-\frac{a}{b} = -\frac{1}{x^2} < 0$

$$\Rightarrow \frac{b}{a} = x^2 > 0$$

∴ a & b must be of same sign

∴ $a > 0, b > 0$ or $a < 0, b < 0$

$$\mathbf{73. (b, c) : \frac{\vec{u}}{0} \cdot \frac{f}{f} = 0}$$

Dist. between them at any instant is given by

$$x = u \cdot t - \left\{ 0 \cdot t + \frac{1}{2} ft^2 \right\}$$

$$\frac{dx}{dt} = u - \frac{1}{2} \cdot f \cdot 2t = u - ft \Rightarrow \frac{d^2x}{dt^2} = -f < 0$$

For max. distance between them, $\frac{dx}{dt} = 0 \Rightarrow t = \frac{u}{f}$

$$\text{Max. distance} = u \cdot \frac{u}{f} - \frac{1}{2} f \cdot \frac{u^2}{f^2} = \frac{u^2}{f} - \frac{u^2}{2f} = \frac{u^2}{2f}$$

74. (a, c) : $|z - i| = |z + 1| = 1$

$$\Rightarrow x^2 + (y - 1)^2 = (x + 1)^2 + y^2 = 1 \quad \dots(i)$$

$$\Rightarrow -2y + 1 = 2x + 1 \Rightarrow y = -x \quad \dots(ii)$$

When $y = -x$, we have $x^2 + 2x + 1 + x^2 = 1$

$$\Rightarrow 2x(x + 1) = 0 \Rightarrow x = 0, -1$$

when $x = 0 \Rightarrow y = 0$

when $x = -1 \Rightarrow y = 1$

∴ $z = 0, -1 + i$

75. (c) : (i) ∴ $1 + a \cdot a = 1 + a^2 > 0 \Rightarrow$ Reflexive

(ii) If $1 + ab > 0$ then $1 + ba > 0 \Rightarrow$ Symmetric

$$\left. \begin{aligned} \text{(iii) } \therefore 1 + 1\left(\frac{1}{2}\right) &= \frac{3}{2} > 0 \Rightarrow \left(1, \frac{1}{2}\right) \in \rho \\ 1 + \frac{1}{2}(-1) &= \frac{1}{2} > 0 \Rightarrow \left(\frac{1}{2}, -1\right) \in \rho \end{aligned} \right\}$$

But, $1 + (1)(-1) = 0 \not> 0$

$\Rightarrow (1, -1) \notin \rho$

∴ ρ is not transitive.

Your favourite MTG Books/Magazines available in MANIPUR at

- Fame Book House - Imphal Mob: 9774220848, 7085556697
- Jain Book Shop - Imphal Mob: 9856031157
- Job Centre - Imphal Ph: 0385-2440140; Mob: 9856700700, 9436204282
- P.C. Jain And Co. Ph: 0385-2451756, 2225004; Mob: 9856084649

Visit "MTG IN YOUR CITY" on www.mtg.in to locate nearest book seller OR write to info@mtg.in OR call

0124-6601200 for further assistance.

Kerala PET

1. $\begin{vmatrix} 1 & 1 & 1 \\ p & q & r \\ p & q & r+1 \end{vmatrix}$ is equal to
(a) $q - p$ (b) $q + p$ (c) q (d) p
(e) 0

2. Let $A = \begin{bmatrix} 5 & 0 \\ 1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$.

If $4A + 5B - C = O$, then C is

(a) $\begin{bmatrix} 5 & 25 \\ -1 & 0 \end{bmatrix}$ (b) $\begin{bmatrix} 20 & 5 \\ -1 & 0 \end{bmatrix}$
(c) $\begin{bmatrix} 5 & -1 \\ 0 & 25 \end{bmatrix}$ (d) $\begin{bmatrix} 5 & 25 \\ -1 & 5 \end{bmatrix}$
(e) $\begin{bmatrix} 0 & 5 \\ 5 & 25 \end{bmatrix}$

3. If $U = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$, then U^{-1} is

(a) U^T (b) U (c) I (d) O
(e) U^2

4. If $A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}$, then A^{-1} is

(a) A^T (b) A^2 (c) A (d) I
(e) O

5. If $\begin{pmatrix} x+y & x-y \\ 2x+z & x+z \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}$, then the values of x, y and z are respectively

(a) 0, 0, 1 (b) 1, 1, 0 (c) -1, 0, 0 (d) 0, 0, 0
(e) 1, 1, 1

6. $\begin{pmatrix} 7 & 1 & 5 \\ 8 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} + 5 \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ is equal to

(a) $\begin{pmatrix} 16 \\ 27 \end{pmatrix}$ (b) $\begin{pmatrix} 27 \\ 16 \end{pmatrix}$ (c) $\begin{pmatrix} 15 \\ 16 \end{pmatrix}$ (d) $\begin{pmatrix} 16 \\ 15 \end{pmatrix}$
(e) $\begin{pmatrix} 16 \\ 16 \end{pmatrix}$

7. If $\begin{pmatrix} 1 & 2 & 4 \\ 1 & 3 & 5 \\ 1 & 4 & a \end{pmatrix}$ is singular, then the value of a is

(a) $a = -6$ (b) $a = 5$
(c) $a = -5$ (d) $a = 6$
(e) $a = 0$

8. If $\begin{pmatrix} 1 & 2 & -3 \\ 0 & 4 & 5 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$, then (x, y, z) is equal to

(a) (1, 6, 6) (b) (1, -6, 1)
(c) (1, 1, 6) (d) (6, -1, 1)
(e) (-1, 6, 1)

9. If $A = \begin{pmatrix} 1 & 5 \\ 0 & 2 \end{pmatrix}$, then

(a) $A^2 - 2A + 2I = O$ (b) $A^2 - 3A + 2I = O$
(c) $A^2 - 5A + 2I = O$ (d) $2A^2 - A + I = O$
(e) $A^2 + 3A + 2I = O$

10. If $\begin{pmatrix} 2x+y & x+y \\ p-q & p+q \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$, then (x, y, p, q) equals

(a) 0, 1, 0, 0 (b) 0, -1, 0, 0
(c) 1, 0, 0, 0 (d) 0, 1, 0, 1
(e) 1, 0, 1, 0

11. The value of $\left| \sqrt{4+2\sqrt{3}} \right| - \left| \sqrt{4-2\sqrt{3}} \right|$ is
 (a) 1 (b) 2 (c) 4 (d) 3
 (e) 5
12. The value of $8^{2/3} - 16^{1/4} - 9^{1/2}$ is
 (a) -1 (b) -2 (c) -3 (d) -4
 (e) -5
13. Let $x = 2$ be a root of $y = 4x^2 - 14x + q = 0$. Then y is equal to
 (a) $(x - 2)(4x - 6)$ (b) $(x - 2)(4x + 6)$
 (c) $(x - 2)(-4x - 6)$ (d) $(x - 2)(-4x + 6)$
 (e) $(x - 2)(4x + 3)$
14. If x_1 and x_2 are the roots of $3x^2 - 2x - 6 = 0$, then $x_1^2 + x_2^2$ is equal to
 (a) $\frac{50}{9}$ (b) $\frac{40}{9}$ (c) $\frac{30}{9}$ (d) $\frac{20}{9}$
 (e) $\frac{10}{9}$
15. Let x_1 and x_2 be the roots of the equation $x^2 + px - 3 = 0$. If $x_1^2 + x_2^2 = 10$, then the value of p is equal to
 (a) -4 or 4 (b) -3 or 3
 (c) -2 or 2 (d) -1 or 1
 (e) 0
16. If the product of roots of the equation $mx^2 + 6x + (2m - 1) = 0$ is -1, then the value of m is
 (a) $\frac{1}{3}$ (b) 1 (c) 3 (d) -1
 (e) -3
17. If $f(x) = \frac{1}{x^2 + 4x + 4} - \frac{4}{x^4 + 4x^3 + 4x^2} + \frac{4}{x^3 + 2x^2}$, then $f\left(\frac{1}{2}\right)$ is equal to
 (a) 1 (b) 2 (c) -1 (d) 3
 (e) 4
18. If x and y are the roots of the equation $x^2 + bx + 1 = 0$, then the value of $\frac{1}{x+b} + \frac{1}{y+b}$ is
 (a) $\frac{1}{b}$ (b) b (c) $\frac{1}{2b}$ (d) $2b$
 (e) 1
19. The equations $x^5 + ax + 1 = 0$ and $x^6 + ax^2 + 1 = 0$ have a common root. Then a is equal to
 (a) -4 (b) -2 (c) -3 (d) -1
 (e) 0
20. The roots of $ax^2 + x + 1 = 0$, where $a \neq 0$, are in the ratio 1 : 1. Then a is equal to
 (a) $\frac{1}{4}$ (b) $\frac{1}{2}$ (c) $\frac{3}{4}$ (d) 1
 (e) 0
21. If $z^2 + z + 1 = 0$ where z is a complex number, then the value of $\left(z + \frac{1}{z}\right)^2 + \left(z^2 + \frac{1}{z^2}\right)^2 + \left(z^3 + \frac{1}{z^3}\right)^2$ equals
 (a) 4 (b) 5 (c) 6 (d) 7
 (e) 8
22. Let $\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & -1-w^2 & w^2 \\ 1 & w & w^4 \end{vmatrix}$, where $w \neq 1$ is a complex number such that $w^3 = 1$. Then Δ equals
 (a) $3w + w^2$ (b) $3w^2$
 (c) $3(w - w^2)$ (d) $-3w^2$
 (e) $3w^2 + 1$
23. If $\begin{vmatrix} 3i & -9i & 1 \\ 2 & 9i & -1 \\ 10 & 9 & i \end{vmatrix} = x + iy$, then
 (a) $x = 1, y = 1$ (b) $x = 0, y = 1$
 (c) $x = 1, y = 0$ (d) $x = 0, y = 0$
 (e) $x = -1, y = 0$
24. If $z = \cos\left(\frac{\pi}{3}\right) - i \sin\left(\frac{\pi}{3}\right)$, then $z^2 - z + 1$ is equal to
 (a) 0 (b) 1 (c) -1 (d) $\frac{\pi}{2}$
 (e) π
25. $\left(\frac{1 + \cos\left(\frac{\pi}{12}\right) + i \sin\left(\frac{\pi}{12}\right)}{1 + \cos\left(\frac{\pi}{12}\right) - i \sin\left(\frac{\pi}{12}\right)} \right)^{72}$ is equal to
 (a) 0 (b) -1 (c) 1 (d) $\frac{1}{2}$
 (e) $-\frac{1}{2}$
26. If $A = \begin{vmatrix} 4 & k & k \\ 0 & k & k \\ 0 & 0 & k \end{vmatrix}$ and $\det(A) = 256$, then $|k|$ equals
 (a) 4 (b) 5 (c) 6 (d) 7
 (e) 8

27. If $A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, then $A^n + nI$ is equal to
 (a) I (b) nA
 (c) $I + nA$ (d) $I - nA$
 (e) $nA - I$
28. If $|z| = 5$ and $w = \frac{z-5}{z+5}$, then $\text{Re}(w)$ is equal to
 (a) 0 (b) $\frac{1}{25}$ (c) 25 (d) 1
 (e) -1
29. If $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$, then A^{2017} is equal to
 (a) $2^{2015}A$ (b) $2^{2016}A$
 (c) $2^{2014}A$ (d) $2^{2017}A$
 (e) $2^{2020}A$
30. If $a = e^{i\theta}$, then $\frac{1+a}{1-a}$ is equal to
 (a) $\cot \frac{\theta}{2}$ (b) $\tan \theta$
 (c) $i \cot \frac{\theta}{2}$ (d) $i \tan \frac{\theta}{2}$
 (e) $2 \tan \theta$
31. Three numbers x , y and z are in arithmetic progression. If $x + y + z = -3$ and $xyz = 8$, then $x^2 + y^2 + z^2$ is equal to
 (a) 9 (b) 10 (c) 21 (d) 20
 (e) 1
32. The 30th term of the arithmetic progression 10, 7, 4, ... is
 (a) -90 (b) -87 (c) -77 (d) -67
 (e) -57
33. The arithmetic mean of two numbers x and y is 3 and geometric mean is 1. Then $x^2 + y^2$ is equal to
 (a) 30 (b) 31 (c) 32 (d) 33
 (e) 34
34. The solution of $3^{2x-1} = 81^{1-x}$ is
 (a) $\frac{2}{3}$ (b) $\frac{1}{6}$ (c) $\frac{7}{6}$ (d) $\frac{5}{6}$
 (e) $\frac{1}{3}$
35. The sixth term in the sequence is 3, 1, $\frac{1}{3}$, ... is
 (a) $\frac{1}{27}$ (b) $\frac{1}{9}$ (c) $\frac{1}{81}$ (d) $\frac{1}{17}$
 (e) $\frac{1}{7}$
36. Three numbers are in arithmetic progression. Their sum is 21 and the product of the first number and the third number is 45. Then the product of these three numbers is
 (a) 315 (b) 90 (c) 180 (d) 270
 (e) 450
37. If $a+1$, $2a+1$, $4a-1$ are in arithmetic progression, then the value of a is
 (a) 1 (b) 2 (c) 3 (d) 4
 (e) 5
38. Two numbers x and y have arithmetic mean 9 and geometric mean 4. Then x and y are the roots of
 (a) $x^2 - 18x - 16 = 0$ (b) $x^2 - 18x + 16 = 0$
 (c) $x^2 + 18x - 16 = 0$ (d) $x^2 + 18x + 16 = 0$
 (e) $x^2 - 17x + 16 = 0$
39. Three unbiased coins are tossed. The probability of getting at least 2 tails is
 (a) $\frac{3}{4}$ (b) $\frac{1}{4}$ (c) $\frac{1}{2}$ (d) $\frac{1}{3}$
 (e) $\frac{2}{3}$
40. A single letter is selected from the word TRICKS. The probability that it is either T or R is
 (a) $\frac{1}{36}$ (b) $\frac{1}{4}$ (c) $\frac{1}{2}$ (d) $\frac{2}{3}$
 (e) $\frac{1}{3}$
41. From 4 red balls, 2 white balls and 4 black balls, four balls are selected. The probability of getting 2 red balls is
 (a) $\frac{7}{21}$ (b) $\frac{8}{21}$ (c) $\frac{9}{21}$ (d) $\frac{10}{21}$
 (e) $\frac{11}{21}$
42. In a class, 60% of the students know lesson I, 40% know lesson II and 20% know lesson I and lesson II. A student is selected at random. The probability that the student does not know lesson I and lesson II is
 (a) 0 (b) $\frac{4}{5}$ (c) $\frac{3}{5}$ (d) $\frac{1}{5}$
 (e) $\frac{2}{5}$
43. Two distinct numbers x and y are chosen from 1, 2, 3, 4, 5. The probability that the arithmetic mean of x and y is an integer is

- (a) 0 (b) $\frac{1}{5}$ (c) $\frac{3}{5}$ (d) $\frac{2}{5}$
 (e) $\frac{4}{5}$
44. The number of 3×3 matrices with entries -1 or $+1$ is
 (a) 2^4 (b) 2^5 (c) 2^6 (d) 2^7
 (e) 2^9
45. Let S be the set of all 2×2 symmetric matrices whose entries are either zero or one. A matrix X is chosen from S . The probability that the determinant of X is not zero is
 (a) $\frac{1}{3}$ (b) $\frac{1}{2}$ (c) $\frac{3}{4}$ (d) $\frac{1}{4}$
 (e) $\frac{2}{9}$
46. The number of words that can be formed by using all the letters of the word PROBLEM only once is
 (a) $5!$ (b) $6!$ (c) $7!$ (d) $8!$
 (e) $9!$
47. The number of diagonals in a hexagon is
 (a) 8 (b) 9 (c) 10 (d) 11
 (e) 12
48. The sum of odd integers from 1 to 2001 is
 (a) 1001^2 (b) 1000^2 (c) 1002^2 (d) 1003^2
 (e) 999^2
49. Two balls are selected from two black and two red balls. The probability that the two balls will have no black ball is
 (a) $\frac{1}{7}$ (b) $\frac{1}{5}$ (c) $\frac{1}{4}$ (d) $\frac{1}{3}$
 (e) $\frac{1}{6}$
50. If $z = i^9 + i^{19}$, then z is equal to
 (a) $0 + 0i$ (b) $1 + 0i$ (c) $0 + i$ (d) $1 + 2i$
 (e) $1 + 3i$
51. The mean for the data 6, 7, 10, 12, 13, 4, 8, 12 is
 (a) 9 (b) 8 (c) 7 (d) 6
 (e) 5
52. The set of all real numbers satisfying the inequality $x - 2 < 1$ is
 (a) $(3, \infty)$ (b) $[3, \infty)$
 (c) $[-3, \infty)$ (d) $(-\infty, -3)$
 (e) $(-\infty, 3)$
53. If $\frac{|x-3|}{x-3} > 0$, then
 (a) $x \in (-3, \infty)$ (b) $x \in (3, \infty)$
 (c) $x \in (2, \infty)$ (d) $x \in (1, \infty)$
 (e) $x \in (-1, \infty)$
54. The mode of the data 8, 11, 9, 8, 11, 9, 7, 8, 7, 3, 2, 8 is
 (a) 11 (b) 9 (c) 8 (d) 3
 (e) 7
55. If the mean of six numbers is 41, then the sum of these numbers is
 (a) 246 (b) 236 (c) 226 (d) 216
 (e) 206
56. If $\int_0^x f(t)dt = x^2 + e^x (x > 0)$, then $f(1)$ is equal to
 (a) $1 + e$ (b) $2 + e$ (c) $3 + e$ (d) e
 (e) 0
57. $\int \frac{x+1}{x^{1/2}} dx =$
 (a) $-x^{3/2} + x^{1/2} + c$ (b) $x^{1/2}$
 (c) $\frac{2}{3}x^{3/2} + 2x^{1/2} + c$ (d) $x^{3/2} + x^{1/2} + c$
 (e) $x^{3/2}$
58. In a flight 50 people speak Hindi, 20 speak English and 10 speak both English and Hindi. The number of people who speak at least one of the two languages is
 (a) 40 (b) 50 (c) 20 (d) 80
 (e) 60
59. If $f(x) = \frac{x+1}{x-1}$, then the value of $f(f(x))$ is equal to
 (a) x (b) 0 (c) $-x$ (d) 1
 (e) 2
60. Two dice are thrown simultaneously. What is the probability of getting two numbers whose product is even?
 (a) $\frac{3}{4}$ (b) $\frac{1}{4}$ (c) $\frac{1}{2}$ (d) $\frac{2}{3}$
 (e) $\frac{1}{16}$
61. $\lim_{x \rightarrow 0} \frac{\sqrt{2+x} - \sqrt{2-x}}{x}$ is equal to
 (a) $\frac{1}{\sqrt{2}}$ (b) $\sqrt{2}$
 (c) 0 (d) Does not exist
 (e) $\frac{1}{2\sqrt{2}}$

62. $\int \frac{dx}{e^x + e^{-x} + 2}$ is equal to
 (a) $\frac{1}{e^x + 1} + c$ (b) $\frac{-1}{e^x + 1} + c$
 (c) $\frac{1}{1 + e^{-x}} + c$ (d) $\frac{1}{e^{-x} - 1} + c$
 (e) $\frac{1}{e^x - 1} + c$
63. $\tan\left(\frac{\pi}{4} + \frac{\theta}{2}\right) + \tan\left(\frac{\pi}{4} - \frac{\theta}{2}\right)$ is equal to
 (a) $\sec \theta$ (b) $2 \sec \theta$ (c) $\sec \frac{\theta}{2}$ (d) $\sin \theta$
 (e) $\cos \theta$
64. $\int_{-1}^0 \frac{dx}{x^2 + x + 2}$ is equal to
 (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{2}$ (c) π (d) 0
 (e) $-\pi$
65. $\int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$ is equal to
 (a) 0 (b) $-\pi$ (c) $\frac{3\pi}{2}$ (d) $\frac{\pi}{2}$
 (e) $\frac{\pi}{4}$
66. If (x, y) is equidistant from $(a + b, b - a)$ and $(a - b, a + b)$, then
 (a) $x + y = 0$ (b) $bx - ay = 0$
 (c) $ax - by = 0$ (d) $bx + ay = 0$
 (e) $ax + by = 0$
67. If the points $(1, 0)$, $(0, 1)$ and $(x, 8)$ are collinear, then the value of x is equal to
 (a) 5 (b) -6 (c) 6 (d) 7
 (e) -7
68. The minimum value of the function $\max\{x, x^2\}$ is equal to
 (a) 0 (b) 1 (c) 2 (d) $\frac{1}{2}$
 (e) $\frac{3}{2}$
69. Let $f(x + y) = f(x)f(y)$ for all x and y . If $f(0) = 1$, $f(3) = 3$ and $f'(0) = 11$, then $f'(3)$ is equal to
 (a) 11 (b) 22 (c) 33 (d) 44
 (e) 55
70. If $f(9) = f'(9) = 0$, then $\lim_{x \rightarrow 9} \frac{\sqrt{f(x)} - 3}{\sqrt{x} - 3}$ is equal to
 (a) 0 (b) $f(0)$ (c) $f'(3)$ (d) $f(9)$
 (e) 1
71. The value of $\cos\left(\frac{\pi}{4} + x\right) + \cos\left(\frac{\pi}{4} - x\right)$ is
 (a) $\sqrt{2} \sin^2 x$ (b) $\sqrt{2} \sin x$
 (c) $\sqrt{2} \cos^2 x$ (d) $\sqrt{3} \cos x$
 (e) $\sqrt{2} \cos x$
72. Area of the triangle with vertices $(-2, 2)$, $(1, 5)$ and $(6, -1)$ is
 (a) 15 (b) $\frac{3}{5}$ (c) $\frac{29}{2}$ (d) $\frac{33}{2}$
 (e) $\frac{35}{2}$
73. The equation of the line passing through $(-3, 5)$ and perpendicular to the line through the points $(1, 0)$ and $(-4, 1)$ is
 (a) $5x + y + 10 = 0$ (b) $5x - y + 20 = 0$
 (c) $5x - y - 10 = 0$ (d) $5x + y + 20 = 0$
 (e) $5y - x - 10 = 0$
74. The coefficient of x^5 in the expansion of $(1 + x^2)^5 (1 + x)^4$ is
 (a) 30 (b) 60 (c) 40 (d) 10
 (e) 45
75. The coefficient of x^4 in the expansion of $(1 - 2x)^5$ is equal to
 (a) 40 (b) 320 (c) -320 (d) -32
 (e) 80
76. The equation $5x^2 + y^2 + y = 8$ represents
 (a) an ellipse (b) a parabola
 (c) a hyperbola (d) a circle
 (e) a straight line
77. The centre of the ellipse $4x^2 + y^2 - 8x + 4y - 8 = 0$ is
 (a) $(0, 2)$ (b) $(2, -1)$ (c) $(2, 1)$ (d) $(1, 2)$
 (e) $(1, -2)$
78. The area bounded by the curves $y = -x^2 + 3$ and $y = 0$ is
 (a) $\sqrt{3} + 1$ (b) $\sqrt{3}$
 (c) $4\sqrt{3}$ (d) $5\sqrt{3}$
 (e) $6\sqrt{3}$
79. The order of the differential equation $\left(\frac{d^3 y}{dx^3}\right)^2 + \left(\frac{d^2 y}{dx^2}\right)^2 + \left(\frac{dy}{dx}\right)^5 = 0$ is

- (a) 3 (b) 4 (c) 1 (d) 5
(e) 6
80. If $f(x) = \sqrt{2x} + \frac{4}{\sqrt{2x}}$, then $f'(2)$ is equal to
(a) 0 (b) -1 (c) 1 (d) 2
(e) -2
81. The area of the circle $x^2 - 2x + y^2 - 10y + k = 0$ is 25π . The value of k is equal to
(a) -1 (b) 1 (c) 0 (d) 2
(e) 3
82. $\int_{2016}^{2017} \frac{\sqrt{x}}{\sqrt{x} + \sqrt{4033-x}} dx$ is equal to
(a) $\frac{1}{4}$ (b) $\frac{3}{2}$ (c) $\frac{2017}{2}$ (d) $\frac{1}{2}$
(e) 508
83. The solution of $\frac{dy}{dx} + y \tan x = \sec x$, $y(0) = 0$ is
(a) $y \sec x = \tan x$ (b) $y \tan x = \sec x$
(c) $\tan x = y \tan x$ (d) $x \sec x = \tan y$
(e) $y \cot x = \sec x$
84. If the vectors $2\hat{i} + 2\hat{j} + 6\hat{k}$, $2\hat{i} + \lambda\hat{j} + 6\hat{k}$, $2\hat{i} - 3\hat{j} + \hat{k}$ are coplanar, then the value of λ is
(a) -10 (b) 1 (c) 0 (d) 10
(e) 2
85. The distance between $(2, 1, 0)$ and $2x + y + 2z + 5 = 0$ is
(a) 10 (b) $\frac{10}{3}$ (c) $\frac{10}{9}$ (d) 5
(e) 1
86. The equation of the hyperbola with vertices $(0, \pm 15)$ and foci $(0, \pm 20)$ is
(a) $\frac{x^2}{175} - \frac{y^2}{225} = 1$ (b) $\frac{x^2}{625} - \frac{y^2}{125} = 1$
(c) $\frac{y^2}{225} - \frac{x^2}{125} = 1$ (d) $\frac{y^2}{65} - \frac{x^2}{65} = 1$
(e) $\frac{y^2}{225} - \frac{x^2}{175} = 1$
87. The value of $\frac{15^3 + 6^3 + 3 \cdot 6 \cdot 15 \cdot 21}{1 + 4(6) + 6(36) + 4(216) + 1296}$ is equal to
(a) $\frac{29}{7}$ (b) $\frac{7}{19}$ (c) $\frac{6}{17}$ (d) $\frac{21}{19}$
(e) $\frac{27}{7}$
88. The equation of the plane that passes through the points $(1, 0, 2)$, $(-1, 1, 2)$, $(5, 0, 3)$ is
(a) $x + 2y - 4z + 7 = 0$ (b) $x + 2y - 3z + 7 = 0$
(c) $x - 2y + 4z + 7 = 0$ (d) $2y - 4z - 7 + x = 0$
(e) $x + 2y + 3z + 7 = 0$
89. The vertex of the parabola $y^2 - 4y - x + 3 = 0$ is
(a) $(-1, 3)$ (b) $(-1, 2)$
(c) $(2, -1)$ (d) $(3, -1)$
(e) $(1, 2)$
90. If $\vec{a}, \vec{b}, \vec{c}$ are vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ and $|\vec{a}| = 7, |\vec{b}| = 5, |\vec{c}| = 3$, then the angle between \vec{c} and \vec{b} is
(a) $\frac{\pi}{3}$ (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{4}$ (d) π
(e) 0
91. Let $f(x) = 2x^3 - 9ax^2 + 12a^2x + 1$, where $a > 0$. The minimum of f is attained at a point q and the maximum is attained at a point p . If $p^3 = q$, then a is equal to
(a) 1 (b) 3 (c) 2 (d) $\sqrt{2}$
(e) $\frac{1}{2}$
92. For all real numbers x and y , it is known that the real valued function f satisfies $f(x) + f(y) = f(x + y)$. If $f(1) = 7$, then $\sum_{r=1}^{100} f(r)$ is equal to
(a) $7 \times 51 \times 102$ (b) $6 \times 50 \times 102$
(c) $7 \times 50 \times 102$ (d) $6 \times 25 \times 102$
(e) $7 \times 50 \times 101$
93. The eccentricity of the ellipse $\frac{(x-1)^2}{2} + \left(y + \frac{3}{4}\right)^2 = \frac{1}{16}$ is
(a) $\frac{1}{\sqrt{2}}$ (b) $\frac{1}{2\sqrt{2}}$ (c) $\frac{1}{2}$ (d) $\frac{1}{4}$
(e) $\frac{1}{4\sqrt{2}}$
94. $\int_{-1}^1 \max\{x, x^3\} dx$ is equal to
(a) $\frac{3}{4}$ (b) $\frac{1}{4}$ (c) $\frac{1}{2}$ (d) 1
(e) 0
95. If $x \in \left[0, \frac{\pi}{2}\right], y \in \left[0, \frac{\pi}{2}\right]$ and $\sin x + \cos y = 2$, then the value of $x + y$ is equal to

- (a) 2π (b) π (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{2}$
(e) 0

96. Let a , $a + r$ and $a + 2r$ be positive real numbers such that their product is 64. Then the minimum value of $a + 2r$ is equal to

- (a) 4 (b) 3 (c) 2 (d) $\frac{1}{2}$
(e) 1

97. The sum $S = \frac{1}{9!} + \frac{1}{3!7!} + \frac{1}{5!5!} + \frac{1}{7!3!} + \frac{1}{9!}$ is equal to

- (a) $\frac{2^{10}}{8!}$ (b) $\frac{2^9}{10!}$ (c) $\frac{2^7}{10!}$ (d) $\frac{2^6}{10!}$
(e) $\frac{2^5}{8!}$

98. If $f(x) = \begin{vmatrix} x & x^2 & x^3 \\ 1 & 2x & 3x^2 \\ 0 & 2 & 6x \end{vmatrix}$, then $f'(x)$ is equal to

- (a) $x^3 + 6x^2$ (b) $6x^3$
(c) $3x$ (d) $6x^2$
(e) 0

99. $\int \frac{x^2}{1+(x^3)^2} dx$ is equal to

- (a) $\tan^{-1}(x^2) + c$ (b) $\frac{2}{3}\tan^{-1}(x^3) + c$
(c) $\frac{1}{3}\tan^{-1}(x^3) + c$ (d) $\frac{1}{2}\tan^{-1}(x^2) + c$
(e) $\tan^{-1}(x^3) + c$

100. Let $f_n(x)$ be the n^{th} derivative of $f(x)$. The least value of n so that $f_n = f_{n+1}$, where $f(x) = x^2 + e^x$ is

- (a) 4 (b) 5 (c) 2 (d) 3
(e) 6

101. $\sin 765^\circ$ is equal to

- (a) 1 (b) 0 (c) $\frac{\sqrt{3}}{2}$ (d) $\frac{1}{2}$
(e) $\frac{1}{\sqrt{2}}$

102. The distance of the point $(3, -5)$ from the line $3x - 4y - 26 = 0$ is

- (a) $\frac{3}{7}$ (b) $\frac{2}{5}$ (c) $\frac{7}{5}$ (d) $\frac{3}{5}$
(e) 1

103. The difference between the maximum and minimum value of the function

$$f(x) = \int_0^x (t^2 + t + 1)dt \text{ on } [2, 3] \text{ is}$$

- (a) $\frac{39}{6}$ (b) $\frac{49}{6}$ (c) $\frac{59}{6}$ (d) $\frac{69}{6}$
(e) $\frac{79}{6}$

104. If a and b are the non zero distinct roots of $x^2 + ax + b = 0$, then the minimum value of $x^2 + ax + b$ is

- (a) $\frac{2}{3}$ (b) $\frac{9}{4}$ (c) $\frac{-9}{4}$ (d) $\frac{-2}{3}$
(e) 1

105. If the straight line $y = 4x + c$ touches the ellipse

$$\frac{x^2}{4} + y^2 = 1 \text{ then } c \text{ is equal to}$$

- (a) 0 (b) $\pm\sqrt{65}$ (c) $\pm\sqrt{62}$ (d) $\pm\sqrt{2}$
(e) ± 13

106. The equations $\lambda x - y = 2$, $2x - 3y = -\lambda$ and $3x - 2y = -1$ are consistent for

- (a) $\lambda = -4$ (b) $\lambda = 1, 4$
(c) $\lambda = 1, -4$ (d) $\lambda = -1, 4$
(e) $\lambda = -1$

107. The set $\{(x, y): |x| + |y| = 1\}$ in the xy plane represents

- (a) a square (b) a circle
(c) an ellipse
(d) a rectangle which is not a square
(e) a rhombus which is not a square

108. The value of $\cos \left(\tan^{-1} \left(\frac{3}{4} \right) \right)$ is

- (a) $\frac{4}{5}$ (b) $\frac{3}{5}$ (c) $\frac{3}{4}$ (d) $\frac{2}{5}$
(e) 0

109. Let $A(6, -1)$, $B(1, 3)$ and $C(x, 8)$ be three points such that $AB = BC$. The values of x are

- (a) 3, 5 (b) -3, 5 (c) 3, -5 (d) 4, 5
(e) -3, -5

110. In an experiment with 15 observations on x , the following results were available $\sum x^2 = 2830$ and $\sum x = 170$. One observation that was 20, was found to be wrong and was replaced by the correct value 30. Then the corrected variance is

- (a) 9.3 (b) 8.3 (c) 188.6 (d) 177.3
(e) 78

111. The angle between the pair of lines

$$\frac{x-2}{2} = \frac{y-1}{5} = \frac{z+3}{-3} \text{ and } \frac{x+2}{-1} = \frac{y-4}{8} = \frac{z-5}{4} \text{ is}$$

- (a) $\cos^{-1}\left(\frac{21}{9\sqrt{38}}\right)$ (b) $\cos^{-1}\left(\frac{23}{9\sqrt{38}}\right)$
 (c) $\cos^{-1}\left(\frac{24}{9\sqrt{38}}\right)$ (d) $\cos^{-1}\left(\frac{25}{9\sqrt{38}}\right)$
 (e) $\cos^{-1}\left(\frac{26}{9\sqrt{38}}\right)$

112. Let \vec{a} be a unit vector. If $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 12$, then the magnitude of \vec{x} is

- (a) $\sqrt{8}$ (b) $\sqrt{9}$ (c) $\sqrt{10}$ (d) $\sqrt{13}$
 (e) $\sqrt{12}$

113. The area of the triangular region whose sides are $y = 2x + 1$, $y = 3x + 1$ and $x = 4$ is

- (a) 5 (b) 6 (c) 7 (d) 8
 (e) 9

114. If ${}^nC_{r-1} = 36$, ${}^nC_r = 84$ and ${}^nC_{r+1} = 126$, then the value of r is

- (a) 9 (b) 3 (c) 4 (d) 5
 (e) 6

115. Let $f(x+y) = f(x)f(y)$ and $f(x) = 1 + \sin(3x)g(x)$, where g is differentiable. Then $f'(x)$ is equal to

- (a) $3f(x)$ (b) $g(0)$
 (c) $f(x)g(0)$ (d) $3g(x)$
 (e) $3f(x)g(0)$

116. The roots of the equation $\begin{vmatrix} x-1 & 1 & 1 \\ 1 & x-1 & 1 \\ 1 & 1 & x-1 \end{vmatrix} = 0$ are

- (a) 1, 2 (b) -1, 2 (c) -1, -2 (d) 1, -2
 (e) 1, 1

117. If the 7th and 8th term of the binomial expansion

$$(2a-3b)^n \text{ are equal, then } \frac{2a+3b}{2a-3b} \text{ is equal to}$$

- (a) $\frac{13-n}{n+1}$ (b) $\frac{n+1}{13-n}$
 (c) $\frac{6-n}{13-n}$ (d) $\frac{n-1}{13-n}$
 (e) $\frac{2n-1}{13-n}$

118. Standard deviation of first n odd natural numbers is

- (a) \sqrt{n} (b) $\sqrt{\frac{(n+2)(n+1)}{3}}$
 (c) $\sqrt{\frac{n^2-1}{3}}$ (d) n
 (e) $2n$

119. Let $S = \{1, 2, 3, \dots, 10\}$. The number of subsets of S containing only odd numbers is

- (a) 15 (b) 31 (c) 63 (d) 7
 (e) 5

120. The area of the parallelogram with vertices $(0, 0)$, $(7, 2)$, $(5, 9)$ and $(12, 11)$ is

- (a) 50 (b) 54 (c) 51 (d) 52
 (e) 53

SOLUTIONS

1. (a) : We have,

$$\begin{vmatrix} 1 & 1 & 1 \\ p & q & r \\ p & q & r+1 \end{vmatrix} = \begin{vmatrix} 1 & 1 & 1 \\ p & q & r \end{vmatrix} + \begin{vmatrix} 1 & 1 & 0 \\ p & q & 0 \\ p & q & 1 \end{vmatrix}$$

$$= 0 + \begin{vmatrix} 1 & 1 & 0 \\ p & q & 0 \\ p & q & 1 \end{vmatrix} \quad (\because R_2 \sim R_3 \text{ in 1st determinant})$$

$$= \begin{vmatrix} 1 & 1 & 0 \\ p & q & 0 \\ p & q & 1 \end{vmatrix} = 1(q-p)$$

2. (b) : Given, $A = \begin{bmatrix} 5 & 0 \\ 1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$

$$\text{Now, } 4A + 5B - C = O$$

$$\Rightarrow C = 4A + 5B = 4 \begin{bmatrix} 5 & 0 \\ 1 & 0 \end{bmatrix} + 5 \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 20 & 0 \\ 4 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 5 \\ -5 & 0 \end{bmatrix} = \begin{bmatrix} 20 & 5 \\ -1 & 0 \end{bmatrix}$$

3. (a) : Given, $U = \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$

$$\text{Since, } U^{-1} = \frac{1}{|U|} \text{adj}(U)$$

$$\text{Hence, } \text{adj}(U) = \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}^T = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

$$|U| = \left(\frac{1}{\sqrt{2}} \cdot \frac{1}{\sqrt{2}} \right) - \left(\frac{-1}{\sqrt{2}} \cdot \frac{1}{\sqrt{2}} \right) = \frac{1}{2} + \frac{1}{2} = 1$$

$$\therefore U^{-1} = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} = U^T$$

4. (a) : We have, $A = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}$

$$\text{Hence, } \text{adj}(A) = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}^T = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{Now, } |A| = 0 + (1)(-1) = -1$$

$$\therefore A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix} = A^T$$

5. (a) : We have, $\begin{pmatrix} x+y & x-y \\ 2x+z & x+z \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}$

On comparing, we get

$$x+y=0 \quad \dots(\text{i}), \quad x-y=0 \quad \dots(\text{ii})$$

$$2x+z=1 \quad \dots(\text{iii}), \quad x+z=1 \quad \dots(\text{iv})$$

On solving (iii) and (iv), we get $x=0, z=1$.

Hence, from (i), we get $y=0$

$$\therefore x=0, y=0, z=1.$$

6. (b) : We have, $\begin{pmatrix} 7 & 1 & 5 \\ 8 & 0 & 0 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} + 5 \begin{pmatrix} 1 \\ 0 \end{pmatrix}$

$$= \begin{pmatrix} 14+3+5 \\ 16+0+0 \end{pmatrix} + \begin{pmatrix} 5 \\ 0 \end{pmatrix} = \begin{pmatrix} 22 \\ 16 \end{pmatrix} + \begin{pmatrix} 5 \\ 0 \end{pmatrix} = \begin{pmatrix} 27 \\ 16 \end{pmatrix}$$

7. (d) : Let $A = \begin{pmatrix} 1 & 2 & 4 \\ 1 & 3 & 5 \\ 1 & 4 & a \end{pmatrix}$

Since A is singular matrix $\therefore |A| = 0$

$$\text{Hence, } \begin{vmatrix} 1 & 2 & 4 \\ 1 & 3 & 5 \\ 1 & 4 & a \end{vmatrix} = 0$$

$$\Rightarrow 1(3a-20) - 2(a-5) + 4(4-3) = 0 \Rightarrow a = 6$$

8. (d) : We have, $\begin{pmatrix} 1 & 2 & -3 \\ 0 & 4 & 5 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$

$$\Rightarrow \begin{pmatrix} x+2y-3z \\ 4y+5z \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

On comparing, we get

$$x+2y-3z=1 \quad \dots(\text{i}) \quad 4y+5z=1 \quad \dots(\text{ii})$$

$$\text{and } z=1 \quad \dots(\text{iii})$$

On solving (i), (ii) and (iii), we get $z=1, y=-1$ and $x=6$

9. (b) : We have, $A = \begin{pmatrix} 1 & 5 \\ 0 & 2 \end{pmatrix}$

$$A^2 = A \cdot A = \begin{pmatrix} 1 & 5 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & 5 \\ 0 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 15 \\ 0 & 4 \end{pmatrix}$$

$$3A = 3 \begin{pmatrix} 1 & 5 \\ 0 & 2 \end{pmatrix} = \begin{pmatrix} 3 & 15 \\ 0 & 6 \end{pmatrix}$$

$$2I = 2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$$

$$\therefore A^2 - 3A + 2I = \begin{pmatrix} 1 & 15 \\ 0 & 4 \end{pmatrix} - \begin{pmatrix} 3 & 15 \\ 0 & 6 \end{pmatrix} + \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$$

$$= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = O$$

10. (a) : We have, $\begin{pmatrix} 2x+y & x+y \\ p-q & p+q \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$

On comparing, we get

$$2x+y=1 \quad \dots(\text{i}), \quad x+y=1 \quad \dots(\text{ii})$$

$$p-q=0 \quad \dots(\text{iii}), \quad p+q=0 \quad \dots(\text{iv})$$

On solving (i) and (ii), we get $x=0$ and $y=1$

And, on solving (iii) and (iv), we get $p=0$ and $q=0$

$$\therefore (x, y, p, q) = (0, 1, 0, 0)$$

11. (b) : Let $|\sqrt{4+2\sqrt{3}}| - |\sqrt{4-2\sqrt{3}}| = x$

On squaring both sides, we get

$$(\sqrt{4+2\sqrt{3}} - \sqrt{4-2\sqrt{3}})^2 = x^2$$

$$\Rightarrow (4+2\sqrt{3}) + (4-2\sqrt{3}) - 2(\sqrt{4+2\sqrt{3}})(\sqrt{4-2\sqrt{3}}) = x^2$$

$$\Rightarrow 8 - 2\sqrt{16-12} = x^2 \Rightarrow 8 - 4 = x^2$$

$$\Rightarrow x^2 = 4 \Rightarrow x = \pm 2$$

$$\text{Hence, } |\sqrt{4+2\sqrt{3}}| - |\sqrt{4-2\sqrt{3}}| = \pm 2$$

12. (a) : We have, $8^{2/3} - 16^{1/4} - 9^{1/2}$
 $= ((2^3)^{2/3}) - ((2^4)^{1/4}) - ((3^2)^{1/2})$
 $= 4 - 2 - 3 = 4 - 5 = -1$

13. (a) : Given, $x = 2$ is a root of $y = 4x^2 - 14x + q = 0$
then, $x = 2$ satisfy the given equation i.e., $y(2) = 0$
 $\Rightarrow 4(2)^2 - 14(2) + q = 0$
 $\Rightarrow 16 - 28 + q = 0 \Rightarrow q = 12$
 $\therefore y = 4x^2 - 14x + 12 = 0$
 $\Rightarrow y = 4x^2 - 6x - 8x + 12$
 $\Rightarrow y = 2x(2x - 3) - 4(2x - 3)$
 $\Rightarrow y = (2x - 4)(2x - 3) \Rightarrow y = (x - 2)(4x - 6)$

14. (b) : We have, $3x^2 - 2x - 6 = 0$
 \therefore Roots of given equation (x_1, x_2)

$$= \frac{-(-2) \pm \sqrt{4 + 72}}{6} = \frac{2 \pm \sqrt{76}}{6} = \frac{2 \pm 2\sqrt{19}}{6}$$

$$= \frac{1 \pm \sqrt{19}}{3}$$

Hence, $x_1 = \frac{1 + \sqrt{19}}{3}$ and $x_2 = \frac{1 - \sqrt{19}}{3}$

Now, $x_1^2 + x_2^2 = \left(\frac{1 + \sqrt{19}}{3}\right)^2 + \left(\frac{1 - \sqrt{19}}{3}\right)^2$
 $= \frac{1}{9}(1 + 19 + 2\sqrt{19} + 1 + 19 - 2\sqrt{19}) = \frac{40}{9}$

15. (c) : Given, x_1 and x_2 are roots of $x^2 + px - 3 = 0$.
Also, $x_1^2 + x_2^2 = 10$

\therefore Sum of roots $= (x_1 + x_2) = \frac{-p}{1} = -p$

And, product of roots $= x_1 x_2 = \frac{-3}{1} = -3$

$\therefore (x_1 + x_2)^2 = p^2$

$\Rightarrow x_1^2 + x_2^2 + 2x_1 x_2 = p^2 \Rightarrow 10 + 2(-3) = p^2$

$\Rightarrow p^2 = 4 \Rightarrow p = \pm 2$

Hence, the value of p is 2 or -2.

16. (a) : Given, $mx^2 + 6x + (2m - 1) = 0$ and product of roots is -1.

$\Rightarrow \frac{(2m-1)}{m} = -1 \Rightarrow 2m - 1 = -m \Rightarrow 3m = 1 \Rightarrow m = \frac{1}{3}$

17. (e) : We have,

$$f(x) = \frac{1}{x^2 + 4x + 4} - \frac{4}{x^4 + 4x^3 + 4x^2} + \frac{4}{x^3 + 2x^2}$$

$$= \frac{1}{(x+2)^2} - \frac{4}{x^2(x+2)^2} + \frac{4}{x^2(x+2)}$$

$$\therefore f(1/2) = \frac{1}{\left(\frac{1}{2} + 2\right)^2} - \frac{4}{\left(\frac{1}{2}\right)^2 \left(\frac{1}{2} + 2\right)^2} + \frac{4}{\left(\frac{1}{2}\right)^2 \left(\frac{1}{2} + 2\right)}$$

$$= \frac{1}{\left(\frac{5}{2}\right)^2} - \frac{4}{\frac{1}{4} \left(\frac{5}{2}\right)^2} + \frac{4}{\frac{1}{4} \left(\frac{5}{2}\right)}$$

$$= \frac{4}{25} - \frac{64}{25} + \frac{32}{5} = \frac{-60 + 160}{25} = \frac{100}{25} = 4$$

18. (b) : We have, $x^2 + bx + 1 = 0$ and x, y are its roots.
 \therefore Sum of roots $= (x + y) = -b$... (i)
And product of roots $= (xy) = 1$... (ii)

Now, $\frac{1}{x+b} + \frac{1}{y+b} = \frac{y+b+x+b}{(x+b)(y+b)}$
 $= \frac{(x+y)+2b}{xy+b(x+y)+b^2} = \frac{(-b)+2b}{1-b^2+b^2}$
 $= b$ (Using (i) and (ii))

19. (b) : Let y be the common root of
 $x^5 + ax + 1 = 0$ and $x^6 + ax^2 + 1 = 0$
Then, $y^5 + ay + 1 = 0$ and $y^6 + ay^2 + 1 = 0$

$\Rightarrow y^5 + ay + 1 = y^6 + ay^2 + 1$

$\Rightarrow y^5 - y^6 + ay - ay^2 = 0$

$\Rightarrow y^5(1 - y) + ay(1 - y) = 0$

$\Rightarrow (y^5 + ay)(1 - y) = 0 \Rightarrow y = 1$

Hence, the common root is 1.

i.e., $1 + a + 1 = 0 \Rightarrow a = -2$

20. (a) : Let x_1 and x_2 be the roots of $ax^2 + x + 1 = 0$ then,

$x_1 + x_2 = \frac{-1}{a}$... (i) and $x_1 x_2 = \frac{1}{a}$... (ii)

Also, $x_1 : x_2 = 1 : 1 \Rightarrow x_1 = x_2$... (iii)

Using (iii) in (i), we get $2x_1 = \frac{-1}{a} \Rightarrow x_1 = \frac{-1}{2a}$

Using (iii) in (ii), we get $x_1^2 = \frac{1}{a} \therefore \frac{1}{4a^2} = \frac{1}{a}$

$\Rightarrow 4a = 1 \Rightarrow a = \frac{1}{4}$

21. (c) : We have, $z^2 + z + 1 = 0$

$\Rightarrow z = \omega$ or ω^2

where ω, ω^2 are complex cube roots of unity.

Now, $\left(z + \frac{1}{z}\right)^2 + \left(z^2 + \frac{1}{z^2}\right)^2 + \left(z^3 + \frac{1}{z^3}\right)^2$
 $= \left(\omega + \frac{1}{\omega}\right)^2 + \left(\omega^2 + \frac{1}{\omega^2}\right)^2 + \left(\omega^3 + \frac{1}{\omega^3}\right)^2$

$$= \left(\frac{\omega^2 + 1}{\omega} \right)^2 + \left(\frac{\omega^4 + 1}{\omega^2} \right)^2 + (1 + 1)^2$$

[using $\omega^3 = 1$]

$$= \left(\frac{-\omega}{\omega} \right)^2 + \left(\frac{-\omega^2}{\omega^2} \right)^2 + 4 \quad [\text{using } 1 + \omega + \omega^2 = 0]$$

$$= 1 + 1 + 4 = 6$$

22. (b) : We have, $\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & -1-w^2 & w^2 \\ 1 & w & w^4 \end{vmatrix}$

$$\Rightarrow \Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & -1-w^2 & w^2 \\ 1 & w & w \end{vmatrix} = \begin{vmatrix} 1 & 1 & 1 \\ 1 & w & w^2 \\ 1 & w & w \end{vmatrix}$$

($\because 1 + w + w^2 = 0$)

Applying $C_1 \rightarrow C_1 - C_2$ and $C_2 \rightarrow C_2 - C_3$, we get

$$\Delta = \begin{vmatrix} 0 & 0 & 1 \\ 1-w & w-w^2 & w^2 \\ 1-w & 0 & w \end{vmatrix}$$

$$= 1(0 - (1-w)(w-w^2)) \quad (\text{on expanding along } R_1)$$

$$= -(w - w^2 - w^2 + w^3)$$

$$= -(-1 - w^2 - w^2 - w^2 + 1) = -(-3w^2) = 3w^2$$

23. (d) : We have, $\begin{vmatrix} 3i & -9i & 1 \\ 2 & 9i & -1 \\ 10 & 9 & i \end{vmatrix} = (x + iy)$

$$\Rightarrow 3i(9i^2 + 9) + 9i(2i + 10) + (18 - 90i) = x + iy$$

$$\Rightarrow 3i(-9 + 9) + 18i^2 + 90i + 18 - 90i = x + iy$$

$$\Rightarrow -18 + 18 = x + iy$$

$$\Rightarrow 0 + 0i = x + iy$$

On comparing, we get $x = 0$ and $y = 0$

24. (a) : We have, $z = \cos\left(\frac{\pi}{3}\right) - i \sin\left(\frac{\pi}{3}\right)$

$$= \frac{1}{2} - \frac{\sqrt{3}i}{2} = \frac{1 - \sqrt{3}i}{2}$$

then, $z^2 - z + 1 = \left(\frac{1 - \sqrt{3}i}{2}\right)^2 - \left(\frac{1 - \sqrt{3}i}{2}\right) + 1$

$$= \frac{1}{4}(1 - 3 - 2\sqrt{3}i) + \frac{1}{2} + \frac{\sqrt{3}i}{2}$$

$$= \frac{-1}{2} - \frac{\sqrt{3}i}{2} + \frac{1}{2} + \frac{\sqrt{3}i}{2} = 0$$

25. (c) : Let $z = \cos\left(\frac{\pi}{12}\right) + i \sin\left(\frac{\pi}{12}\right)$

$$\therefore \frac{1}{z} = \left(\cos\frac{\pi}{12} - i \sin\frac{\pi}{12} \right)$$

Now, from given expression, we have

$$\left(\frac{1 + \cos\left(\frac{\pi}{12}\right) + i \sin\left(\frac{\pi}{12}\right)}{1 + \cos\left(\frac{\pi}{12}\right) - i \sin\left(\frac{\pi}{12}\right)} \right)^{72} = \left(\frac{1+z}{1+z^{-1}} \right)^{72}$$

$$= \left(\frac{(1+z)z}{(z+1)} \right)^{72} = (z)^{72} = \left(\cos\left(\frac{\pi}{12}\right) + i \sin\left(\frac{\pi}{12}\right) \right)^{72}$$

$$= \cos\left(\frac{72\pi}{12}\right) + i \sin\left(\frac{72\pi}{12}\right)$$

(Using De-Moivre's theorem)

$$= \cos 6\pi + i \sin 6\pi = 1$$

26. (e) : We have, $A = \begin{vmatrix} 4 & k & k \\ 0 & k & k \\ 0 & 0 & k \end{vmatrix}$

Now, expanding along C_1 , we get $|A| = 4(k^2)$

But $\det(A) = 256$

(Given)

\therefore On comparing, we get

$$4k^2 = 256 \Rightarrow k^2 = 64 \Rightarrow k = \pm 8$$

Hence, $|k| = 8$

27. (c) : We have, $A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$

$$\therefore A^2 = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$$

$$A^3 = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix} \therefore A^n = \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}$$

Hence, $A^n + nI = \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix} + n \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

$$= \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix} + \begin{pmatrix} n & 0 \\ 0 & n \end{pmatrix} = \begin{pmatrix} 1+n & 0 \\ n & 1+n \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} n & 0 \\ n & n \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + n \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$$

$$= I + nA$$

28. (a) : Let $z = x + iy$

$$\therefore |z| = 5 \Rightarrow \sqrt{x^2 + y^2} = 5 \Rightarrow x^2 + y^2 = 25$$

$$\text{Now, } w = \frac{z-5}{z+5} = \frac{x+iy-5}{x+iy+5} = \frac{(x-5)+iy}{(x+5)+iy}$$

On rationalizing the denominator, we get

$$\begin{aligned} \frac{((x-5)+iy)((x+5)-iy)}{(x+5)^2+(y)^2} &= \frac{x^2-25+y^2+10yi}{(x+5)^2+y^2} \\ &= \frac{10yi}{(x+5)^2+y^2} \quad [\text{Using } x^2+y^2=25] \\ &= 0 + \frac{10yi}{(x+5)^2+y^2} \quad \therefore R(w) = 0 \end{aligned}$$

29. (b) : We have, $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

$$\begin{aligned} \therefore A^2 &= A \cdot A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix} \\ &= 2 \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = 2A \end{aligned}$$

$$\begin{aligned} A^3 &= A^2 \cdot A = \begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 4 & 4 \\ 4 & 4 \end{pmatrix} \\ &= 4 \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = 2^2 \cdot A \end{aligned}$$

Similarly $A^n = 2^{n-1} \cdot A$

$$\therefore A^{2017} = 2^{2017-1} \cdot A = 2^{2016} \cdot A$$

30. (c) : We have, $a = e^{i\theta} = \cos \theta + i \sin \theta$ (polar form)

$$\begin{aligned} \therefore \frac{1+a}{1-a} &= \frac{1+\cos \theta + i \sin \theta}{1-(\cos \theta + i \sin \theta)} \\ &= \frac{2 \cos^2 \frac{\theta}{2} + i 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{2 \sin^2 \frac{\theta}{2} - i 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}} \\ &= \frac{2 \cos \frac{\theta}{2} \left[\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} \right]}{2 \sin \frac{\theta}{2} \left[\sin \frac{\theta}{2} - i \cos \frac{\theta}{2} \right]} \\ &= \cot \frac{\theta}{2} \left[\frac{\left(\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} \right) \times \left(\sin \frac{\theta}{2} + i \cos \frac{\theta}{2} \right)}{\left(\sin \frac{\theta}{2} - i \cos \frac{\theta}{2} \right) \left(\sin \frac{\theta}{2} + i \cos \frac{\theta}{2} \right)} \right] \end{aligned}$$

[Rationalizing the denominator]

$$\begin{aligned} &= \cot \frac{\theta}{2} \left[\frac{\cos \frac{\theta}{2} \sin \frac{\theta}{2} + i \cos^2 \frac{\theta}{2} + i \sin^2 \frac{\theta}{2} - \cos \frac{\theta}{2} \sin \frac{\theta}{2}}{\sin^2 \frac{\theta}{2} + \cos^2 \frac{\theta}{2}} \right] \\ &= i \cot \frac{\theta}{2} \end{aligned}$$

31. (c) : We have, $x + y + z = -3$... (i)

On squaring both sides, we get

$$(x+y+z)^2 = (-3)^2$$

$$\Rightarrow x^2 + y^2 + z^2 + 2xy + 2yz + 2zx = 9 \quad \dots (ii)$$

Also, x, y, z are in A.P. $\Rightarrow 2y = x + z$

$$\Rightarrow 2y = -3 - y \Rightarrow y = -1 \quad \dots (iii)$$

$$\text{So, } xyz = 8 \Rightarrow xz = -8 \quad \dots (iv)$$

Now, using (iii) and (iv) in (ii), we get

$$x^2 + y^2 + z^2 - 2x - 2z - 16 = 9$$

$$\Rightarrow x^2 + y^2 + z^2 - 2(x+z) - 16 = 9$$

$$\Rightarrow x^2 + y^2 + z^2 + 4 - 16 = 9$$

$$\Rightarrow x^2 + y^2 + z^2 = 9 + 12 = 21$$

Hence, $x^2 + y^2 + z^2 = 21$

32. (c) : We have the series of an A.P. as 10, 7, 4, ...

\therefore First term (a) = 10

Common difference (d) = $7 - 10 = -3$

$$\therefore a_{30} = a + (30-1)d = 10 + 29(-3) = 10 - 87 = -77$$

33. (e) : Given, arithmetic mean of x and y is 3

$$\text{i.e., } \frac{x+y}{2} = 3 \Rightarrow x+y = 6 \quad \dots (i)$$

and geometric mean of x and y is 1

$$\text{i.e., } \sqrt{xy} = 1 \Rightarrow xy = 1 \quad \dots (ii)$$

Squaring (i) on both sides, we get

$$(x+y)^2 = (6)^2 \Rightarrow x^2 + y^2 + 2xy = 36$$

$$\Rightarrow x^2 + y^2 + 2 = 36 \quad (\text{Using (ii)})$$

$$\Rightarrow x^2 + y^2 = 34$$

34. (d) : We have, $3^{2x-1} = 81^{1-x}$

$$\Rightarrow (3)^{2x-1} = ((3)^4)^{1-x} \Rightarrow (3)^{2x-1} = (3)^{4-4x}$$

\therefore On comparing, we get, $2x-1 = 4-4x$

$$\Rightarrow 6x = 5 \quad \text{or} \quad x = \frac{5}{6}$$

35. (c) : Given series $3, 1, \frac{1}{3}, \dots$ forms a G.P.

where first term (a) = 3 and common ratio (r) = $\frac{1}{3}$

$$\therefore \text{Sixth term, } a_6 = ar^5$$

$$= (3) \left(\frac{1}{3} \right)^5 = \left(\frac{1}{3} \right)^4 = \frac{1}{81}$$

36. (a) : Let three numbers in A.P. be a, b and c

Then, according to question

$$a+b+c = 21 \quad \dots (i) \quad \text{and} \quad ac = 45 \quad \dots (ii)$$

$$\therefore a, b \text{ and } c \text{ are in A.P.} \quad \therefore 2b = c + a \quad \dots (iii)$$

Substituting (iii) in (i), we get

$$3b = 21 \Rightarrow b = 7$$

Hence, product of these three numbers = abc

$$= 7(45) = 315 \quad (\text{Using (ii)})$$

37. (b) : Given, $a + 1, 2a + 1, 4a - 1$ are in A.P.

$$\therefore 2(2a + 1) = 4a - 1 + a + 1$$

$$\Rightarrow 4a + 2 = 5a \Rightarrow a = 2$$

38. (b) : Given, arithmetic mean of x and y is 9

$$\text{i.e., } \frac{x+y}{2} = 9 \Rightarrow x + y = 18$$

Geometric mean of x and y is 4 i.e., $\sqrt{xy} = 4 \Rightarrow xy = 16$

Now, sum of roots $(x + y) = 18$

Product of roots $(xy) = 16$

\therefore Required quadratic equation is $x^2 - 18x + 16 = 0$

39. (c) : Total number of outcomes = 8 i.e.,

{HHH, HHT, HTH, THH, TTH, THT, HTT, TTT}

Number of favourable outcomes = 4

i.e., {TTH, THT, HTT, TTT} = 4

$$\therefore P(\text{getting at least 2 tails}) = \frac{4}{8} = \frac{1}{2}$$

40. (e) : Number of letters in TRICKS = 6

Number of favourable outcomes = {T, R} = 2

$$\therefore P(\text{either T or R}) = \frac{2}{6} = \frac{1}{3}$$

41. (c) : Total number of outcomes = ${}^{10}C_4$

2 red balls can be selected from 4 red balls in 4C_2 ways

And, remaining 2 balls can be selected from 2 white balls and 4 black balls in 6C_2 ways.

$$\therefore \text{Required probability} = \frac{{}^4C_2 \times {}^6C_2}{{}^{10}C_4} = \frac{3}{7} \text{ or } \frac{9}{21}$$

42. (d) : Let A and B be two events of knowing lesson I and lesson II respectively.

\therefore According to question,

$$P(A) = \frac{60}{100}; \quad P(B) = \frac{40}{100}$$

$$P(A \cap B) = \frac{20}{100}$$

$$\therefore P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{60}{100} + \frac{40}{100} - \frac{20}{100} = \frac{80}{100}$$

Hence, required probability = $P(A^C \cap B^C)$

$$= P(A \cup B)^C = 1 - P(A \cup B)$$

$$= 1 - \frac{80}{100} = \frac{20}{100} = \frac{1}{5}$$

43. (d) : Two distinct numbers can be chosen from 1, 2, 3, 4, 5 in 5C_2 ways.

Number of outcomes having arithmetic mean an integer i.e., $\{(1, 3), (1, 5), (2, 4), (3, 5)\} = 4$

$$\therefore \text{Required probability} = \frac{4}{{}^5C_2} = \frac{4}{10} = \frac{2}{5}$$

44. (e) : There are 9 elements in 3×3 matrices and each element can be filled in two ways either -1 or 1.

\therefore Total possible matrices = 2^9

45. (b) : Total possible set of 2×2 symmetric matrices of entries either zero or one = 8

Possible set of matrices having determinant not zero = 4

$$\therefore \text{Required probability} = \frac{4}{8} = \frac{1}{2}$$

46. (c) : Number of words that can be formed by using all 7 letters of word PROBLEM only once is 7!.

47. (b) : Number of diagonals in n -sided polygon

$$= \frac{n(n-3)}{2}$$

$$\therefore \text{Number of diagonals in hexagon} = \frac{6(6-3)}{2} = 9$$

48. (a) : We know sum of n odd numbers = n^2

Number of odd terms from 1 to 2001 = 1001

$$\therefore \text{Sum of 1001 odd terms} = (1001)^2$$

49. (e) : Total number of ways of selecting two balls = 4C_2

Number of ways of selecting 2 balls in which no ball is black = 2C_2

$$\therefore \text{Required probability} = \frac{{}^2C_2}{{}^4C_2} = \frac{1}{6}$$

50. (a) : We have, $z = i^9 + i^{19} = (i^2)^4 \cdot i + (i^2)^9 \cdot i$
 $= i + (-i) = 0 = 0 + 0i$

51. (a) : Given data is 6, 7, 10, 12, 13, 4, 8, 12

$$\therefore \text{Mean} = \frac{6+7+10+12+13+4+8+12}{8} = \frac{72}{8} = 9$$

52. (e) : Given, $x - 2 < 1 \Rightarrow x < 1 + 2 = 3$

Hence, set of all real numbers satisfying the inequality $x - 2 < 1$ is $(-\infty, 3)$

53. (b) : Given, $\frac{|x-3|}{x-3} > 0$

$$\Rightarrow |x-3| > 0 \Rightarrow x-3 > 0 \Rightarrow x > 3$$

$$\therefore x \in (3, \infty)$$

54. (c) : From the given data 8 has highest frequency.

\therefore Mode of the given data is 8.

55. (a) : Let the six numbers be $x_1, x_2, x_3, x_4, x_5, x_6$

Given, mean of six numbers = 41

$$\therefore \text{Mean} = \frac{x_1 + x_2 + x_3 + x_4 + x_5 + x_6}{6} = 41$$

$$\Rightarrow x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 246$$

56. (b) : Given, $\int_0^x f(t) dt = x^2 + e^x (x > 0)$

$\Rightarrow f(x) = 2x + e^x$

$\therefore f(1) = 2(1) + e = 2 + e$

57. (c) : $\int \frac{x+1}{x^{1/2}} dx = \int \sqrt{x} dx + \int \frac{1}{\sqrt{x}} dx$
 $= \frac{2}{3} x^{3/2} + 2x^{1/2} + c$

58. (e) : Let H and E be the two events of people speaking Hindi and English respectively.

$\therefore n(H) = 50, n(E) = 20$

$n(H \cap E) = 10$

$\therefore n(H \cup E) = n(H) + n(E) - n(H \cap E)$
 $= 50 + 20 - 10 = 60$

i.e., number of people who speak atleast one of two languages is 60.

59. (a) : Given, $f(x) = \frac{x+1}{x-1}$

$\therefore f(f(x)) = f\left(\frac{x+1}{x-1}\right) = \frac{\frac{x+1}{x-1} + 1}{\frac{x+1}{x-1} - 1}$
 $= \frac{(x+1+x-1)}{(x+1-x-1)} = \frac{2x}{2} = x$

60. (a) : Total number of outcomes = $6 \times 6 = 36$
 Favourable number of outcomes = 27

\therefore Required probability = $\frac{27}{36} = \frac{3}{4}$

61. (a) : We have, $\lim_{x \rightarrow 0} \frac{\sqrt{2+x} - \sqrt{2-x}}{x}$

On rationalizing the numerator, we get

$\lim_{x \rightarrow 0} \frac{(2+x) - (2-x)}{x(\sqrt{2+x} + \sqrt{2-x})} = \lim_{x \rightarrow 0} \frac{2x}{x(\sqrt{2+x} + \sqrt{2-x})}$
 $= \lim_{x \rightarrow 0} \frac{2}{\sqrt{2+x} + \sqrt{2-x}} = \frac{2}{2\sqrt{2}} = \frac{1}{\sqrt{2}}$

62. (b) : We have, $\int \frac{dx}{e^x + e^{-x} + 2} = \int \frac{e^x dx}{e^{2x} + 1 + 2e^x}$
 $= \int \frac{e^x dx}{(e^x + 1)^2} = \int \frac{dt}{t^2}$
 [Put $(e^x + 1) = t \Rightarrow e^x dx = dt]$
 $= \frac{-1}{t} + c = \frac{-1}{e^x + 1} + c$

63. (b) : We have, $\tan\left(\frac{\pi}{4} + \frac{\theta}{2}\right) + \tan\left(\frac{\pi}{4} - \frac{\theta}{2}\right)$

$= \frac{\tan \frac{\pi}{4} + \tan \frac{\theta}{2}}{1 - \tan \frac{\pi}{4} \tan \frac{\theta}{2}} + \frac{\tan \frac{\pi}{4} - \tan \frac{\theta}{2}}{1 + \tan \frac{\pi}{4} \tan \frac{\theta}{2}}$
 $= \frac{1 + \tan \frac{\theta}{2}}{1 - \tan \frac{\theta}{2}} + \frac{1 - \tan \frac{\theta}{2}}{1 + \tan \frac{\theta}{2}}$
 $= \frac{\left(1 + \tan \frac{\theta}{2}\right)^2 + \left(1 - \tan \frac{\theta}{2}\right)^2}{\left(1 - \tan^2 \frac{\theta}{2}\right)} = \frac{2\left(1 + \tan^2 \frac{\theta}{2}\right)}{1 - \tan^2 \frac{\theta}{2}}$
 $= 2\left(\frac{1}{\cos \theta}\right) = 2 \sec \theta$

64. (None of the options is correct) :

We have, $\int_{-1}^0 \frac{dx}{x^2 + x + 2}$
 $= \int_{-1}^0 \frac{dx}{-1x^2 + x + 2 + \frac{1}{4} - \frac{1}{4}} = \int_{-1}^0 \frac{dx}{\left(x + \frac{1}{2}\right)^2 + \frac{7}{4}}$
 $= \int_{-1}^0 \frac{dx}{-1\left(x + \frac{1}{2}\right)^2 + \left(\frac{\sqrt{7}}{2}\right)^2} = \frac{1}{\sqrt{7}/2} \tan^{-1} \frac{x+1/2}{\sqrt{7}/2} \Big|_{-1}^0$
 $= \frac{2}{\sqrt{7}} \left[\tan^{-1} \left(\frac{1/2}{\sqrt{7}/2} \right) - \tan^{-1} \left(\frac{-1/2}{\sqrt{7}/2} \right) \right]$
 $= \frac{2}{\sqrt{7}} \left(\tan^{-1} \frac{1}{\sqrt{7}} + \tan^{-1} \frac{1}{\sqrt{7}} \right) = \frac{4}{\sqrt{7}} \tan^{-1} \left(\frac{1}{\sqrt{7}} \right)$

65. (e) : Let $I = \int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$... (i)

$\Rightarrow I = \int_0^{\pi/2} \frac{\sqrt{\sin\left(\frac{\pi}{2} - x\right)} dx}{\sqrt{\sin\left(\frac{\pi}{2} - x\right)} + \sqrt{\cos\left(\frac{\pi}{2} - x\right)}}$
 $\Rightarrow I = \int_0^{\pi/2} \frac{\sqrt{\cos x} dx}{\sqrt{\cos x} + \sqrt{\sin x}}$... (ii)

Adding (i) and (ii), we get

$$2I = \int_0^{\pi/2} \left(\frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} \right) dx$$

$$\Rightarrow I = \frac{1}{2} \int_0^{\pi/2} 1 \cdot dx = \frac{1}{2} \left[\frac{\pi}{2} \right] = \frac{\pi}{4}$$

66. (b) : Given, (x, y) is equidistant from $(a+b, b-a)$ and $(a-b, a+b)$

\therefore Using distance formula, we have

$$\begin{aligned} \sqrt{(x-(a+b))^2 + (y-(b-a))^2} \\ = \sqrt{(x-(a-b))^2 + (y-(a+b))^2} \end{aligned}$$

On squaring both sides, we get


$$\begin{aligned} (x-(a+b))^2 + (y-(b-a))^2 \\ = (x-(a-b))^2 + (y-(a+b))^2 \\ \Rightarrow x^2 + a^2 + b^2 + 2ab - 2ax - 2bx + y^2 + b^2 + a^2 \\ - 2ab - 2by + 2ay \\ = x^2 + a^2 + b^2 - 2ab + 2bx - 2ax + y^2 + a^2 + b^2 \\ + 2ab - 2by - 2ay \\ \Rightarrow -4bx + 4ay = 0 \Rightarrow ay - bx = 0 \text{ or } bx - ay = 0 \end{aligned}$$

67. (e) : Given, $(1, 0)$, $(0, 1)$ and $(x, 8)$ are collinear
 \therefore Area of Δ formed by these points is zero

$$\text{i.e., } \frac{1}{2} |1(1-8) + 0(8-0) + x(0-1)| = 0$$

$$\Rightarrow \frac{1}{2} |-7-x| = 0 \Rightarrow x = -7$$

68. (a) : Graph of $\max\{x, x^2\}$ is shown below



Hence, min value of $\max\{x, x^2\}$ is 0.

69. (c) : Given, $f(x+y) = f(x)f(y)$

$$\begin{aligned} f'(3) &= \lim_{h \rightarrow 0} \frac{f(3+h) - f(3)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(3)f(h) - f(3)}{h} = f(3) \lim_{h \rightarrow 0} \frac{f(h) - 1}{h} \\ &= f(3) \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} \quad [\because f(0) = 1] \end{aligned}$$

$$\begin{aligned} &= f(3) \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = f(3) f'(0) \\ &= 3 \times 11 = 33 \quad \therefore f'(3) = 33 \end{aligned}$$

70. (a) : We have, $f'(9) = \lim_{x \rightarrow 9} \frac{f(x) - f(9)}{x - 9}$

$$\Rightarrow 0 = \lim_{x \rightarrow 9} \frac{f(x)}{x-9} \times \frac{\sqrt{f(x)} - 3}{\sqrt{f(x)} - 3}$$

$$\Rightarrow 0 = \lim_{x \rightarrow 9} \left\{ \frac{\sqrt{f(x)} - 3}{(\sqrt{x} - 3)(\sqrt{x} + 3)} \times \frac{f(x) + 9 - 9}{\sqrt{f(x)} - 3} \right\}$$

$$\begin{aligned} \Rightarrow 0 &= \lim_{x \rightarrow 9} \left(\frac{\sqrt{f(x)} - 3}{(\sqrt{x} - 3)} \right) \times \left[\lim_{x \rightarrow 9} \left\{ \frac{(\sqrt{f(x)})^2 - (3)^2}{(\sqrt{f(x)} - 3)(\sqrt{x} + 3)} \right\} \right. \\ &\quad \left. - \lim_{x \rightarrow 9} \left(\frac{9}{(\sqrt{f(x)} - 3)(\sqrt{x} + 3)} \right) \right] \end{aligned}$$

$$\Rightarrow 0 = \lim_{x \rightarrow 9} \left(\frac{\sqrt{f(x)} + 3}{\sqrt{x} - 3} \right) \times$$

$$\left[\lim_{x \rightarrow 9} \frac{\sqrt{f(x)} + 3}{\sqrt{x} + 3} - \frac{9}{(\sqrt{f(9)} - 3)(\sqrt{9} + 3)} \right]$$

$$\Rightarrow 0 = \lim_{x \rightarrow 9} \left(\frac{\sqrt{f(x)} + 3}{\sqrt{x} - 3} \right) \times \left[\frac{3}{6} + \frac{9}{3 \times 6} \right]$$

$$\Rightarrow \lim_{x \rightarrow 9} \left(\frac{\sqrt{f(x)} + 3}{\sqrt{x} - 3} \right) = 0$$

$$\mathbf{71. (e) :} \cos \left(\frac{\pi}{4} + x \right) + \cos \left(\frac{\pi}{4} - x \right)$$

$$\begin{aligned} &= \cos \frac{\pi}{4} \cos x - \sin \frac{\pi}{4} \sin x + \cos \frac{\pi}{4} \cos x \\ &\quad + \sin \frac{\pi}{4} \sin x \end{aligned}$$

$$= 2 \cos \frac{\pi}{4} \cos x = \frac{2}{\sqrt{2}} \cos x = \sqrt{2} \cos x$$

72. (d) : Area of triangle

$$\begin{aligned} &= \frac{1}{2} |(-2)(5+1) + 1(-1-2) + 6(2-5)| \\ &= \frac{1}{2} |(-2)(6) - 3 - 18| = \frac{1}{2} |-12 - 3 - 18| \\ &= \frac{1}{2} |-33| = \frac{33}{2} \text{ sq. units} \end{aligned}$$

73. (b) : Equation of line passing through (1, 0) and (-4, 1) is

$$\frac{y-0}{1-0} = \frac{x-1}{-4-1} \Rightarrow y = \frac{x-1}{-5}$$

or $x + 5y - 1 = 0$

Now, equation of line perpendicular to $x + 5y - 1 = 0$ is

$5x - y + \lambda = 0$ is for some constant λ

Also, $5x - y + \lambda = 0$ passes through $(-3, 5)$

$$\therefore 5(-3) - 5 + \lambda = 0 \Rightarrow \lambda = 20$$

Hence, $5x - y + 20 = 0$ is the required equation of line

74. (b) : Given expansion is $(1 + x^2)^5 (1 + x)^4$

$$= [{}^5C_0(x^2)^0 + {}^5C_1(x^2)^1 + {}^5C_2(x^2)^2 + {}^5C_3(x^2)^3 + {}^5C_4(x^2)^4 + {}^5C_5(x^2)^5] [{}^4C_0x^0 + {}^4C_1x + {}^4C_2x^2 + {}^4C_3x^3 + {}^4C_4x^4]$$

$$= [{}^5C_0 + {}^5C_1x^2 + {}^5C_2x^4 + {}^5C_3x^6 + {}^5C_4x^8 + {}^5C_5x^{10}]$$

$$[{}^4C_0 + {}^4C_1x + {}^4C_2x^2 + {}^4C_3x^3 + {}^4C_4x^4]$$

$$\therefore \text{Coefficient of } x^5 = {}^5C_1 \cdot {}^4C_3 + {}^5C_2 \cdot {}^4C_1$$

$$= 20 + 40 = 60$$

75. (e) : Given expansion is

$$(1 - 2x)^5 = {}^5C_0 - {}^5C_1(2x) + {}^5C_2(2x)^2 - {}^5C_3(2x)^3 + {}^5C_4(2x)^4 - {}^5C_5(2x)^5$$

$$\therefore \text{Coefficient of } x^4 = {}^5C_4 \cdot (2)^4 = 80$$

76. (a) : We have, $5x^2 + y^2 + y = 8$

$$\Rightarrow 5x^2 + y^2 + y - 8 + \frac{1}{4} - \frac{1}{4} = 0$$

$$\Rightarrow (\sqrt{5}x)^2 + (y + 1/2)^2 - \frac{33}{4} = 0$$

$$\Rightarrow (\sqrt{5}x)^2 + (y + 1/2)^2 = \left(\frac{\sqrt{33}}{2}\right)^2$$

which represents an ellipse.

77. (e) : We have, $4x^2 + y^2 - 8x + 4y - 8 = 0$

$$\Rightarrow 4(x^2 - 2x + 1 - 1) + (y^2 + 4y + 4 - 4) - 8 = 0$$


$$\Rightarrow 4(x - 1)^2 - 4 + (y + 2)^2 - 4 - 8 = 0$$

$$\Rightarrow \frac{(x-1)^2}{1/4} + (y+2)^2 = 16$$

$$\Rightarrow \frac{(x-1)^2}{4} + \frac{(y+2)^2}{16} = 1$$

$$\therefore \text{Centre} \equiv (1, -2).$$

78. (c) :



$$\text{Required area} = 2 \int_0^{\sqrt{3}} (-x^2 + 3) dx$$

$$= 2 \left[\frac{-x^3}{3} + 3x \right]_0^{\sqrt{3}} = 2 \left[\frac{-3\sqrt{3}}{3} + 3\sqrt{3} \right]$$

$$= 2(2\sqrt{3}) = 4\sqrt{3}$$

79. (a) : Order of differential equation

$$\left(\frac{d^3 y}{dx^3} \right)^2 + \left(\frac{d^2 y}{dx^2} \right)^2 + \left(\frac{dy}{dx} \right)^5 = 0 \text{ is } 3.$$

80. (a) : We have, $f(x) = \sqrt{2x} + \frac{4}{\sqrt{2x}}$

$$\therefore f'(x) = \sqrt{2} \cdot \frac{1}{2\sqrt{x}} + \frac{4}{\sqrt{2}} \cdot \left(\frac{-1}{2x^{3/2}} \right)$$

$$= \frac{1}{\sqrt{2x}} - \frac{\sqrt{2}}{(x)^{3/2}}$$

$$\Rightarrow f'(2) = \frac{1}{2} - \frac{1}{2} = 0$$

81. (b) : Let r be the radius of given circle.

Given, area of circle $x^2 - 2x + y^2 - 10y + k = 0$ is 25π

$$\text{i.e., } \pi r^2 = 25\pi \Rightarrow r^2 = 25 \quad \dots(i)$$

Also, radius from the given equation is

$$r = \sqrt{(1)^2 + (5)^2 - k} \Rightarrow r^2 = 26 - k$$

$$\Rightarrow 25 = 26 - k$$

[Using (i)]

$$\Rightarrow k = 1$$

$$\text{82. (d) : Let } I = \int_{2016}^{2017} \frac{\sqrt{x}}{\sqrt{x} + \sqrt{4033-x}} dx \quad \dots(i)$$

$$\text{Also, } I = \int_{2016}^{2017} \frac{\sqrt{4033-x}}{\sqrt{4033-x} + \sqrt{x}} dx \quad \dots(ii)$$

Adding (i) and (ii), we get

$$2I = \int_{2016}^{2017} \frac{(\sqrt{x} + \sqrt{4033-x})}{(\sqrt{x} + \sqrt{4033-x})} dx = \int_{2016}^{2017} 1 \cdot dx$$

$$\Rightarrow I = \frac{1}{2} [2017 - 2016] = \frac{1}{2}$$

83. (a) : We have, $\frac{dy}{dx} + y \tan x = \sec x$, $y(0) = 0$

This is a linear differential equation

$$\therefore \text{I.F.} = e^{\int \tan x \, dx} = e^{\log|\sec x|} = \sec x$$

$$\therefore \text{Solution is given by } y \cdot \sec x = \int \sec x \cdot \sec x \, dx$$

$$\Rightarrow y \cdot \sec x = \int \sec^2 x \, dx \Rightarrow y \sec x = \tan x + c$$

Now, we have $y(0) = 0$

$$\Rightarrow (0) \cdot \sec 0 = \tan(0) + c \Rightarrow c = 0$$

$$\therefore \text{Particular solution is, } y \sec x = \tan x$$

84. (e) : Since the vectors $2\hat{i} + 2\hat{j} + 6\hat{k}$, $2\hat{i} + \lambda\hat{j} + 6\hat{k}$, $2\hat{i} - 3\hat{j} + \hat{k}$ are coplanar.

$$\therefore \begin{vmatrix} 2 & 2 & 6 \\ 2 & \lambda & 6 \\ 2 & -3 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 2(\lambda + 18) - 2(2 - 12) + 6(-6 - 2\lambda) = 0$$

$$\Rightarrow -10\lambda + 20 = 0 \Rightarrow \lambda = 2$$

85. (b) : Distance of the point $(2, 1, 0)$ from the plane $2x + y + 2z + 5 = 0$ is given by

$$\left| \frac{(2 \cdot 2) + (1 \cdot 1) + (0 \cdot 2) + 5}{\sqrt{2^2 + 1^2 + 2^2}} \right| = \left| \frac{4 + 1 + 5}{3} \right| = \frac{10}{3}$$

86. (e) : Coordinates of vertices of hyperbola $(0, \pm 15) = (0, \pm b)$

and foci $(0, \pm 20) = (0, \pm be)$

$$\therefore b = 15 \text{ and } be = 20$$

$$\Rightarrow e = \frac{20}{15} = \frac{4}{3}$$

$$\text{Now, } e^2 = 1 + \frac{a^2}{b^2} \Rightarrow \frac{16}{9} = 1 + \frac{a^2}{225}$$

$$\Rightarrow a^2 = \frac{7}{9} \times 225 = 175$$

$$\therefore \text{Equation of hyperbola } \frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$$

$$\text{i.e., } \frac{y^2}{225} - \frac{x^2}{175} = 1$$

$$\text{87. (e) : Given, } \frac{15^3 + 6^3 + 3 \cdot 6 \cdot 15 \cdot 21}{1 + 4(6) + 6(36) + 4(216) + 1296}$$

$$= \frac{(15+6)^3}{(1+6)^4} = \frac{(21)^3}{(7)^4} = \frac{27}{7}$$

88. (a) : The equation of the plane passing through the points $(1, 0, 2)$, $(-1, 1, 2)$ and $(5, 0, 3)$ is

$$\begin{vmatrix} x-1 & y-0 & z-2 \\ -1-1 & 1-0 & 2-2 \\ 5-1 & 0-0 & 3-2 \end{vmatrix} = 0 \Rightarrow \begin{vmatrix} x-1 & y & z-2 \\ -2 & 1 & 0 \\ 4 & 0 & 1 \end{vmatrix} = 0$$

$$\Rightarrow (x-1)(1) - y(-2) + (z-2)(-4) = 0$$

$$\Rightarrow x - 1 + 2y - 4z + 8 = 0$$

$$\Rightarrow x + 2y - 4z + 7 = 0$$

89. (b) : Given equation is $y^2 - 4y - x + 3 = 0$

$$\Rightarrow (y-2)^2 - x - 1 = 0 \Rightarrow (y-2)^2 = x + 1$$

Now, shifting the origin to the point $(-1, 2)$ without rotating the axes and denoting the new coordinates w.r.t. these axes by X and Y , we get

$$y - 2 = Y \text{ and } x + 1 = X$$

Now, substituting $(X = 0, Y = 0)$

$$y = 2, x = -1$$

Hence, vertex w.r.t to old axes $(-1, 2)$.

90. (a) : Given, $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

$$\Rightarrow \vec{a} = -(\vec{b} + \vec{c}) \Rightarrow |\vec{a}|^2 = |-(\vec{b} + \vec{c})|^2$$

$$\Rightarrow |\vec{a}|^2 = |\vec{b}|^2 + |\vec{c}|^2 + 2|\vec{b}||\vec{c}|\cos\theta$$

(θ is angle between \vec{b} and \vec{c})

$$\Rightarrow 49 = 25 + 9 + 30 \cos \theta$$

$$\Rightarrow \cos \theta = \frac{15}{30} = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$$

91. (d) : Let $f(x) = 2x^3 - 9ax^2 + 12a^2x + 1$

$$\therefore f'(x) = 6x^2 - 18ax + 12a^2$$

$$\text{Now, } f'(x) = 0 \Rightarrow 6x^2 - 18ax + 12a^2 = 0$$

$$\Rightarrow (x - 2a)(x - a) = 0$$

$$\Rightarrow x = a \text{ or } 2a$$

$$\text{Also, } f''(x) = 12x - 18a$$

$$\therefore f''(a) = 12a - 18a = -6a < 0 \quad (\because a > 0)$$

$$\text{and } f''(2a) = 24a - 18a = 6a > 0$$

Hence, $p = a$ and $q = 2a$

$$\text{Now, } p^3 = q \Rightarrow a^3 = 2a \Rightarrow a(a^2 - 2) = 0$$

$$\Rightarrow a = \pm \sqrt{2}$$

92. (e) : We have, $f(x) + f(y) = f(x + y)$

Put $x = y = 1$, we get

$$f(1 + 1) = f(1) + f(1) \Rightarrow f(2) = 7 + 7 = 14$$

$$f(2 + 1) = f(2) + f(1) \Rightarrow f(3) = 14 + 7 = 21$$

Continuing in the same way, we get
 $f(4) = 28, f(5) = 35$ and so on.

$$\therefore \sum_{r=1}^{100} f(r) = f(1) + f(2) + f(3) + \dots + f(100)$$

$$= 7 + 14 + 21 + \dots + 700$$

Since, the series forms an A.P.

$$\therefore \sum_{r=1}^{100} f(r) = \frac{100}{2} [7 + 700] = 50 \times 707 = 50 \times 7 \times 101$$

93. (a) : Given, $\frac{(x-1)^2}{2} + \left(y + \frac{3}{4}\right)^2 = \frac{1}{16}$

or $\frac{(x-1)^2}{1/8} + \frac{(y+3/4)^2}{1/16} = 1$

$$\therefore \text{Eccentricity } (e) = \sqrt{1 - \frac{b^2}{a^2}} = \sqrt{1 - \frac{(1/16)}{(1/8)}}$$

$$= \sqrt{1 - \frac{1}{2}} = \frac{1}{\sqrt{2}}$$

94. (b) : We have, $\int_{-1}^1 \max\{x, x^3\} dx$

$$= \int_{-1}^0 x^3 dx + \int_0^1 x dx = \left[\frac{x^4}{4} \right]_{-1}^0 + \left[\frac{x^2}{2} \right]_0^1 = \frac{1}{4}$$

95. (d) : We have, $\sin x + \cos y = 2$

It is possible only if $\sin x = 1$ and $\cos y = 1$

$$\Rightarrow x = \frac{\pi}{2} \text{ and } y = 0 \therefore x + y = \frac{\pi}{2} + 0 = \frac{\pi}{2}$$

96. (a) : Given numbers $a, a + r, a + 2r$ are in A.P.
 Also their product = 64. This is possible only when three numbers are equal to 4.

i.e., $a = a + r = a + 2r = 4$

\therefore Minimum value of $a + 2r$ is 4.

97. (b) : We have, $S = \frac{1}{9!} + \frac{1}{3!7!} + \frac{1}{5!5!} + \frac{1}{7!3!} + \frac{1}{9!}$

$$= \frac{1}{9!} \left[2 + \frac{2 \times 8 \times 9}{3!} + \frac{9 \times 8 \times 7 \times 6}{5!} \right]$$

$$= \frac{1}{9!} \left[\frac{2 \times 120 + 2(72)(20) + (3024)}{120} \right] = \frac{1 \times 6144}{10! \times 12} = \frac{2^9}{10!}$$

98. (d) : Given, $f(x) = \begin{vmatrix} x & x^2 & x^3 \\ 1 & 2x & 3x^2 \\ 0 & 2 & 6x \end{vmatrix}$

$$= x(12x^2 - 6x^2) - x^2(6x) + x^3(2 - 0)$$

$$= 6x^3 - 6x^3 + 2x^3 = 2x^3 \therefore f'(x) = 6x^2$$

99. (c) : Let $I = \int \frac{x^2}{1+(x^3)^2} dx$

Put $x^3 = t \Rightarrow 3x^2 dx = dt$

$$\therefore I = \frac{1}{3} \int \frac{dt}{1+t^2} = \frac{1}{3} [\tan^{-1} t] + c = \frac{1}{3} \tan^{-1}(x^3) + c$$

100. (d) : We have, $f(x) = x^2 + e^x$

$$\therefore f'(x) = 2x + e^x$$

$$f''(x) = 2 + e^x$$

$$f'''(x) = e^x$$

Hence, the least value of n so that $f_n = f_{n+1}$ is 3.

101. (e) : $\sin 765^\circ = \sin(2 \times 360^\circ + 45^\circ)$

$$= \sin 45^\circ = \frac{1}{\sqrt{2}}$$

102. (d) : Distance of the point $(3, -5)$ from the line $3x - 4y - 26 = 0$ is

$$\left| \frac{3 \cdot 3 + (-4)(-5) - 26}{\sqrt{3^2 + 4^2}} \right| = \left| \frac{9 + 20 - 26}{5} \right| = \frac{3}{5}$$

103. (c) : Given, $f(x) = \int_0^x (t^2 + t + 1) dt = \left[\frac{t^3}{3} + \frac{t^2}{2} + t \right]_0^x$

$$\therefore f(x) = \left[\frac{x^3}{3} + \frac{x^2}{2} + x \right]$$

$$\therefore f(x) = \left[\frac{x^3}{3} + \frac{x^2}{2} + x \right] \text{ on } [2, 3] \text{ gives}$$

$$f(2) = \frac{8}{3} + \frac{4}{2} + 2 = \frac{20}{3} \quad (\text{minimum})$$

$$\text{and } f(3) = \frac{27}{3} + \frac{9}{2} + 3 = \frac{33}{2} \quad (\text{maximum})$$

\therefore Difference between the maximum and minimum value is $\frac{33}{2} - \frac{20}{3} = \frac{59}{6}$

104. (c) : Let, $f(x) = x^2 + ax + b = 0$

Since a and b are roots of $f(x)$

$$\therefore a + b = -a \text{ and } ab = b \Rightarrow a = 1$$

$$\therefore 1 + b = -1 \Rightarrow b = -2$$

So, $f(x) = x^2 + x - 2$

Also, $f'(x) = 2x + 1$. For maximum/minimum $f'(x) = 0$

$$\Rightarrow x = -\frac{1}{2}$$

Now, $f''(x) = 2 > 0$

FIND
MORE
FREE
MAGAZINES

FREEMAGS.CC

$\therefore x = \frac{-1}{2}$ is the minimum point.

$$\begin{aligned}\therefore \text{Minimum value} &= f\left(-\frac{1}{2}\right) = \left(\frac{-1}{2}\right)^2 + \left(\frac{-1}{2}\right) - 2 \\ &= \frac{1}{4} - \frac{1}{2} - 2 = \frac{-9}{4}\end{aligned}$$

105. (b) : We have, $y = 4x + c$ and $\frac{x^2}{4} + y^2 = 1$

$\therefore y = 4x + c$ touches the given ellipse

$$\therefore \frac{x^2}{4} + (4x + c)^2 = 1$$

$$\Rightarrow x^2 + 4(16x^2 + c^2 + 8xc) = 4$$

$$\Rightarrow x^2 + 64x^2 + 4c^2 + 32xc - 4 = 0$$

$$\Rightarrow 65x^2 + 32xc + 4c^2 - 4 = 0$$

$$\text{Now, discriminant } D = 0 \Rightarrow (32c)^2 - 4(65)(4c^2 - 4) = 0$$

$$\Rightarrow 1024c^2 - 1040c^2 + 1040 = 0$$

$$\Rightarrow 16c^2 = 1040 \Rightarrow c^2 = 65$$

$$\Rightarrow c = \pm\sqrt{65}$$

106. (d) : Given equations $\lambda x - y = 2$, $2x - 3y = -\lambda$ and $3x - 2y = -1$ are consistent

$$\therefore \begin{vmatrix} \lambda & -1 & -2 \\ 2 & -3 & \lambda \\ 3 & -2 & 1 \end{vmatrix} = 0$$

$$\Rightarrow \lambda(-3 + 2\lambda) + 1(2 - 3\lambda) - 2(-4 + 9) = 0$$


$$\Rightarrow -3\lambda + 2\lambda^2 + 2 - 3\lambda + 8 - 18 = 0$$

$$\Rightarrow \lambda^2 - 3\lambda - 4 = 0 \Rightarrow (\lambda - 4)(\lambda + 1) = 0$$

$$\therefore \lambda = -1, 4$$

107. (a) : The set $\{(x, y) : |x| + |y| = 1\}$ in xy plane represents

$$\begin{cases} x + y = 1; & x > 0, y > 0 \\ x - y = 1; & x > 0, y < 0 \\ -x + y = 1; & x < 0, y > 0 \\ -x - y = 1; & x < 0, y < 0 \end{cases}$$



Hence, the given set represents a square.

108. (a) : Let $\tan^{-1}\left(\frac{3}{4}\right) = \theta \Rightarrow \tan \theta = \frac{3}{4}$

$$\therefore \cos \theta = \frac{4}{5}$$

$$\text{Now, } \cos\left(\tan^{-1}\left(\frac{3}{4}\right)\right) = \cos \theta = \frac{4}{5}$$

109. (b) : $\therefore AB = BC$

\therefore By using distance formula, we have

$$\sqrt{(1-6)^2 + (3+1)^2} = \sqrt{(x-1)^2 + (8-3)^2}$$

On squaring both sides, we get

$$(-5)^2 + (4)^2 = (x-1)^2 + (5)^2$$

$$\Rightarrow x^2 + 1 - 2x = 16 \Rightarrow x^2 - 2x - 15 = 0$$

$$\Rightarrow (x+3)(x-5) = 0 \Rightarrow x = -3, 5$$

110. (e) : We have, $n = 15$, Incorrect $\Sigma x^2 = 2830$,

$$\text{Incorrect } \Sigma(x) = 170$$

$$\begin{aligned}\therefore \text{Correct } \Sigma(x) &= (\text{Incorrect } \Sigma x - \text{Incorrect value}) \\ &\quad + \text{Correct value} \\ &= (170 - 20) + 30 = 180\end{aligned}$$

$$\therefore \text{Correct Mean} = \text{Correct } \frac{\Sigma x}{15} = \frac{180}{15} = 12$$

$$\begin{aligned}\text{Similarly, Correct } \Sigma x^2 &= \text{Incorrect } \Sigma x^2 - (\text{Incorrect value})^2 \\ &\quad + (\text{Correct value})^2 \\ &= 2830 - (20)^2 + (30)^2 = 2830 + 500 = 3330\end{aligned}$$

$$\begin{aligned}\therefore \text{Correct variance} &= \text{Correct } \frac{(\Sigma x^2)}{n} - (\text{Correct mean})^2 \\ &= \frac{3330}{15} - (12)^2 = 222 - 144 = 78\end{aligned}$$

111. (e) : Let θ be the angle between the lines

$$\frac{x-2}{2} = \frac{y-1}{5} = \frac{z+3}{-3} \text{ and } \frac{x+2}{-1} = \frac{y-4}{8} = \frac{z-5}{4}$$

$$\begin{aligned}\therefore \cos \theta &= \frac{|2(-1) + 5(8) + (-3)(4)|}{\sqrt{(2)^2 + (5)^2 + (-3)^2} \sqrt{(-1)^2 + (8)^2 + (4)^2}} \\ &= \frac{|-2 + 40 - 12|}{\sqrt{38}\sqrt{81}} = \frac{26}{9\sqrt{38}}\end{aligned}$$

$$\text{i.e., } \theta = \cos^{-1}\left(\frac{26}{9\sqrt{38}}\right)$$


112. (d) : We have, $(\vec{x} - \vec{a}) \cdot (\vec{x} + \vec{a}) = 12$

$$\Rightarrow |\vec{x}|^2 - |\vec{a}|^2 = 12$$

Since, \vec{a} be a unit vector $\therefore |\vec{a}| = 1$

$$\Rightarrow |\vec{x}|^2 - 1 = 12 \Rightarrow |\vec{x}|^2 = 13 \Rightarrow |\vec{x}| = \sqrt{13}$$

113. (d) : Sides of triangular region are $y = 2x + 1$, $y = 3x + 1$ and $x = 4$.



∴ Area of shaded region i.e., ΔABC where
 $A = (0, 1)$, $B = (4, 9)$, $C = (4, 13)$
 $= \frac{1}{2} |0(9 - 13) + 4(13 - 1) + 4(1 - 9)| = \frac{1}{2} |4(12) - 32|$
 $= \frac{1}{2} \times 16 = 8 \text{ sq. units}$

114. (b) : Given, ${}^nC_{r-1} = 36$, ${}^nC_r = 84$, ${}^nC_{r+1} = 126$

Since, $\frac{{}^nC_r}{{}^nC_{r-1}} = \frac{n-r+1}{r}$

∴ $\frac{84}{36} = \frac{n-r+1}{r} \Rightarrow \frac{n-r+1}{r} = \frac{7}{3}$
 $\Rightarrow 7r = 3n - 3r + 3 \Rightarrow 10r = 3n + 3 \quad \dots(i)$

Also, $\frac{{}^nC_r}{{}^nC_{r+1}} = \frac{r+1}{n-r} \Rightarrow \frac{84}{126} = \frac{r+1}{n-r} \Rightarrow \frac{r+1}{n-r} = \frac{2}{3}$
 $\Rightarrow 3r + 3 = 2n - 2r \Rightarrow 5r = 2n - 3 \quad \dots(ii)$

Solving (i) & (ii), we get $r = 3$

115. (e) : $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 $= \lim_{h \rightarrow 0} \frac{f(x)f(h) - f(x)}{h} \quad [\because f(x+y) = f(x)f(y)]$
 $= f(x) \lim_{h \rightarrow 0} \frac{f(h) - 1}{h} = f(x) \lim_{h \rightarrow 0} \frac{1 + \sin 3h g(h) - 1}{h}$
 $= f(x) \lim_{h \rightarrow 0} g(h) \frac{\sin 3h}{h}$
 $= f(x) g(0) \lim_{h \rightarrow 0} \frac{\sin 3h}{3h} \times 3 = 3 f(x) g(0)$

116. (b) : Given, $\begin{vmatrix} x-1 & 1 & 1 \\ 1 & x-1 & 1 \\ 1 & 1 & x-1 \end{vmatrix} = 0$

$\Rightarrow (x-1)[(x-1)^2 - 1] - 1[x-1-1] + 1[1-x+1] = 0$
 $\Rightarrow (x-1)[x^2 + 1 - 2x - 1] - x + 2 + 2 - x = 0$
 $\Rightarrow (x-1)x(x-2) - 2x + 4 = 0$
 $\Rightarrow x(x-1)(x-2) - 2(x-2) = 0$
 $\Rightarrow (x-2)[x^2 - x - 2] = 0 \Rightarrow (x-2)(x-2)(x+1) = 0$
 $\Rightarrow x = 2, -1$

117. (None of the options is correct) :

Given, $T_7 = T_8$
 $\Rightarrow {}^nC_6(2a)^{n-6}(-3b)^6 = {}^nC_7(2a)^{n-7}(-3b)^7$
 $\Rightarrow \frac{n!}{(n-6)!6!}(2a)^{n-6}(-3b)^6 = \frac{n!}{(n-7)!7!}(2a)^{n-7}(-3b)^7$
 $\Rightarrow \frac{1}{(n-6)}(2a) = \frac{(-3b)}{7} \Rightarrow \frac{2a}{3b} = \frac{-(n-6)}{7}$

Applying componendo & dividendo, we get

$\frac{2a+3b}{2a-3b} = \frac{6-n+7}{6-n-7} = \frac{13-n}{-n-1} = \frac{n-13}{n+1}$

118. (c) : Mean of first n odd natural numbers

$\bar{x} = \frac{1+3+5+\dots+(2n-1)}{n} = \frac{n^2}{n} = n$

Sum of square of first n odd natural numbers i.e.,

$1^2 + 3^2 + 5^2 + \dots + (2n-1)^2 = \frac{n(2n+1)(2n-1)}{3}$

∴ Standard deviation $= \sqrt{\frac{n(2n+1)(2n-1)}{3n} - n^2}$
 $= \sqrt{\frac{n^2-1}{3}}$


119. (b) : Given, $S = \{1, 2, 3, \dots, 10\}$.

∴ Set containing odd numbers of $S = \{1, 3, 5, 7, 9\}$

∴ Number of subsets of S containing only odd numbers $= (2)^5 - 1 = 32 - 1 = 31$

120. (e) : Let the vertices of parallelogram be

$A(0, 0)$, $B(7, 2)$, $C(5, 9)$ and $D(12, 11)$



Area of || gm ABCD = area of ΔABC + area of ΔADC

Now, area of $\Delta ABC = \frac{1}{2} |0(2-9) + 7(9-0) + 5(0-2)|$
 $= \frac{1}{2} |63 - 10| = \frac{53}{2}$

∴ Area of $\Delta ADC = \frac{1}{2} |0(11-9) + 12(9-0) + 5(0-11)|$
 $= \frac{1}{2} |108 - 55| = \frac{53}{2}$

∴ Area of || gm ABCD $= \frac{53}{2} + \frac{53}{2} = 53 \text{ sq. units}$



Karnataka CET

- The distance of the point $(-2, 4, -5)$ from the line $\frac{x+3}{3} = \frac{y-4}{5} = \frac{z+8}{6}$ is
(a) $\frac{\sqrt{37}}{10}$ (b) $\frac{37}{\sqrt{10}}$ (c) $\sqrt{\frac{37}{10}}$ (d) $\frac{37}{10}$
- If A is a square matrix of order 3×3 , then $|KA|$ is equal to
(a) $K|A|$ (b) $K^2|A|$
(c) $3K|A|$ (d) $K^3|A|$
- Equation of line passing through the point $(1, 2)$ and perpendicular to the line $y = 3x - 1$ is
(a) $x - 3y = 0$ (b) $x + 3y = 0$
(c) $x + 3y - 7 = 0$ (d) $x + 3y + 7 = 0$
- General solution of differential equation $\frac{dy}{dx} + y = 1 (y \neq 1)$ is
(a) $\log \left| \frac{1}{1-y} \right| = x + C$ (b) $\log |1-y| = x + C$
(c) $\log |1+y| = x + C$ (d) $\log \left| \frac{1}{1-y} \right| = -x + C$
- The value of C in mean value theorem for the function $f(x) = x^2$ in $[2, 4]$ is
(a) 2 (b) 4 (c) $\frac{7}{2}$ (d) 3
- The value of $\lim_{\theta \rightarrow 0} \frac{1 - \cos 4\theta}{1 - \cos 6\theta}$ is
(a) $\frac{9}{4}$ (b) $\frac{4}{9}$ (c) $\frac{9}{3}$ (d) $\frac{3}{4}$
- If $y = \tan^{-1} \left(\frac{\sin x + \cos x}{\cos x - \sin x} \right)$, then $\frac{dy}{dx}$ is equal to
(a) 0 (b) $\frac{1}{2}$ (c) $\frac{\pi}{4}$ (d) 1
- If $\left(\frac{1+i}{1-i} \right)^m = 1$, then the least positive integral value of m is
(a) 4 (b) 1 (c) 2 (d) 3
- $\int_{-5}^5 |x+2| dx$ is equal to
(a) 28 (b) 29 (c) 27 (d) 30
- $\int \frac{\cos 2x - \cos 2\theta}{\cos x - \cos \theta} dx$ is equal to
(a) $2(\sin x + x \cos \theta) + C$
(b) $2(\sin x - x \cos \theta) + C$
(c) $2(\sin x + 2x \cos \theta) + C$
(d) $2(\sin x - 2x \cos \theta) + C$
- The area of the region bounded by the curve $y = x^2$ and the line $y = 16$ is
(a) $\frac{256}{3}$ sq. units (b) $\frac{128}{3}$ sq. units
(c) $\frac{32}{3}$ sq. units (d) $\frac{64}{3}$ sq. units
- If A and B are finite sets and $A \subset B$, then
(a) $n(A \cup B) = n(B)$ (b) $n(A \cap B) = n(B)$
(c) $n(A \cap B) = \phi$ (d) $n(A \cup B) = n(A)$
- If a matrix A is both symmetric and skew symmetric, then
(a) A is diagonal matrix
(b) A is a zero matrix
(c) A is scalar matrix
(d) A is square matrix
- If $\begin{vmatrix} 3 & x \\ x & 1 \end{vmatrix} = \begin{vmatrix} 3 & 2 \\ 4 & 1 \end{vmatrix}$ then x is equal to
(a) 8 (b) 4
(c) $\pm 2\sqrt{2}$ (d) 2

15. The integrating factor of the differential equation


$$x \cdot \frac{dy}{dx} + 2y = x^2 \text{ is } (x \neq 0)$$

- (a) $e^{\log x}$ (b) $\log|x|$
(c) x (d) x^2

16. The perpendicular distance of the point $P(6, 7, 8)$ from XY -plane is

- (a) 7 (b) 6 (c) 8 (d) 5

17. The shaded region in the figure is the solution set of the inequations



- (a) $5x + 4y \leq 20, x \leq 6, y \leq 3, x \geq 0, y \geq 0$
(b) $5x + 4y \geq 20, x \leq 6, y \geq 3, x \geq 0, y \geq 0$
(c) $5x + 4y \geq 20, x \leq 6, y \leq 3, x \geq 0, y \geq 0$
(d) $5x + 4y \geq 20, x \geq 6, y \leq 3, x \geq 0, y \geq 0$

18. If an LPP admits optimal solution at two consecutive vertices of a feasible region, then

- (a) the required optimal solution is at the midpoint of the line joining two points.
(b) the optimal solution occurs at every point on the line joining these two points.
(c) the LPP under consideration is not solvable.
(d) the LPP under consideration must be reconstructed.

19. $3 + 5 + 7 + \dots$ to n terms is

- (a) n^2 (b) $n(n-2)$
(c) $n(n+2)$ (d) $(n+1)^2$

20. If $2 \begin{bmatrix} 1 & 3 \\ 0 & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$, then the value of x and y are

- (a) $x = 3, y = 3$ (b) $x = -3, y = 3$
(c) $x = 3, y = -3$ (d) $x = -3, y = -3$

21. The derivative of $\cos^{-1}(2x^2 - 1)$ w.r.t $\cos^{-1} x$ is

- (a) 2 (b) $\frac{2}{x}$
(c) $1 - x^2$ (d) $\frac{-1}{2\sqrt{1-x^2}}$

22. A box has 100 pens of which 10 are defective. The probability that out of a sample of 5 pens drawn one by one with replacement and atmost one is defective is

- (a) $\frac{9}{10}$ (b) $\frac{1}{2} \left(\frac{9}{10} \right)^4$
(c) $\left(\frac{9}{10} \right)^5 + \frac{1}{2} \left(\frac{9}{10} \right)^4$ (d) $\frac{1}{2} \left(\frac{9}{10} \right)^5$

23. If $y = \log(\log x)$, then $\frac{d^2 y}{dx^2}$ is equal to

- (a) $\frac{(1 + \log x)}{x^2 \log x}$ (b) $\frac{-(1 + \log x)}{(x \log x)^2}$
(c) $\frac{(1 + \log x)}{(x \log x)^2}$ (d) $\frac{-(1 + \log x)}{x^2 \log x}$

24. $\int \frac{(x+3)e^x}{(x+4)^2} dx$ is equal to

- (a) $\frac{e^x}{(x+4)} + C$ (b) $\frac{e^x}{(x+4)^2} + C$
(c) $\frac{e^x}{(x+3)} + C$ (d) $\frac{1}{(x+4)^2} + C$

25. $\int_0^{\pi/2} \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x} dx$ is equal to

- (a) $\frac{\pi a}{4b}$ (b) $\frac{\pi b}{4a}$ (c) $\frac{\pi}{2ab}$ (d) $\frac{\pi a}{2b}$

26. Let $f: R \rightarrow R$ be defined by $f(x) = x^4$, then

- (a) f is one-one but not onto
(b) f is neither one-one nor onto
(c) f is one-one and onto
(d) f may be one-one and onto

27. The point on the curve $y^2 = x$ where the tangent makes an angle $\frac{\pi}{4}$ with X -axis is

- (a) $(1, 1)$ (b) $\left(\frac{1}{4}, \frac{1}{2} \right)$
(c) $\left(\frac{1}{2}, \frac{1}{4} \right)$ (d) $(4, 2)$

28. The total number of terms in the expansion of $(x+a)^{47} - (x-a)^{47}$ after simplification is

- (a) 24 (b) 96 (c) 47 (d) 48

29. The function $f(x) = x^2 + 2x - 5$ is strictly increasing in the interval

(a) $[-1, \infty)$ (b) $(-\infty, -1)$
(c) $(-\infty, -1]$ (d) $(-1, \infty)$

30. The degree of the differential equation

$$\left[1 + \left(\frac{dy}{dx}\right)^2\right]^2 = \frac{d^2y}{dx^2} \text{ is}$$

(a) 1 (b) 4 (c) 2 (d) 3

31. Binary operation $*$ on $R - \{-1\}$ defined by

$$a * b = \frac{a}{b+1} \text{ is}$$

(a) $*$ is associative and commutative
(b) $*$ is neither associative nor commutative
(c) $*$ is commutative but not associative
(d) $*$ is associative but not commutative

32. The plane $2x - 3y + 6z - 11 = 0$ makes an angle $\sin^{-1}(\alpha)$ with X -axis. The value of α is equal to

(a) $\frac{\sqrt{3}}{2}$ (b) $\frac{2}{7}$ (c) $\frac{\sqrt{2}}{3}$ (d) $\frac{3}{7}$

33. If coefficient of variation is 60 and standard deviation is 24, then arithmetic mean is

(a) $\frac{20}{7}$ (b) $\frac{7}{20}$ (c) $\frac{1}{40}$ (d) 40

34. The contrapositive statement of the statement "If x is prime number, then x is odd" is

(a) If x is not a prime number, then x is odd.
(b) If x is not a prime number, then x is not odd.
(c) If x is a prime number, then x is not odd.
(d) If x is not odd, then x is not a prime number.

35. The probability distribution of X is

X	0	1	2	3
$P(X)$	0.3	k	$2k$	$2k$

The value of k is

(a) 0.7 (b) 0.3 (c) 1 (d) 0.14

36. $\int \sqrt{x^2 + 2x + 5} dx$ is equal to

(a) $(x+1)\sqrt{x^2 + 2x + 5} - 2\log|x+1 + \sqrt{x^2 + 2x + 5}| + C$
(b) $\frac{1}{2}(x+1)\sqrt{x^2 + 2x + 5} + 2\log|x+1 + \sqrt{x^2 + 2x + 5}| + C$

$$(c) (x+1)\sqrt{x^2 + 2x + 5} + 2\log|x+1 + \sqrt{x^2 + 2x + 5}| + C$$

$$(d) (x+1)\sqrt{x^2 + 2x + 5} + \frac{1}{2}\log|x+1 + \sqrt{x^2 + 2x + 5}| + C$$

37. If ${}^nC_{12} = {}^nC_8$ then n is equal to

(a) 12 (b) 26 (c) 6 (d) 20

38. If $y = \begin{vmatrix} f(x) & g(x) & h(x) \\ l & m & n \\ a & b & c \end{vmatrix}$, then $\frac{dy}{dx}$ is equal to

$$(a) \begin{vmatrix} f'(x) & g'(x) & h'(x) \\ l & m & n \\ a & b & c \end{vmatrix}$$

$$(b) \begin{vmatrix} l & m & n \\ f'(x) & g'(x) & h'(x) \\ a & b & c \end{vmatrix}$$

$$(c) \begin{vmatrix} f'(x) & l & a \\ g'(x) & m & b \\ h'(x) & n & c \end{vmatrix} \quad (d) \begin{vmatrix} l & m & n \\ a & b & c \\ f'(x) & g'(x) & h'(x) \end{vmatrix}$$

39. If $\tan^{-1}x + \tan^{-1}y = \frac{4\pi}{5}$, then $\cot^{-1}x + \cot^{-1}y$ is equal to

(a) $\frac{\pi}{5}$ (b) $\frac{3\pi}{5}$ (c) $\frac{2\pi}{5}$ (d) π

40. The range of the function $f(x) = \sqrt{9 - x^2}$ is

(a) $[0, 3]$ (b) $(0, 3]$
(c) $(0, 3)$ (d) $[0, 3)$

41. Two events A and B will be independent if

(a) $P(A' \cap B') = (1 - P(A))(1 - P(B))$
(b) A and B are mutually exclusive
(c) $P(A) + P(B) = 1$
(d) $P(A) = P(B)$

42. The eccentricity of the ellipse $\frac{x^2}{36} + \frac{y^2}{16} = 1$ is

(a) $\frac{2\sqrt{5}}{6}$ (b) $\frac{2\sqrt{13}}{4}$ (c) $\frac{2\sqrt{5}}{4}$ (d) $\frac{2\sqrt{13}}{6}$

43. If \vec{a} & \vec{b} are unit vectors, then angle between \vec{a} and \vec{b} for $\sqrt{3}\vec{a} - \vec{b}$ to be unit vector is
(a) 45° (b) 30° (c) 90° (d) 60°

44. If $\vec{a} = 2\hat{i} + \lambda\hat{j} + \hat{k}$ and $\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$ are orthogonal, then value of λ is

(a) $\frac{3}{2}$ (b) 1 (c) $-\frac{5}{2}$ (d) 0

45. The value of $\cos^2 45^\circ - \sin^2 15^\circ$ is

(a) $\frac{\sqrt{3}+1}{2\sqrt{2}}$ (b) $\frac{\sqrt{3}}{4}$
(c) $\frac{\sqrt{3}-1}{2\sqrt{2}}$ (d) $\frac{\sqrt{3}}{2}$

46. The range of $\sec^{-1}x$ is

(a) $[0, \pi]$ (b) $[0, \pi] - \left\{\frac{\pi}{2}\right\}$
(c) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (d) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

47. If $\vec{a}, \vec{b}, \vec{c}$ are unit vectors such that $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, then the value of $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$ is equal to

(a) $-\frac{3}{2}$ (b) 3 (c) $\frac{3}{2}$ (d) 1

48. $\int_{-\pi/2}^{\pi/2} \frac{dx}{e^{\sin x} + 1}$ is equal to

(a) 1 (b) 0 (c) $\frac{\pi}{2}$ (d) $-\frac{\pi}{2}$

49. The rate of change of volume of a sphere with respect to its surface area when the radius is 4 cm is

(a) $2 \text{ cm}^3/\text{cm}^2$ (b) $4 \text{ cm}^3/\text{cm}^2$
(c) $8 \text{ cm}^3/\text{cm}^2$ (d) $6 \text{ cm}^3/\text{cm}^2$

50. $\int_0^{\pi/2} \frac{\tan^7 x}{\cot^7 x + \tan^7 x} dx$ is equal to

(a) $\frac{\pi}{4}$ (b) $\frac{\pi}{2}$ (c) $\frac{\pi}{6}$ (d) $\frac{\pi}{3}$

51. If $|x-2| \leq 1$, then

(a) $x \in (1, 3)$ (b) $x \in (-1, 3)$
(c) $x \in [1, 3]$ (d) $x \in [-1, 3]$

52. $\int_{0.2}^{3.5} [x] dx$ is equal to

(a) 3.5 (b) 4.5 (c) 3 (d) 4

53. The area of triangle with vertices $(K, 0), (4, 0), (0, 2)$ is 4 square units, then value of K is

(a) 8 (b) 0 or -8 (c) 0 (d) 0 or 8

54. If $f(x) = \begin{cases} Kx^2 & \text{if } x \leq 2 \\ 3 & \text{if } x > 2 \end{cases}$ is continuous at $x = 2$, then the value of K is

(a) $\frac{4}{3}$ (b) $\frac{3}{4}$ (c) 3 (d) 4

55. If $A = \frac{1}{\pi} \begin{bmatrix} \sin^{-1}(\pi x) & \tan^{-1}\left(\frac{x}{\pi}\right) \\ \sin^{-1}\left(\frac{x}{\pi}\right) & \cot^{-1}(\pi x) \end{bmatrix}$

$$B = \begin{bmatrix} -\cos^{-1}(\pi x) & \tan^{-1}\left(\frac{x}{\pi}\right) \\ \sin^{-1}\left(\frac{x}{\pi}\right) & -\tan^{-1}(\pi x) \end{bmatrix}$$

then $A - B$ is equal to

(a) $\frac{1}{2}I$ (b) I (c) O (d) $2I$

56. If $f(x) = 8x^3, g(x) = x^{1/3}$, then $f \circ g(x)$ is

(a) 8^3x (b) $8x^3$
(c) $8x$ (d) $(8x)^{1/3}$

57. Let $\Delta = \begin{vmatrix} Ax & x^2 & 1 \\ By & y^2 & 1 \\ Cz & z^2 & 1 \end{vmatrix}$ and $\Delta_1 = \begin{vmatrix} A & B & C \\ x & y & z \\ zy & zx & xy \end{vmatrix}$ then

(a) $\Delta_1 = -\Delta$ (b) $\Delta_1 = \Delta$
(c) $\Delta_1 = 2\Delta$ (d) $\Delta_1 \neq \Delta$

58. If $\sin x = \frac{2t}{1+t^2}, \tan y = \frac{2t}{1-t^2}$, then $\frac{dy}{dx}$ is equal to

(a) 1 (b) -1 (c) 2 (d) 0

59. Reflection of the point (α, β, γ) in XY plane is

(a) $(0, 0, \gamma)$ (b) $(\alpha, \beta, -\gamma)$
(c) $(-\alpha, -\beta, \gamma)$ (d) $(\alpha, \beta, 0)$

60. Area of the region bounded by the curve $y = \cos x$, $x = 0$ and $x = \pi$ is

(a) 2 sq. units (b) 3 sq. units
(c) 4 sq. units (d) 1 sq. units

MPP-2 CLASS XI

ANSWER KEY

1. (c) 2. (a) 3. (d) 4. (a) 5. (b)
6. (c) 7. (b) 8. (a,c) 9. (b,c) 10. (d)
11. (b,c) 12. (a,c,d) 13. (a) 14. (a) 15. (b)
16. (d) 17. (4) 18. (5) 19. (1) 20. (1)


SOLUTIONS

1. (c): Given, point is $A(-2, 4, -5)$,

Line (l) is $\frac{x+3}{3} = \frac{y-4}{5} = \frac{z+8}{6} = \lambda$ (say)

Co-ordinates of B are $(3\lambda - 3, 5\lambda + 4, 6\lambda - 8)$

\therefore Direction ratios of AB are $(3\lambda - 1, 5\lambda, 6\lambda - 3)$



Now, $(3\lambda - 1)(3) + (5\lambda)(5) + (6\lambda - 3)(6) = 0$

$\Rightarrow \lambda = \frac{3}{10}$

$\therefore \overrightarrow{AB} = -\frac{1}{10}\hat{i} + \frac{3}{2}\hat{j} - \frac{6}{5}\hat{k}$

$\therefore d = |\overrightarrow{AB}| = \sqrt{\frac{1}{100} + \frac{9}{4} + \frac{36}{25}}$
 $= \sqrt{\frac{1+225+144}{100}} = \sqrt{\frac{370}{100}} = \sqrt{\frac{37}{10}}$

2. (d): We have $|KA| = K^n|A|$, Here $n = 3$

$\therefore |KA| = K^3|A|$

3. (c): Equation of required line is $y - 2 = -\frac{1}{3}(x - 1)$

$\Rightarrow x + 3y - 7 = 0$

4. (a): Given, $\frac{dy}{dx} + y = 1$

This is a linear differential equation.

\therefore I.F. $= e^{\int 1 dx} = e^x$

\therefore Solution is given by, $ye^x = \int e^x \cdot 1 dx = e^x + C_1$

$\Rightarrow e^x(y - 1) = C_1 \Rightarrow x + \log|y - 1| = \log C_1$

$\Rightarrow -x - \log|y - 1| = -\log C_1$

$\Rightarrow \log\left|\frac{1}{y-1}\right| = x + C$ [where $-\log C_1 = C$]

or $\log\left|\frac{1}{1-y}\right| = x + C$

5. (d): We have, $f(x) = x^2$ in $[2, 4]$

\therefore According to mean value theorem,

We have, $f'(C) = \frac{f(b) - f(a)}{b - a}$

[where $a = 2$ and $b = 4$]

$\therefore 2C = \frac{f(4) - f(2)}{4 - 2} = \frac{(4)^2 - (2)^2}{2} = \frac{12}{2}$

$\Rightarrow 2C = 6 \Rightarrow C = 3$

6. (b): We have, $\lim_{\theta \rightarrow 0} \frac{1 - \cos 4\theta}{1 - \cos 6\theta}$

$= \lim_{\theta \rightarrow 0} \frac{\sin^2 2\theta}{\sin^2 3\theta} = \frac{(2)^2}{(3)^2} = \frac{4}{9}$

7. (d): We have, $y = \tan^{-1}\left(\frac{\sin x + \cos x}{\cos x - \sin x}\right)$

$= \tan^{-1}\left(\frac{1 + \tan x}{1 - \tan x}\right) = \tan^{-1}\left(\tan\left(\frac{\pi}{4} + x\right)\right) = \frac{\pi}{4} + x$

$\therefore \frac{dy}{dx} = 1$

8. (a): Given, $\left(\frac{1+i}{1-i}\right)^m = 1 \Rightarrow i^m = i^4 \Rightarrow m = 4$

9. (b): Let $I = \int_{-5}^5 |x + 2| dx$

$= -\int_{-5}^{-2} (x + 2) dx + \int_{-2}^5 (x + 2) dx$
 $= -\left[\frac{x^2}{2} + 2x\right]_{-5}^{-2} + \left[\frac{x^2}{2} + 2x\right]_{-2}^5$
 $= \frac{9}{2} + \frac{49}{2} = \frac{58}{2} = 29$


10. (a): Let $I = \int \frac{\cos 2x - \cos 2\theta}{\cos x - \cos \theta} dx$

$= \int \frac{(2\cos^2 x - 1) - (2\cos^2 \theta - 1)}{\cos x - \cos \theta} dx$

$= \int \frac{2(\cos^2 x - \cos^2 \theta)}{\cos x - \cos \theta} dx = 2 \int (\cos x + \cos \theta) dx$

$= 2(\sin x + x \cos \theta) + C$

11. (a): Required area = Area of shaded portion



$$= 2 \int_0^{16} \sqrt{y} dy = 2 \cdot \frac{2}{3} [y^{3/2}]_0^{16} = \frac{4}{3} [4^3] = \frac{256}{3} \text{ sq. units}$$

12. (a): Given, $A \subset B \Rightarrow A \cap B = A$

$$\therefore n(A \cup B) = n(A) + n(B) - n(A \cap B) \\ = n(A) + n(B) - n(A) = n(B)$$

13. (b): Given, A is symmetric $\Rightarrow a_{ij} = a_{ji} \dots$ (i) $i \neq j$
 A is skew symmetric $\Rightarrow a_{ij} = -a_{ji} \dots$ (ii) and $a_{ii} = 0$
 Adding (i) and (ii) we get $2a_{ij} = 0 \Rightarrow a_{ij} = 0$

$\therefore A$ is a zero matrix.

14. (c): We have, $\begin{vmatrix} 3 & x \\ x & 1 \end{vmatrix} = \begin{vmatrix} 3 & 2 \\ 4 & 1 \end{vmatrix}$

$$\Rightarrow 3 - x^2 = 3 - 8 \Rightarrow x^2 = 8 \Rightarrow x = \pm 2\sqrt{2}$$

15. (d): Given, $x \cdot \frac{dy}{dx} + 2y = x^2 \Rightarrow \frac{dy}{dx} + \frac{2y}{x} = x$

$$\therefore \text{I.F.} = e^{\int \frac{2}{x} dx} = e^{\log x^2} = x^2$$

16. (c): Perpendicular distance of the point $P(6,7,8)$ from XY -plane is 8.

17. (c): Clearly, shaded region represents

$$5x + 4y \geq 20, x \leq 6, y \leq 3, x \geq 0, y \geq 0$$

18. (b): The optimal solution occurs at every point on the line joining these two points.

19. (c): Given series is in A.P. with first term $(a) = 3$, common difference $(d) = 2$

$$\therefore S_n = \frac{n}{2} [2 \times 3 + (n-1)2] = n(n+2)$$

20. (a): Given $\begin{bmatrix} 2+y & 6 \\ 1 & 2x+2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$

$$\Rightarrow 2+y=5 \text{ and } 2x+2=8 \Rightarrow y=3, x=3$$

21. (a): Let $u = \cos^{-1}(2x^2 - 1)$ and $v = \cos^{-1} x \Rightarrow \cos v = x$

$$\therefore u = \cos^{-1}(2\cos^2 v - 1) = \cos^{-1}(\cos 2v) = 2v$$

$$\therefore \frac{du}{dv} = 2$$

22. (c): $p = \frac{10}{100} = \frac{1}{10}, q = 1 - p = \frac{9}{10}, n = 5$

$$P(X = x) = {}^5C_x \left(\frac{1}{10}\right)^x \left(\frac{9}{10}\right)^{5-x}$$

$$\therefore P(X \leq 1) = P(X = 0) + P(X = 1)$$

$$= {}^5C_0 \left(\frac{1}{10}\right)^0 \left(\frac{9}{10}\right)^5 + {}^5C_1 \left(\frac{1}{10}\right)^1 \left(\frac{9}{10}\right)^4$$

23. (b): We have, $y = \log(\log x) \Rightarrow \frac{dy}{dx} = \frac{1}{x \log x}$

$$\Rightarrow \frac{d^2 y}{dx^2} = \frac{-1}{(x \log x)^2} \left(x \cdot \frac{1}{x} + \log x \cdot 1 \right) = \frac{-(1 + \log x)}{(x \log x)^2}$$

24. (a): Let $I = \int \frac{(x+3)e^x}{(x+4)^2} dx = \int \frac{(x+4-1)e^x}{(x+4)^2} dx$

$$= \int \left[\frac{1}{x+4} - \frac{1}{(x+4)^2} \right] e^x dx = \frac{e^x}{x+4} + C$$

$$\left[\because \int e^x (f(x) + f'(x)) dx = e^x f(x) + C \right]$$

25. (c): Let $I = \int_0^{\pi/2} \frac{1}{a^2 \sin^2 x + b^2 \cos^2 x} dx$

$$= \int_0^{\pi/2} \frac{\sec^2 x dx}{a^2 \tan^2 x + b^2}$$

$$\text{Put } \tan x = t \Rightarrow \sec^2 x dx = dt$$

$$\therefore I = \int_0^{\infty} \frac{dt}{a^2 t^2 + b^2} = \int_0^{\infty} \frac{dt}{a^2 \left(t^2 + \frac{b^2}{a^2} \right)}$$

$$= \frac{1}{a^2} \left[\frac{a}{b} \tan^{-1} \frac{at}{b} \right]_0^{\infty} = \frac{\pi}{2ab}$$

26. (b): Given $f(x) = x^4$

Now, $f(1) = f(-1)$ but $1 \neq -1$

$\therefore f$ is not one-one

Also, co-domain of f is R and range of f is $[0, \infty)$

$\therefore f$ is not onto

27. (b): We have, $y^2 = x \Rightarrow 2yy' = 1$

$$\Rightarrow y' = \frac{1}{2y} = \tan \frac{\pi}{4} \Rightarrow y = \frac{1}{2}$$

$$\text{When } y = \frac{1}{2}, x = y^2 = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$$

28. (a): Number of terms in $(x+a)^{47} - (x-a)^{47}$

$$= \frac{47+1}{2} = 24$$

[Number of terms in $(a+b)^n - (a-b)^n$ is $\frac{n+1}{2}$, when n is odd number]

29. (d): f is strictly increasing $\Rightarrow f'(x) > 0$
 $\Rightarrow 2x + 2 > 0 \Rightarrow x > -1$

30. (a): Highest order derivative is $\frac{d^2y}{dx^2}$ and its power is 1.

Hence, degree of differential equation is 1.

31. (b): We have, $a * b = \frac{a}{b+1}$

$$1 * 2 = \frac{1}{3} \text{ but } 2 * 1 = \frac{2}{2} = 1$$

Thus $1 * 2 \neq 2 * 1 \therefore *$ is not commutative

$$\text{Now, } (1 * 2) * 3 = \frac{1}{3} * 3 = \frac{1/3}{4} = \frac{1}{12}$$

$$\text{and } 1 * (2 * 3) = 1 * \frac{1}{2} = \frac{1}{\frac{1}{2} + 1} = \frac{2}{3}$$

$\therefore *$ is not associative

[Infact, $*$ is not a binary operation !!!]

32. (b): Let ϕ be the angle made by plane
 $2x - 3y + 6z - 11 = 0$ and X -axis i.e., $(1, 0, 0)$

$$\therefore \sin \phi = \frac{|2 \times 1 - 3 \times 0 + 6 \times 0|}{\sqrt{4 + 9 + 36} \sqrt{1 + 0 + 0}}$$

$$\Rightarrow \phi = \sin^{-1} \left(\frac{2}{7} \right) = \sin^{-1}(\alpha)$$

33. (d): We have C.V. = $\frac{\sigma}{\bar{x}} \times 100$

$$\Rightarrow 60 = \frac{24}{\bar{x}} \times 100 \Rightarrow \bar{x} = 40$$

34. (d): The contrapositive statement of the statement
 "If x is prime number, then x is odd" is "If x is not odd, then x is not a prime number."

35. (d): We know, $\sum P(X) = 1$

$$\Rightarrow 0.3 + k + 2k + 2k = 1 \Rightarrow 5k = 0.7 \Rightarrow 0.14$$

36. (b): Let $I = \int \sqrt{x^2 + 2x + 5} dx = \int \sqrt{(x+1)^2 + 2^2} dx$

$$= \frac{(x+1)}{2} \sqrt{x^2 + 2x + 5} + 2 \log \left| x+1 + \sqrt{x^2 + 2x + 5} \right| + C$$

37. (d): Given, ${}^nC_{12} = {}^nC_8 \Rightarrow 8 + 12 = n \Rightarrow n = 20$

$$[\because {}^nC_x = {}^nC_y \Rightarrow x = y \text{ or } x + y = n]$$

38. (a, c, d)

39. (a): Given, $\tan^{-1} x + \tan^{-1} y = \frac{4\pi}{5}$

$$\Rightarrow \cot^{-1} x + \cot^{-1} y = \pi - \frac{4\pi}{5} = \frac{\pi}{5}$$

40. (a): Let $y = \sqrt{9 - x^2} \Rightarrow y^2 = 9 - x^2$

$$\Rightarrow x^2 = 9 - y^2 \Rightarrow x = \sqrt{9 - y^2}$$

$$\text{Clearly, } 9 - y^2 \geq 0 \Rightarrow y^2 \leq 9$$

$$\Rightarrow -3 \leq y \leq 3$$

But $y \geq 0$. Hence $0 \leq y \leq 3$.

41. (a): If A and B are independent

Then A' and B' are also independent

$$\Rightarrow P(A' \cap B') = P(A')P(B') \\ = (1 - P(A))(1 - P(B))$$

$$42. (a): e = \frac{\sqrt{a^2 - b^2}}{a} = \frac{\sqrt{36 - 16}}{6} = \frac{2\sqrt{5}}{6}$$

43. (b): Given, $|\vec{a}| = |\vec{b}| = 1$

$$\text{Now, } |\sqrt{3}\vec{a} - \vec{b}|^2 = 3|\vec{a}|^2 + |\vec{b}|^2 - 2\sqrt{3}(\vec{a} \cdot \vec{b})$$

$$\Rightarrow 1 = 3(1) + 1 - 2\sqrt{3}|\vec{a}||\vec{b}|\cos\theta$$

$$\Rightarrow 1 = 4 - 2\sqrt{3}(1)(1)\cos\theta$$

$$\Rightarrow \cos\theta = \frac{\sqrt{3}}{2} \Rightarrow \theta = 30^\circ$$

44. (c): For, \vec{a} and \vec{b} to be orthogonal $\vec{a} \cdot \vec{b} = 0$

$$\Rightarrow (2)(1) + (\lambda)(2) + (1)(3) = 0$$

$$\Rightarrow 5 + 2\lambda = 0 \Rightarrow \lambda = -\frac{5}{2}$$

45. (b): $\cos^2 45^\circ - \sin^2 15^\circ = \cos(45^\circ + 15^\circ)\cos(45^\circ - 15^\circ)$

$$= \cos 60^\circ \cos 30^\circ = \frac{1}{2} \cdot \frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{4}$$

46. (b): Range of $\sec^{-1} x$ is $[0, \pi] - \left\{ \frac{\pi}{2} \right\}$

47. (a): Here, $\vec{a}, \vec{b}, \vec{c}$ are unit vectors and $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

$$\therefore (\vec{a} + \vec{b} + \vec{c})^2 = 0$$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow 1 + 1 + 1 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = -\frac{3}{2}$$

48. (c): Let $I = \int_{-\pi/2}^{\pi/2} \frac{dx}{e^{\sin x} + 1}$... (i)

$$\Rightarrow I = \int_{-\pi/2}^{\pi/2} \frac{dx}{e^{-\sin x} + 1} = \int_{-\pi/2}^{\pi/2} \frac{e^{\sin x} dx}{1 + e^{\sin x}} \quad \dots (ii)$$

Adding (i) and (ii), we get

$$2I = \int_{-\pi/2}^{\pi/2} \frac{e^{\sin x} + 1}{e^{\sin x} + 1} dx$$

$$\Rightarrow 2I = \int_{-\pi/2}^{\pi/2} 1 dx = [x]_{-\pi/2}^{\pi/2} = \pi \Rightarrow I = \frac{\pi}{2}$$

49. (a): $\frac{dV}{dS} = \frac{dV/dt}{dS/dt} = \frac{4\pi r^2 \frac{dr}{dt}}{8\pi r \frac{dr}{dt}} = \frac{r}{2} = \frac{4}{2} = 2 \text{ cm}^3/\text{cm}^2$

50. (a): $\int_0^{\pi/2} \frac{\tan^7 x}{\cot^7 x + \tan^7 x} dx = \frac{\pi}{4}$

$$\left[\because \int_0^{\pi/2} \frac{f(x)}{f(x) + f\left(\frac{\pi}{2} - x\right)} dx = \frac{\pi}{4} \right]$$

51. (c): Here, $|x - 2| \leq 1 \Rightarrow -1 \leq x - 2 \leq 1 \Rightarrow 1 \leq x \leq 3$
 $\Rightarrow x \in [1, 3]$

52. (b): Here, $\int_{0.2}^{3.5} [x] dx = \int_{0.2}^1 0 dx + \int_1^2 1 dx + \int_2^3 2 dx + \int_3^{3.5} 3 dx$
 $= 0 + [x]_1^2 + [2x]_2^3 + [3x]_3^{3.5} = 1 + 2(1) + 3(0.5) = 4.5$

53. (d): Given, area of triangle with vertices $(K, 0)$, $(4, 0)$, $(0, 2)$ is 4 sq. units i.e.,

$$\pm 4 = \frac{1}{2} \begin{vmatrix} K & 0 & 1 \\ 4 & 0 & 1 \\ 0 & 2 & 1 \end{vmatrix} \Rightarrow \frac{-2K + 8}{2} = \pm 4$$

$$\Rightarrow K = 0 \text{ or } 8$$

54. (b): Given, $f(x)$ is continuous

$$\Rightarrow \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = f(2)$$

$$\Rightarrow \lim_{x \rightarrow 2} (Kx^2) = 3 \Rightarrow 4K = 3 \Rightarrow K = \frac{3}{4}$$

55. (None of the options is correct):

$$\text{If matrix } B \text{ will be, } B = \frac{1}{\pi} \begin{bmatrix} -\cos^{-1} \pi x & \tan^{-1} \frac{x}{\pi} \\ \sin^{-1} \left(\frac{x}{\pi} \right) & -\tan^{-1} \pi x \end{bmatrix}$$

$$\text{Then, } A - B = \frac{1}{2} I$$

56. (c): $f \circ g(x) = f(g(x)) = f(x^{1/3}) = 8(x^{1/3})^3 = 8x$

57. (b): We have, $\Delta = \begin{vmatrix} Ax & x^2 & 1 \\ By & y^2 & 1 \\ Cz & z^2 & 1 \end{vmatrix} = xyz \begin{vmatrix} A & x & \frac{1}{x} \\ B & y & \frac{1}{y} \\ C & z & \frac{1}{z} \end{vmatrix}$

$$= \begin{vmatrix} A & x & yz \\ B & y & xz \\ C & z & xy \end{vmatrix} = \begin{vmatrix} A & B & C \\ x & y & z \\ zy & zx & xy \end{vmatrix} = \Delta_1$$


58. (a): We have, $x = \sin^{-1} \frac{2t}{1+t^2} = 2 \tan^{-1} t$

$$\text{and } y = \tan^{-1} \frac{2t}{1-t^2} = 2 \tan^{-1} t$$

$$\therefore y = x \Rightarrow \frac{dy}{dx} = 1.$$

59. (b): Reflection of the point (α, β, γ) in XY plane is $(\alpha, \beta, -\gamma)$.

60. (a): Given, curves are $y = \cos x$; $x = 0$ and $x = \pi$



$$\therefore \text{Required area} = 2 \int_0^{\pi/2} \cos x dx$$

$$= 2 [\sin x]_0^{\pi/2} = 2 \text{ sq. units}$$



Solution Sender of Maths Musing

SET-173

1. V. Damodhar Reddy (Telangana)

SET-172

1. Satyadev. P (Bangalore)
2. Gouri Sankar Adhikary (West Bengal)
3. P.N.L Bhavana (Hyderabad)
4. Shubham Agarwal (Hyderabad)
5. K. Poojasree (Hyderabad)
6. N. Jayanthi (Hyderabad)

Minimise your efforts to
excel in **Boards & Competitive Exams**




✓ Error-free ✓ Easy Language ✓ Authentic Solutions ✓ Fully Solved by Experts

NCERT

EXERCISES+EXEMPLAR SOLUTIONS

A number of questions in School Examinations/Boards, Medical, Engineering Entrance Exams and Olympiads are directly picked from NCERT books. Going by Experts' and Toppers' advice, a thorough revision of NCERT textbooks will definitely widen your chances of selection in any exam.

Available for Class 6 to 12



You may find free NCERT Solutions at many places
but why MTG is better-than-the-rest

- Comprehensive Guide that covers entire solutions of NCERT Textbook Questions and NCERT Exemplar Books questions
- Expert answers by experienced teachers who check CBSE Papers
- Error free, Authentic solutions with the accurate information
- Complete solution for all the questions



Available at all leading book shops throughout the country.
For more information or for help in placing your order:
Call 0124-6601200 or email: info@mtg.in

MATHS MUSING

Maths Musing was started in January 2003 issue of Mathematics Today. The aim of Maths Musing is to augment the chances of bright students seeking admission into IITs with additional study material.

During the last 10 years there have been several changes in JEE pattern. To suit these changes Maths Musing also adopted the new pattern by changing the style of problems. Some of the Maths Musing problems have been adapted in JEE benefitting thousand of our readers. It is heartening that we receive solutions of Maths Musing problems from all over India.

Maths Musing has been receiving tremendous response from candidates preparing for JEE and teachers coaching them. We do hope that students will continue to use Maths Musing to boost up their ranks in JEE Main and Advanced.

PROBLEM Set 174

JEE MAIN

- If $a_1 = 11, a_2 = 75, a_3 = 20, a_4 = 23$ and $a_n = a_{n-1} - a_{n-2} + a_{n-3} - a_{n-4}, n \geq 5$, then $a_5 - a_3 + a_2 =$
(a) 53 (b) 58
(c) 65 (d) 75
- If $xyz = 1, x + \frac{1}{z} = 5, y + \frac{1}{x} = 29$, then $z + \frac{1}{y} =$
(a) $\frac{1}{3}$ (b) $\frac{1}{4}$
(c) 2 (d) $\frac{3}{4}$
- In a triangle $ABC, \Delta \left(\frac{1}{r^2} + \frac{1}{r_1^2} + \frac{1}{r_2^2} + \frac{1}{r_3^2} \right) =$
(a) $2 \sum \tan A$ (b) $2 \sum \cot A$
(c) $4 \sum \tan A$ (d) $4 \sum \cot A$
- A straight line through the point (a, b) meets x -axis at A and y -axis at B . O is the origin. If $(a, b) = (4, 1)$, then the minimum value of $OA + OB$ is
(a) 7 (b) 8 (c) 9 (d) 10

- If $x \neq a, y \neq b, z \neq c$ and $\begin{vmatrix} a & y & z \\ x & b & z \\ x & y & c \end{vmatrix} = 0$, then

$$\frac{x+a}{x-a} + \frac{y+b}{y-b} + \frac{z+c}{z-c} =$$

- (a) -1 (b) 0
(c) 1 (d) 2

JEE ADVANCED

- A bag contains 30 tokens numbered serially from 0 to 30. The number of ways of selecting 3 tokens from the bag, so that the sum of the numbers on them is 30, is divisible by
(a) 2 (b) 3 (c) 5 (d) 7

COMPREHENSION

- Let $f(x) = [\sin^{-1} x] + [\cos^{-1} x], x \in [0, 1]$
- The number of points at which $f(x)$ is not differentiable is
(a) 0 (b) 1 (c) 2 (d) 3
 - The solution set of the equation $f(x) = 0$ is an interval of length
(a) $\sqrt{2} \sin \left(1 - \frac{\pi}{4} \right)$ (b) $\sqrt{2} \sin \left(1 + \frac{\pi}{4} \right)$
(c) $\sqrt{2} \cos \left(1 - \frac{\pi}{4} \right)$ (d) $\sqrt{2} \cos \left(1 + \frac{\pi}{4} \right)$

INTEGER TYPE

- Ten boys and two girls are to be seated in a row such that there are atleast 3 boys between the girls. The number of ways this can be done is $\lambda \cdot 12!$, where $\lambda =$

MATRIX MATCH

- The curve $f(x, y) = 0$ lies in the first quadrant. The tangent at a point on it meets the positive x and y axes at A and B and O is the origin

	List-I		List-II
P.	$f(x, y) = 4xy - 1$	1.	$AB = 1$
Q.	$f(x, y) = x^2 + y^2 - 1$	2.	$OA + OB = 1$
R.	$f(x, y) = \sqrt{x} + \sqrt{y} - 1$	3.	$OA \cdot OB = 1$
S.	$f(x, y) = x^{2/3} + y^{2/3} - 1$	4.	$\frac{1}{OA^2} + \frac{1}{OB^2} = 1$

P	Q	R	S
(a) 2	1	3	4
(b) 1	2	4	3
(c) 4	3	2	1
(d) 3	4	2	1

See Solution Set of Maths Musing 173 on page no. 85

20

CHALLENGING PROBLEMS

For Entrance Exams

- Let x, y, z be real numbers such that $\cos x + \cos y + \cos z = 0$ and $\cos 3x + \cos 3y + \cos 3z = 0$, then $\cos 2x \cos 2y \cos 2z$
 - ≤ 0
 - ≥ 0
 - depends on x, y, z values
 - data insufficient
- Eliminate θ from the system: $\lambda \cos 2\theta = \cos(\theta + \alpha)$ and $\lambda \sin 2\theta = 2 \sin(\theta + \alpha)$
 - $(\cos \alpha)^{2/3} - (\sin \alpha)^{2/3} = \lambda^{2/3}$
 - $(\cos \alpha)^{2/3} + (\sin \alpha)^{2/3} = \lambda^{2/3}$
 - $(\cos \alpha)^{1/3} - (\sin \alpha)^{1/3} = \lambda^{1/3}$
 - $(\cos \alpha)^{1/3} + (\sin \alpha)^{1/3} = \lambda^{1/3}$
- The bisector of $\angle BAC$ intersects the circumcircle of $\triangle ABC$ at D . If $AB^2 + AC^2 = 2AD^2$ then angle of intersection of AD and BC is
 - 30°
 - 45°
 - 60°
 - 90°
- The laws of points (x, y) satisfying the equations $x^2 + y \cos^2 \alpha = x \sin \alpha \cos \alpha$ and $x \cos 2\alpha + y \sin 2\alpha = 0$ lies on (α is a parameter)
 - circle
 - parabola
 - ellipse
 - hyperbola
- In $\triangle ABC$, $\angle ABC = \angle ACB = 40^\circ$. If P is a point in the interior of the triangle such that $\angle PBC = 20^\circ$ and $\angle PCB = 30^\circ$ then
 - $BP = BA$
 - $BP = 2BA$
 - $BP = \frac{1}{2} BA$
 - $BP = 3BA$
- Let ABC be a triangle of area $\frac{1}{2}$, then minimum value of $a^2 + \csc A$ is
 - $\sqrt{2}$
 - $\sqrt{3}$
 - $\sqrt{4}$
 - $\sqrt{5}$
- Let $\alpha, \beta, \gamma, \delta$ be positive numbers such that for all x , $\sin \alpha x + \sin \beta x = \sin \gamma x + \sin \delta x$, then $\gamma + \delta =$
 - α
 - 2α
 - 3α
 - 4α
- Let S be the set of all triangles ABC for which $5 \left[\frac{1}{AP} + \frac{1}{BQ} + \frac{1}{CR} \right] - \frac{3}{\min\{AP, BQ, CR\}} = \frac{6}{r}$, where r is inradius and P, Q, R are points of tangency of incircle with sides AB, BC, CA respectively then all the triangles in the set S are
 - scalene
 - isosceles
 - equilateral
 - right angled
- Let ABC be a triangle such that $\max\{A, B\} = C + 30^\circ$ and $\frac{R}{r} = \sqrt{3} + 1$, R is circumradius, r is inradius then $\triangle ABC$ is
 - scalene
 - isosceles
 - equilateral
 - right angled
- Let $ABCDEFGHIJKL$ be a regular do-decagon then $\frac{AB}{AF} + \frac{AF}{AB} =$
 - 1
 - 2
 - 3
 - 4
- Let $a, b, c, d \in [0, \pi]$ such that $2 \cos a + 6 \cos b + 7 \cos c + 9 \cos d = 0$ and $2 \sin a - 6 \sin b + 7 \sin c - 9 \sin d = 0$ then $\frac{\cos(a+d)}{\cos(b+c)} =$
 - $\frac{7}{3}$
 - $\frac{3}{7}$
 - $\frac{3}{5}$
 - $\frac{5}{3}$
- If $\sin x \cos y + \sin y \cos z + \sin z \cos x = \frac{3}{2}$, then
 - $\sin x = \cos 2y$
 - $\sin x = \sin y$
 - $\sin x = \cos y$
 - $\sin x = \cos z$

13. In $\triangle ABC$, $\lambda = \sin A \sin B + \sin B \sin C + \sin C \sin A$ and $(1 + \sin A)(1 + \sin B)(1 + \sin C) = 2(\lambda + 1)$, then $\triangle ABC$ is

(a) scalene (b) isosceles
(c) equilateral (d) right angled

14. Let ABC be a triangle such that $\sin^2 B + \sin^2 C = 1 + 2 \sin B \sin C \cos A$, then $\triangle ABC$ is
- (a) scalene (b) isosceles
(c) equilateral (d) right angled

15. In $\triangle ABC$, $\left(\cot \frac{A}{2}\right)^2 + 4 \cot^2 \frac{B}{2} + 9 \cot^2 \frac{C}{2} = \left(\frac{6s}{7r}\right)^2$ where $s \rightarrow$ semiperimeter and $r \rightarrow$ inradius, then the ratio of the sides of the triangle is
- (a) 45 : 40 : 13 (b) 50 : 55 : 45
(c) 55 : 65 : 70 (d) 60 : 65 : 80

16. Let n be a positive integer and for real numbers λ and a_{kl} ($k, l = 1, 2, 3, \dots, n$) ($k > l$), we have

$$\frac{\sin^2 nx}{\sin^2 x} = \lambda + \sum_{1 \leq l < k \leq n} a_{kl} \cos(2k - 2l)x$$

($x \neq m\pi, m \in I$) then $\lambda =$

(a) l (b) k (c) n (d) $k - l$

17. In $\triangle ABC$, $\angle A = 30^\circ$ and an inscribed circle of fixed radius r is drawn. If $\triangle ABC$ has least perimeter then $\angle B =$

(a) 30° (b) 45° (c) 60° (d) 75°

18. If A_1, A_2, A_3, A_4 are the angles of a convex quad. then max. value of

$$\sin\left(\frac{A_1}{2}\right) + \sin\left(\frac{A_2}{2}\right) + \sin\left(\frac{A_3}{2}\right) + \sin\left(\frac{A_4}{2}\right) \text{ is}$$

(a) $\sqrt{2}$ (b) $2\sqrt{2}$ (c) $3\sqrt{2}$ (d) $4\sqrt{2}$

19. If a, b, c, k are constants and α, β, γ are variables subject to the relation $a \tan \alpha + b \tan \beta + c \tan \gamma = k$ then minimum value of $\tan^2 \alpha + \tan^2 \beta + \tan^2 \gamma$ is

(a) $\frac{k^2}{(a+b+c)^2}$ (b) $\frac{k^2}{a^2+b^2+c^2}$

(c) $\frac{(a+b+c)^2}{k^2}$ (d) $\frac{a^2+b^2+c^2}{k^2}$

20. For $x \in R$, the minimum value of $|\sin x + \cos x + \tan x + \sec x + \operatorname{cosec} x + \cot x|$ is

(a) $2\sqrt{2}$ (b) $2\sqrt{2} - 1$

(c) $2\sqrt{2} + 1$ (d) $\sqrt{2} - 1$

SOLUTIONS

1. (a) : Using $\cos 3x = 4 \cos^3 x - 3 \cos x$.

We have, $\sum \cos^3 x = 0$

using identity $\sum a^3 - 3abc = (\sum a)(\sum a^2 - \sum ab)$

We have, $abc = \cos x \cos y \cos z = 0$

Let $\cos z = 0$. So, $\cos x = -\cos y$ and

$\cos 2x = \cos 2y$ and $\cos 2z = -1$

So, $\cos 2x \cos 2y \cos 2z = -\cos^2 2x \leq 0$.

2. (b) : Expanding the given equations, we have $(\cos \theta \cos \alpha - \sin \theta \sin \alpha) = \lambda(\cos^2 \theta - \sin^2 \theta)$ and $\sin \theta \cos \alpha + \cos \theta \sin \alpha = \lambda \sin \theta \cos \theta$
 $\Rightarrow \cos \alpha = \lambda \cos^3 \theta$ and $\sin \alpha = \lambda \sin^3 \theta$
Hence, $(\cos \alpha)^{2/3} + (\sin \alpha)^{2/3} = \lambda^{2/3}$.

3. (b) : Let $\angle CAB = \alpha$, $\angle ABC = \beta$, $\angle BCA = \gamma$ then using sine rule, we have

$$\frac{AB}{\sin \gamma} = \frac{AC}{\sin \beta} = \frac{AD}{\sin(\alpha/2 + \beta)}$$

So, $AB^2 + AC^2 = 2AD^2$ becomes

$$\sin^2 \gamma + \sin^2 \beta = 2 \sin^2(\alpha/2 + \beta)$$

Simplifying, $\cos(\beta - \gamma)[1 + \cos(\beta + \gamma)] = 0$

i.e., $\cos(\beta - \gamma) = 0$. So, $\alpha + 2\beta = \pi/2$

$$\text{So, } \angle AEC = \frac{\alpha}{2} + \beta = \frac{\pi}{4} = 45^\circ$$

4. (b) : Simplifying the first equation, we have

$$x \sin 2\alpha - y \cos 2\alpha = 2x^2 + y$$

and the other given equation is

$$x \cos 2\alpha + y \sin 2\alpha = 0$$

Solving, we have

$$\sin 2\alpha = \frac{x(2x^2 + y)}{x^2 + y^2} \text{ and } \cos 2\alpha = \frac{-y(2x^2 + y)}{x^2 + y^2}$$

Hence, $\sin^2 2\alpha + \cos^2 2\alpha = 1$ gives

$$\frac{(2x^2 + y)^2}{x^2 + y^2} = 1 \text{ i.e., } 4x^2 + 4y - 1 = 0, \text{ Parabola.}$$

5. (a) : Let us assume that $AB = AC = 1$ unit then

$$BC = 2 \cos 40^\circ \text{ and } \angle BPC = 130^\circ$$

Applying sine rule in $\triangle BPC$, we have

$$\frac{BP}{\sin(30^\circ)} = \frac{BC}{\sin(130^\circ)}$$

$$\Rightarrow BP = \frac{BC \cdot \sin 30^\circ}{\cos 40^\circ} = \frac{2 \cos 40^\circ \cdot \sin 30^\circ}{\cos 40^\circ} = 1$$

i.e., $BP = AB$.

6. (d) : Given, area = $\frac{1}{2} \Rightarrow \frac{1}{2} bc \sin A = \frac{1}{2}$

$\Rightarrow \operatorname{cosec} A = bc$ and $[bc \geq 1]$

Now,

$a^2 + \operatorname{cosec} A = a^2 + bc = b^2 + c^2 - 2bc \cos A + bc$
[cosine rule]

$= b^2 + c^2 - 2bc\sqrt{1 - \sin^2 A} + bc$

$= b^2 + c^2 - 2\sqrt{b^2c^2 - (bc \sin A)^2} + bc$

$= b^2 + bc + c^2 - 2\sqrt{b^2c^2 - 1}$

$\geq 3bc - 2\sqrt{b^2c^2 - 1}$

Let $x = bc (\geq 1)$ then $y = 3x - 2\sqrt{x^2 - 1}$ gives

$5x^2 - 6xy + y^2 + 4 = 0$. As $x \in R, D \geq 0$ gives

$(-6y)^2 - 20(y^2 + 4) \geq 0$ i.e., $y \geq \sqrt{5}$.

7. (b) : Differentiating the given identity three times, we have, $\alpha^3 \cos \alpha x + \beta^3 \cos \beta x = \gamma^3 \cos \gamma x + \delta^3 \cos \delta x$
Also, $\alpha \cos \alpha x + \beta \cos \beta x = \gamma \cos \gamma x + \delta \cos \delta x$

In particular for $x = 0$, we have

$\alpha + \beta = \gamma + \delta$ and $\alpha^3 + \beta^3 = \gamma^3 + \delta^3$
i.e., $(\alpha + \beta)^3 = (\gamma + \delta)^3$

$\Rightarrow \alpha\beta = \gamma\delta$ on simplification.

So, $(\alpha - \gamma)(\alpha - \delta) = \alpha^2 - \alpha(\gamma + \delta) + \gamma\delta$
 $= \alpha^2 - \alpha(\alpha + \beta) + \alpha\beta = 0$

$\Rightarrow \gamma = \alpha$ and $\delta = \alpha$ i.e., $\gamma + \delta = 2\alpha$

8. (b) : Let us assume that $\min. \{AP, BQ, CR\} = AP$ and let $\tan(A/2) = x, \tan(B/2) = y, \tan(C/2) = z$.

So, $AP = \frac{r}{x}, BQ = \frac{r}{y}$ and $CR = \frac{r}{z}$

Now, the relation in question becomes

$2x + 5y + 5z = 6$

and in any Δ , we know that $xy + yz + zx = 1$.

Now, eliminating (x) from these two equations,

we have $5y^2 + 5z^2 + 8yz - 6y - 6z + 2 = 0$

i.e., $(3y - 1)^2 + (3z - 1)^2 = 4(y - z)^2$

or, $5\alpha^2 + 5\beta^2 + 8\alpha\beta = 0$ [where $3y - 1 = \alpha,$
 $3z - 1 = \beta$]

$\Rightarrow \alpha = 0 = \beta$ for real solutions.

i.e., $y = z = \frac{1}{3}$ and so, $x = \frac{4}{3}$

i.e., isosceles triangle.

9. (d) : Let $\max. \{A, B\} = A$ then A.T.Q, $A - C = 30^\circ$ and using the identity

$r = 4R \sin\left(\frac{A}{2}\right) \cdot \sin\left(\frac{B}{2}\right) \cdot \sin\left(\frac{C}{2}\right)$

We have, $r = 4R\sqrt{3+1} \left[\sin\left(\frac{A}{2}\right) \cdot \sin\left(\frac{B}{2}\right) \cdot \sin\left(\frac{C}{2}\right) \right]$

i.e., $\frac{\sqrt{3}-1}{4} = \sin \frac{B}{2} \left[\cos \frac{A-C}{2} - \cos \frac{A+C}{2} \right]$

i.e., $\frac{\sqrt{3}-1}{4} = \sin \frac{B}{2} \left[\cos\left(\frac{30^\circ}{2}\right) - \cos\left(\frac{180^\circ-B}{2}\right) \right]$

i.e., $\sin^2 \frac{B}{2} - \sin\left(\frac{B}{2}\right) \cdot \left(\frac{\sqrt{6}+\sqrt{2}}{4}\right) + \left(\frac{\sqrt{3}-1}{4}\right) = 0$

Solving, $\sin(B/2) = \frac{\sqrt{6}-\sqrt{2}}{4}$ or $\frac{\sqrt{2}}{2}$

i.e., $\frac{B}{2} = 15^\circ$ or 45° .

But $B = 90^\circ$ is not possible. Hence, $B = 30^\circ$ and $A = 90^\circ, C = 60^\circ$, i.e., [Right angled Δ]

10. (d) : Let R be the circumradius then

$AB = 2R \sin\left(\frac{\pi}{12}\right)$ and $AF = 2R \sin\left(\frac{5\pi}{12}\right)$

So, the required quantity is $\frac{2R \sin \theta}{2R \sin 5\theta} + \frac{2R \sin 5\theta}{2R \sin \theta}$

$\left[\text{where } \theta = \frac{\pi}{12} \right]$

$= \frac{\sin^2 \theta + \sin^2 5\theta}{\sin 5\theta \sin \theta} = \frac{1 - \cos 2\theta + 1 - \cos 10\theta}{\cos 4\theta - \cos 6\theta}$
 $= 4$ (on simplification)

11. (a) : Rearranging the two equations, we have

$2\sin a - 9\sin d = 6\sin b - 7\sin c$

and $2\cos a + 9\cos d = -6\cos b - 7\cos c$

Squaring and adding both the equations, we have

$85 + 36\cos(a+d) = 85 + 84\cos(b+c)$

i.e., $\frac{\cos(a+d)}{\cos(b+c)} = \frac{84}{36} = \frac{7}{3}$

12. (c): The given equation can be rewritten as,

$(\sin x - \cos y)^2 + (\sin y - \cos z)^2 + (\sin z - \cos x)^2 = 0$

i.e., $\sin x = \cos y$

13. (d) : Equating the two λ values and simplifying, we have

$(1 - \sin A)(1 - \sin B)(1 - \sin C) = 0$

i.e., $\sin A = 1$ or $\sin B = 1$ or $\sin C = 1$

i.e., ABC is right angled triangle.

14. (d) : Using cosine rule, we have

$$a^2 + 2bc \cos A = b^2 + c^2$$

And using sine rule, here, we have

$$\sin^2 B + \sin^2 C = \sin^2 A + 2 \sin B \sin C \cos A$$

Now comparing this with equation given in question, we have, $\sin^2 A = 1$ i.e., $A = 90^\circ$

15. (a) : In any triangle, we have

$$\cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} = \cot \frac{A}{2} \cdot \cot \frac{B}{2} \cdot \cot \frac{C}{2} = \frac{s}{r}$$

So, the given relation in question becomes,

$$(6^2 + 3^2 + 2^2) \left[\left(\cot \frac{A}{2} \right)^2 + \left(2 \cot \frac{B}{2} \right)^2 + \left(3 \cot \frac{C}{2} \right)^2 \right] \\ = \left(6 \cot \frac{A}{2} + 6 \cot \frac{B}{2} + 6 \cot \frac{C}{2} \right)^2$$

i.e., Equality in Cauchy-Schwarz inequality.

$$\text{So, } \frac{\cot(A/2)}{6} = \frac{2 \cot(B/2)}{3} = \frac{3 \cot(C/2)}{2}$$

$$\text{i.e., } \cot(A/2) = 7, \cot(B/2) = \frac{7}{4} \text{ and } \cot(C/2) = \frac{7}{9}$$

$$\Rightarrow \sin A = \frac{7}{25}, \sin B = \frac{56}{65}, \sin C = \frac{126}{130}$$

i.e., Sides of the triangle are 26, 80 and 90.

16. (c): Using the two identities, we have

$$S_1 = \sum_{m=1}^n \cos(2mx) = \frac{\sin nx \cdot \cos(n+1)x}{\sin x}$$

$$\text{and } S_2 = \sum_{m=1}^n \sin(2mx) = \frac{\sin nx \cdot \sin(n+1)x}{\sin x}$$

$$S_1^2 + S_2^2 = \left(\frac{\sin nx}{\sin x} \right)^2$$

$$\text{But } S_1^2 + S_2^2 = (\cos 2x + \cos 4x + \dots + \cos nx)^2 \\ + (\sin 2x + \sin 4x + \dots + \sin nx)^2 \\ = n + 2 \sum_{1 \leq l < k \leq n} (\cos 2kx \cos 2lx + \sin 2kx \sin 2lx)$$

$$\text{i.e., } S_1^2 + S_2^2 = n + \sum_{1 \leq l < k \leq n} \cos 2(k-l)x$$

$$\text{i.e., } \lambda = n$$

17. (d) : In any triangle, we have $\sum \cot(A/2) = \frac{s}{r}$

So, perimeter $2s = 2r \cdot \sum \cot(A/2)$

Since A, r are fixed, s is min. when $\sum \cot(A/2)$ is min.

i.e., $\cot(B/2) + \cot(C/2)$ is min.

i.e., $\frac{\cos(A/2)}{\sin(B/2) \sin(C/2)}$ is min.

or, $\sin(B/2) \cdot \sin(C/2)$ is max.

i.e., $\frac{1}{2} \left[\cos\left(\frac{B-C}{2}\right) - \sin\left(\frac{A}{2}\right) \right]$ is max.

i.e., $\cos\left(\frac{B-C}{2}\right) = 1$ for max. i.e., $B = C$.

$\therefore \Delta ABC$ is an isosceles Δ with $\angle B = \angle C = 75^\circ$.

18. (b) : $A_1 + A_2 + A_3 + A_4 = 2\pi$,

$$\text{So, } \frac{A_1}{2} + \frac{A_2}{2} + \frac{A_3}{2} + \frac{A_4}{2} = \pi$$

Now, $f(x) = \sin x$ is a concave function.

So, $\sin\left(\frac{A_1}{2}\right) + \sin\left(\frac{A_2}{2}\right) + \sin\left(\frac{A_3}{2}\right) + \sin\left(\frac{A_4}{2}\right)$ is max.

$$\text{When } \frac{A_1}{2} = \frac{A_2}{2} = \frac{A_3}{2} = \frac{A_4}{2} = \text{each} = \frac{\pi}{4}$$

[Result of Jensen's inequality]

So, required max. value is $4 \cdot \sin(\pi/4) = 2\sqrt{2}$.

19. (b) : Use the identity

$$\sum (b \tan \gamma - c \tan \beta)^2 \\ = (a^2 + b^2 + c^2)(\tan^2 \alpha + \tan^2 \beta + \tan^2 \gamma) \\ - (a \tan \alpha + b \tan \beta + c \tan \gamma)^2$$

We have, $\text{RHS} \geq 0$

$$\Rightarrow (\sum a^2) \cdot (\sum \tan^2 \alpha) - k^2 \geq 0$$

$$\text{i.e., } \left(\sum \tan^2 \alpha \right)_{\min} = \frac{k^2}{\sum a^2}$$

20. (b) : Let $E = |\sin x + \cos x + \tan x + \sec x + \operatorname{cosec} x + \cot x|$

Putting $a = \sin x, b = \cos x, c = a + b$

We have on simplification, the given expression

$$E = \left| c + \frac{2}{c-1} \right| = \left| c-1 + \frac{2}{c-1} + 1 \right|$$

where $c \in [-\sqrt{2}, \sqrt{2}]$

Using $\text{AM} \geq \text{GM}$ on $(c-1)$ and $\left(\frac{2}{c-1}\right)$, we have

$$E_{\min} = |-2\sqrt{2} + 1| \quad \text{i.e., } 2\sqrt{2} - 1$$



MATH archives



Math Archives, as the title itself suggests, is a collection of various challenging problems related to the topics of IIT-JEE Syllabus. This section is basically aimed at providing an extra insight and knowledge to the candidates preparing for IIT-JEE. In every issue of MT, challenging problems are offered with detailed solution. The readers' comments and suggestions regarding the problems and solutions offered are always welcome.

1. The sum of all the non-real roots of

$$(x^2 + x - 2)(x^2 + x - 3) = 12 \text{ is}$$

- (a) 1 (b) -1
(c) 6 (d) -2

2. Statement-1 : The equation $\sin x + x \cos x = 0$ has at least one root in the interval $(0, \pi)$.

Statement-2 : Between any two roots of $f(x) = 0$, there exists atleast one root of $f'(x) = 0$.

- (a) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.
(b) Statement-1 is true, Statement-2 is true; Statement-2 is not correct explanation for Statement-1.
(c) Statement-1 is true, Statement-2 is false.
(d) Statement-1 is false, Statement-2 is true.

3. If LCM of p, q is $r^2 t^4 s^2$, where r, s, t are prime numbers and p, q are positive integers. Then the number of ordered pairs (p, q) is

- (a) 252 (b) 254
(c) 225 (d) 224

4. Let $f(x) = \max\{x, x^3\}$ $x \in R$ the set of points where $f(x)$ is not differentiable is

- (a) $\{-1, 1\}$ (b) $\{-1, 0\}$
(c) $\{0, 1\}$ (d) $\{-1, 0, 1\}$

5. Statement-1: The sum of the first 30 terms of the sequence 1, 2, 4, 7, 11, 16, 22, 29, 37, 46 ... is 4520.

Statement-2: The successive differences of the terms of the sequence form an A.P.

- (a) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.
(b) Statement-1 is true, Statement-2 is true; Statement-2 is not correct explanation for Statement-1.
(c) Statement-1 is true, Statement-2 is false.
(d) Statement-1 is false, Statement-2 is true.

6. $\sum_{r=1}^n \tan^{-1} \left(\frac{1}{2r^2} \right) =$

- (a) $\tan^{-1} n$ (b) $\tan^{-1} \frac{n}{n+1}$
(c) $\tan^{-1} \frac{n}{n+2}$ (d) $\tan^{-1} \left(\frac{n+1}{n+2} \right)$

7. The statement $p \rightarrow (q \rightarrow p)$ is equivalent to

- (a) $p \rightarrow (p \rightarrow q)$ (b) $p \rightarrow (p \vee q)$
(c) $p \rightarrow (p \wedge q)$ (d) $p \rightarrow (p \leftrightarrow q)$

8. The remainder left out when $8^{2n} - (62)^{2n+1}$ is divisible by 9 (where $n \in N$)

- (a) 0 (b) 2 (c) 7 (d) 8

9. Consider all functions that can be defined from the set $A = \{1, 2, 3\}$ to the set $B = \{1, 2, 3, 4, 5\}$. A function $f(x)$ is selected at random from these functions. The probability that, selected function satisfies $f(i) \leq f(j)$ for $i < j$, is equal to

- (a) $\frac{6}{25}$ (b) $\frac{12}{25}$
(c) $\frac{2}{5}$ (d) none of these

10. $\int_0^{\pi} [\cot x] dx =$

(where $[\cdot]$ denotes the greatest integer function)

- (a) $\frac{\pi}{2}$ (b) 1 (c) -1 (d) $-\frac{\pi}{2}$

SOLUTIONS

1. (b): Put $x^2 + x = y$ then, we have $y^2 - 5y - 6 = 0$
 $\Rightarrow (y - 6)(y + 1) = 0$

$$\Rightarrow x^2 + x - 6 = 0 \text{ or } x^2 + x + 1 = 0$$


$$\Rightarrow x = -3, 2 \text{ or } x = \omega, \omega^2$$

$$\text{Sum of non-real roots} = \omega + \omega^2 = -1$$

2. (c): Take $f(x) = x \sin x$, which is continuous in $[0, \pi]$ and differentiable in $(0, \pi)$. Also $f(0) = f(\pi) = 0$. By Rolle's theorem, there exists at least one root of $f'(x) = 0 \Rightarrow x \cos x + \sin x = 0$

3. (c): No. of ordered pairs
 $= (2(2) + 1)(2(4) + 1)(2(2) + 1)$
 $= 5 \times 9 \times 5 = 225$

4. (d):



5. (d): $a_2 - a_1 = 1, a_3 - a_2 = 2, a_4 - a_3 = 3, \dots$

$$a_n = 1 + \frac{n(n-1)}{2} = \frac{n^2 - n + 2}{2}$$

$$\therefore \text{Sum} = \frac{1}{2} \left\{ \frac{30 \cdot 31 \cdot 61}{6} - \frac{30 \cdot 31}{2} + 2 \cdot 30 \right\} = 4525$$

$$6. (b): \frac{1}{2r^2} = \frac{2}{4r^2} = \frac{(2r+1) - (2r-1)}{1 + (4r^2 - 1)}$$

$$\text{Sum} = \sum (\tan^{-1}(2r+1) - \tan^{-1}(2r-1))$$

$$= \tan^{-1}(2n+1) - \tan^{-1}(1) = \tan^{-1}\left(\frac{n}{n+1}\right)$$

7. (b): $p \rightarrow (q \rightarrow p) \equiv \sim p \vee (q \rightarrow p)$

$$\equiv \sim p \vee (\sim q \vee p) \equiv p \rightarrow (p \vee q)$$

8. (b): $(1 + 63)^n - (63 - 1)^{2n+1}$

Remainder is 2.

9. (d): Total Function = 5^3


$$10. (d): \text{Let } \ell = \int_0^{\pi} [\cot x] dx \Rightarrow \ell = \int_0^{\pi} [\cot(\pi - x)] dx$$

$$\therefore \ell + \ell = \int_0^{\pi} (-1) dx \left[\because [x] + [-x] = \begin{cases} -1 & \text{if } x \notin \mathbb{Z} \\ 0 & \text{if } x \in \mathbb{Z} \end{cases} \right]$$

$$\therefore \ell = -\frac{\pi}{2}$$




AVAILABLE BOUND VOLUMES




Physics For You 2016

₹ 325




Chemistry Today 2016

₹ 325



Biology Today 2016

₹ 325



Mathematics Today 2015

₹ 325

Mathematics Today 2016

₹ 325

Mathematics Today 2014

₹ 300

Mathematics Today 2013

₹ 300

Physics For You 2016
 April, May, June issues not
 included (9 issues)

₹ 240

buy online at www.mtg.in

of your favourite magazines

How to order : Send money by demand draft/money order. Demand Draft should be drawn in favour of **MTG Learning Media (P) Ltd.** Mention the volume you require along with your name and address.

Add ₹ 60 as postal charges


Mail your order to :

Circulation Manager, MTG Learning Media (P) Ltd.
 Plot 99, Sector 44 Institutional Area, Gurgaon, (HR)
 Tel.: (0124) 6601200
 E-mail : info@mtg.in Web : www.mtg.in

FULLY LOADED & COMPLETELY UPDATED

mtg

MTG's BITSAT Explorer is not only the most exhaustive prep-tool, but also the only book available at present, updated as per the latest BITSAT syllabus for students aspiring for top rank in BITSAT 2017.



Strictly based on revised BITSAT syllabus



Free Interactive CD with MTG's BITSAT Explorer. Simulate the online testing experience with this unique interactive CD.

Runs on both PCs and Macs.

₹ 750


Get MTG's BITSAT Explorer today for a real-world feel of BITSAT. Find out what's different about the BITSAT test, including its pattern of examination and key success factors. Be it with chapter-wise MCQs or model test papers, check how good your chances are for glory in BITSAT 2017.

FEATURES:

- Covers all 5 subjects - Physics, Chemistry, Mathematics, English & Logical Reasoning
- Chapterwise 1,000+ MCQs in each section for practice
- 5 Model Test Papers with detailed solutions
- Free interactive CD

Visit www.MTG.in to buy online. Or visit a leading bookseller near you.
For more information, email info@mtg.in or call 1800 300 23355 (toll-free).

OLYMPIAD CORNER



1. $a_1, \dots, a_k, a_{k+1}, \dots, a_n$ are positive numbers ($k < n$). Suppose that the values of a_{k+1}, \dots, a_n are fixed. How should one choose the values of a_1, \dots, a_k in

order to minimize $\sum_{i,j,i \neq j} \frac{a_i}{a_j}$?

2. Let m be a positive integer. Define the sequence a_0, a_1, a_2, \dots by $a_0 = 0, a_1 = m$ and $a_{n+1} = m^2 a_n - a_{n-1}$ for $n = 1, 2, 3, \dots$. Prove that an ordered pair (a, b) of non-negative integers, with $a \leq b$, gives a solution to the equation

$$\frac{a^2 + b^2}{ab + 1} = m^2$$

if and only if (a, b) is of the form (a_n, a_{n+1}) for some $n \geq 0$.

3. In a $\triangle ABC$, $\angle C = 2\angle B$. P is a point in the interior of $\triangle ABC$ satisfying that $AP = AC$ and $PB = PC$. Show that AP trisects $\angle A$.
4. Determine all the possible values of the sum of the digits of the perfect squares.
5. $ABCD$ is a convex quadrilateral and O is the intersection of its diagonals. Let L, M, N be the mid-points of DB, BC, CA respectively. Suppose that AL, OM, DN are concurrent. Show that either $AD \parallel BC$ or $[ABCD] = 2[OBC]$.

SOLUTIONS

1. To minimize the given rational function, choose

$$a_i = \left(\frac{a_{k+1} + \dots + a_n}{\frac{1}{a_{k+1}} + \dots + \frac{1}{a_n}} \right)^{1/2} = (A \cdot H)^{1/2}, i = 1, 2, \dots, k$$

where A is the arithmetic mean and H is the harmonic mean of a_{k+1}, \dots, a_n .

To prove this, we will be forgiven if we change notation: let $x_i = a_i, i = 1, 2, \dots, k$ and $b_r = a_{k+r}, r = 1, \dots, m$ with $k + m = n$ and denote the given rational function $F(x_1, \dots, x_k)$.

Then we have $F(x_1, \dots, x_k) = X + Y + B$, where

$$X = \sum_{1 \leq i < j \leq k} \left(\frac{x_i}{x_j} + \frac{x_j}{x_i} \right),$$

$$Y = \sum_{1 \leq i \leq k} \sum_{1 \leq r \leq m} \left(\frac{x_i}{b_r} + \frac{b_r}{x_i} \right),$$

$$B = \sum_{1 \leq r < s \leq m} \left(\frac{b_r}{b_s} + \frac{b_s}{b_r} \right).$$

Note that B is fixed and Y can be improved to

$$\begin{aligned} Y &= \sum_{1 \leq i \leq k} \left(\left(\sum_{1 \leq r \leq m} \frac{1}{b_r} \right) x_i + \left(\sum_{1 \leq r \leq m} b_r \right) \frac{1}{x_i} \right) \\ &= \sum_i \left(\frac{m}{H} x_i + \frac{mA}{x_i} \right) \end{aligned}$$

where A is the arithmetic mean and H is the harmonic mean of the b_r .

Now we recall that the simple function $\alpha x + \frac{\beta}{x}$ (with α, β, x all positive) assumes its minimum when $\alpha x = \frac{\beta}{x}$; that is $x = \sqrt{\beta/\alpha}$. Thus each of the terms in Y (and so Y itself) assumes its minimum when we choose, for $i = 1, 2, \dots, k$,

$$x_i = \sqrt{\frac{mA}{m/H}} = \sqrt{AH}, \text{ as asserted.}$$

But there is more. It is also known that each term in X , (and so X itself) assumes its minimum when $x_i = x_j$, with $1 \leq i < j \leq k$. Thus choosing all $x_i = \sqrt{AH}$ minimizes both X and Y and, since B

is fixed, minimizes $F(x_1, \dots, x_k)$ as claimed.

2. Let us first prove by induction that

$$\frac{a_n^2 + a_{n+1}^2}{a_n \cdot a_{n+1} + 1} = m^2 \text{ for all } n \geq 0.$$

$$\text{Proof: Base case } (n=0): \frac{a_0^2 + a_1^2}{a_0 \cdot a_1 + 1} = \frac{0 + m^2}{0 + 1} = m^2.$$

Now, let us assume that it is true for $n = k, k \geq 0$.

$$\text{Then, } \frac{a_k^2 + a_{k+1}^2}{a_k \cdot a_{k+1} + 1} = m^2$$

$$a_k^2 + a_{k+1}^2 = m^2 \cdot a_k \cdot a_{k+1} + m^2$$

$$\begin{aligned} a_{k+1}^2 + m^4 a_{k+1}^2 - 2m^2 \cdot a_k \cdot a_{k+1} + a_k^2 \\ = m^2 + m^4 a_{k+1}^2 - m^2 \cdot a_k \cdot a_{k+1} \end{aligned}$$

$$a_{k+1}^2 + (m^2 a_{k+1} - a_k)^2 = m^2 + m^2 a_{k+1} (m^2 a_{k+1} - a_k)$$

$$a_{k+1}^2 + a_{k+2}^2 = m^2 + m^2 \cdot a_{k+1} \cdot a_{k+2}$$

$$\text{Therefore, } \frac{a_{k+1}^2 + a_{k+2}^2}{a_{k+1} \cdot a_{k+2} + 1} = m^2, \text{ proving the}$$

induction. Hence (a_n, a_{n+1}) is a solution to

$$\frac{a^2 + b^2}{ab + 1} = m^2 \text{ for all } n \geq 0.$$

Now, consider the equation $\frac{a^2 + b^2}{ab + 1} = m^2$ and suppose

$(a, b) = (x, y)$ is a solution with $0 \leq x \leq y$. Then

$$\frac{x^2 + y^2}{xy + 1} = m^2 \quad \dots(1)$$

If $x = 0$ then it is easily seen that $y = m$, so $(x, y) = (a_0, a_1)$. Since we are given $x \geq 0$, suppose now that $x > 0$.

Let us show that $y \leq m^2 x$.

Proof by contradiction: Assume that $y > m^2 x$. Then $y = m^2 x + k$ where $k \geq 1$.

Substituting into (1) we get

$$\frac{x^2 + (m^2 x + k)^2}{(x)(m^2 x + k) + 1} = m^2$$

$$x^2 + m^4 x^2 + 2m^2 xk + k^2 = m^4 x^2 + m^2 kx + m^2$$

$$(x^2 + k^2) + m^2(kx - 1) = 0.$$

Now, $m^2(kx - 1) \geq 0$ since $kx \geq 1$ and $x^2 + k^2 \geq x^2 + 1 \geq 1$ so $(x^2 + k^2) + m^2(kx - 1) \neq 0$.

Thus we have a contradiction, so $y \leq m^2 x$ if $x > 0$.

Now substitute $y = m^2 x - x_1$, where $0 \leq x_1 < m^2 x$, into (1).

We have

$$\frac{x^2 + (m^2 x - x_1)^2}{x(m^2 x - x_1) + 1} = m^2$$

$$\begin{aligned} x^2 + m^4 x^2 - 2m^2 x \cdot x_1 + x_1^2 &= m^4 x^2 - m^2 x \cdot x_1 + m^2 \\ x^2 + x_1^2 &= m^2(x \cdot x_1 + 1) \end{aligned}$$

$$\frac{x^2 + x_1^2}{x \cdot x_1 + 1} = m^2 \quad \dots(2)$$

If $x_1 = 0$, then $x^2 = m^2$. Hence $x = m$ and

$(x_1, x) = (0, m) = (a_0, a_1)$. But $y = m^2 x - x_1 = a_2$, so $(x, y) = (a_1, a_2)$.

Thus suppose $x_1 > 0$.

Let us now show that $x_1 < x$.

Proof by contradiction: Assume $x_1 \geq x$.

Then $m^2 x - y \geq x$, since $y = m^2 x - x_1$, and

$\left(\frac{x^2 + y^2}{xy + 1} \right) x - y \geq x$, since (x, y) is a solution to

$$\frac{a^2 + b^2}{ab + 1} = m^2.$$

So $x^3 + xy^2 \geq x^2 y + xy^2 + x + y$.

Hence $x^3 \geq x^2 y + x + y$, which is a contradiction since $y \geq x > 0$.

With the same proof that $y \leq m^2 x$, we have $x \leq m^2 x_1$. So the substitution $x = m^2 x_1 - x_2$ with $x_2 \geq 0$ is valid.

Substituting $x = m^2 x_1 - x_2$ into (2) gives

$$\frac{x_1^2 + x_2^2}{x_1 \cdot x_2 + 1} = m^2.$$

If $x_2 \neq 0$, then we continue with the substitution

$$x_i = m_{x_{i+1}}^2 - x_{i+2} (*) \text{ until we get } \frac{x_j^2 + x_{j+1}^2}{x_j \cdot x_{j+1} + 1} = m^2$$

and $x_{j+1} = 0$. (The sequence x_i is decreasing, non-negative and integer.)

So, if $x_{j+1} = 0$, then $x_j^2 = m^2$ so $x_j = m$ and

$$(x_{j+1}, x_j) = (0, m) = (a_0, a_1).$$

Then $(x_j, x_{j-1}) = (a_1, a_2)$ since $x_{j-1} = m^2 x_j - x_{j+1}$ (from (*)).

Continuing, we have $(x_1, x) = (a_{n-1}, a_n)$ for some n . Then $(x, y) = (a_n, a_{n+1})$.

Hence $\frac{a^2 + b^2}{ab + 1} = m^2$ has solutions (a, b) if and

only if $(a, b) = (a_n, a_{n+1})$ for some n .

3. Let $\angle PAC$ and $\angle BAP$ be 2α and β respectively. Then, since $\angle C = 2\angle B$, we deduce from $A + B + C = 180^\circ$ that $2\alpha + \beta + 3B = 180^\circ$.

The angles at the base of the isosceles triangle PAC are each $90^\circ - \alpha$. Also $\triangle BPC$ is isosceles, having base angles $C - (90^\circ - \alpha) = 2B + \alpha - 90^\circ$, and so $\angle BPA = 180^\circ - (\angle PBA + \angle BAP) = 180^\circ - [B - (2B + \alpha - 90^\circ) + 180^\circ - 2\alpha - 3B] = 4B + 3\alpha - 90^\circ$.

As usual, let a , b and c denote the lengths of the sides BC , AC and AB . By the Law of Cosines, applied to $\triangle BPA$, where $PA = b$ and $PB = PC = 2b \sin \alpha$,

$$c^2 = b^2 + (2b \sin \alpha)^2 - 2 \cdot b \cdot 2b \sin \alpha \cdot \cos(4B + 3\alpha - 90^\circ),$$

so that

$$c^2 = b^2 [1 + 4 \sin^2 \alpha - 4 \sin \alpha \sin(4B + 3\alpha)] \dots (2)$$

We now use the fact that $\angle C = 2\angle B$ is equivalent to the condition $c^2 = b(b + a)$.

Since $a = 2 \cdot PC \cdot \cos(2B + \alpha - 90^\circ) = 4b \sin \alpha \sin(2B + \alpha)$, we have

$$c^2 = b^2 [1 + 4 \sin \alpha \sin(2B + \alpha)] \dots (3)$$

Therefore, from (2) and (3), we get

$$b^2 [1 + 4 \sin^2 \alpha - 4 \sin \alpha \sin(4B + 3\alpha)] = b^2 [1 + 4 \sin \alpha \sin(2B + \alpha)],$$

which simplifies to

$$\sin \alpha - \sin(4B + 3\alpha) = \sin(2B + \alpha).$$

Since $\sin \alpha - \sin(4B + 3\alpha) = -2 \cos(2B + 2\alpha) \sin(2B + \alpha)$, this equation may be rewritten as $\sin(2B + \alpha) [1 + 2 \cos(2B + 2\alpha)] = 0$

Since, from (1), $2B + \alpha < 180^\circ$, we must have $1 + 2 \cos(2B + 2\alpha) = 0$, giving $\cos(2B + 2\alpha) = -1/2$; that is,

$$2B + 2\alpha = 120^\circ \dots (4)$$

Since, again from (1), $2B + 2\alpha < 180^\circ$


Finally, we may eliminate B between (1) and (4) to obtain $\alpha = \beta$. The result follows.

4. The squares can only be 0, 1, 4 or 7 mod 9.

Thus the sum of the digits of a perfect square cannot be 2, 3, 5, 6 or 8 mod 9, since the number itself would then be 2, 3, 5, 6 or 8 mod 9.

We shall show that the sum of the digits of a perfect square can take every value of the form 0, 1, 4 or 7 mod 9.

$$(10^m - 1)^2 = 10^{2m} - 2 \cdot 10^m + 1 = \underbrace{99 \dots 980}_{m-1} \dots \underbrace{01}_{m-1}, m \geq 1.$$



The sum of the digits is $9m$, giving all the values greater than or equal to 9 congruent to 0 mod 9

$$(10^m - 2)^2 = 10^{2m} - 4 \cdot 10^m + 4 = \underbrace{99 \dots 960}_{m-1} \dots \underbrace{04}_{m-1}, m \geq 1.$$

The sum of the digits is $9m + 1$, which gives all values greater than or equal to 10 congruent to 1 mod 9.

$$(10^m - 3)^2 = 10^{2m} - 6 \cdot 10^m + 9 = \underbrace{99 \dots 940}_{m-1} \dots \underbrace{09}_{m-1}, m \geq 1.$$

The sum of the digits is $9m + 4$, which takes every value greater than or equal to 13 which is congruent to 4 mod 9

$$(10^m - 5)^2 = 10^{2m} - 10^{m+1} + 25 = \underbrace{9 \dots 900}_{m-1} \dots \underbrace{025}_{m-1}.$$

The sum of the digits is $9(m - 1) + 7 = 9m - 2$, from which we get every value greater than or equal to 7 congruent to 7 mod 9.

We have taken care of all the integers apart from 0, 1, 4, which are the sums of the digits of 0^2 , 1^2 and 2^2 respectively.

5. Let O be the origin of a coordinate system where A , B , C , D are represented by $(a, 0)$, $(0, b)$, $(c, 0)$, $(0, d)$ with a, b positive and c, d negative. Thus L is the point

$$\left(0, \frac{(b+d)}{2}\right), M \text{ is } \left(\frac{c}{2}, \frac{b}{2}\right), N \text{ is } \left(\frac{(a+c)}{2}, 0\right) \text{ and}$$

$$AL : (b + d)x + 2ay - a(b + d) = 0$$

$$OM : bx - cy = 0$$

$$DN : 2dx + (a + c)y - d(a + c) = 0.$$

These lines are concurrent if and only if

$$\begin{vmatrix} b & -c & 0 \\ b+d & 2a & -a(b+d) \\ 2d & a+c & -d(a+c) \end{vmatrix} = 0.$$

This equation reduces (after some manipulation) to $(ab - cd) [(a - c)(b - d) + 2bc] = 0$.

Consequently, either

(a) $ab = cd$, in which case $AD \parallel BC$, or

$$(b) \frac{1}{2}(a - c)(b - d) \sin \alpha = 2 \left(-\frac{1}{2} bc \sin \alpha \right)$$

(where $\alpha = \angle AOB$), in which case $[ABCD] = 2 [OBC]$.



VIT Engineering Entrance Exam (VITEEE-2017) Results

VITEEE - 2017 TOP 10 RANK HOLDERS



RANK 1
AASHISH WAIKAR



RANK 2
DIVYANSH TRIPATHI



RANK 3
DIVYANSHU MANDOWARA



RANK 4
ABHISHEK RAO



RANK 5
BHANUTEJA BOLISETTI



RANK 6
HRITWIK SINGHAI



RANK 7
**PRATHEEK D SOUZA
REBELLO**



RANK 8
**AVVARI SAI S V
BHARADWAJ**



RANK 9
**PATEL MANAN
BRIJESH**



RANK 10
SHOURYA AGGARWAL

AASHISH WAIKAR, a student of BHAVAN VIDYALAYA PANCHKULA, Madhya Pradesh has secured the first rank in the VIT Engineering Entrance Examination (VITEEE)-2017 which was held from April 5th to 16 in 119 selected cities across India, as well as Dubai, Kuwait and Muscat. The entrance exam was held for admission to the various B.Tech programmes offered by VIT University at its Vellore, Chennai, Bhopal & Amaravati (AP).

Releasing the results, VIT Chancellor Dr.G.Viswanathan said that a record 2,23,081 candidates had registered for the VITEEE-2017. The other rank holders among the top 10 are 2nd rank: DIVYANSH TRIPATHI (Prabhat Sr Sec Public School, Uttar Pradesh), 3rd rank: DIVYANSHU MANDOWARA (Arcadia Academy Co-Educational English Medium Senior Secondary School, Rajasthan), 4th rank: ABHISHEK RAO (Remal Public School, Uttar Pradesh), 5th rank : BHANUTEJA BOLISETTI (Sri Chaitanya Narayana Jr College, Telengana), 6th rank: HRITWIK SINGHAI (Little Kingdom Senior Secondary School, Madhya Pradesh), 7th rank: PRATHEEK D SOUZA REBELLO (Mushtifund Aryaan Higher Secondary School, GOA), 8th rank : AVVARI SAI S V BHARADWAJ (Sri Chaitanya Junior Kalasala, Telengana), 9th rank: PATEL MANAN BRIJESH (Shree Swaminarayan Secondary School, Gujarat) and 10th

rank : SHOURYA AGGARWAL (Hans Raj Model School, Delhi).

Dr. G. Viswanathan said that admissions would be only on merit, based on the marks obtained by the candidates in the VITEEE. The results have been released through the www.vit.ac.in.

Counselling for candidates, who obtained ranks upto 8,000 was held on May 10 and counseling for ranks 8001 to 14,000 was held on May 11 while for those who secured ranks from 14001 to 20000 was held on May 12. The counselling was held simultaneously in the Vellore, Chennai, Bhopal and Amaravati (AP).

Under the G V School Development Programme central and State board toppers would be given 100 percent fee waiver for all the four years. Candidates with ranks upto 50 would be given a 75% tuition fee waiver, Rank 51 to 100 would be given a 50% tuition fee waiver and Rank 101 to 1000 would be given a 25 % tuition fee waiver.

Each one boy and one girl secured top ranks in "Plus2" at district level from state board schools who also appeared for VIT Engineering Entrance Examination will be given 100% fee concession and free boarding and lodging in the hostels of VIT under STARS scheme.



YOU ASK WE ANSWER

Do you have a question that you just can't get answered?

Use the vast expertise of our MTG team to get to the bottom of the question. From the serious to the silly, the controversial to the trivial, the team will tackle the questions, easy and tough. The best questions and their solutions will be printed in this column each month.

1. For $r = 0, 1, \dots, 10$, let A_r, B_r, C_r denote respectively, the coefficients of x^r in the expansion of $(1+x)^{10}$,

$$(1+x)^{20}, (1+x)^{30}. \text{ Then } \sum_{r=1}^{10} A_r (B_{10} B_r - C_{10} A_r) =$$

(Ram Krishan, West Bengal)

Ans. $(1+x)^{10} = A_0 + A_1 x + A_2 x^2 + \dots + A_{10} x^{10}$
 $(x+1)^{20} = B_0 x^{20} + B_1 x^{19} + B_2 x^{18} + \dots + B_{20}$.
 Considering the coefficient of x^{20} in the product, we get
 $A_0 B_0 + A_1 B_1 + A_2 B_2 + \dots + A_{10} B_{10} = \text{coefficient of } x^{20} \text{ in the expansion of } (1+x)^{10}(x+1)^{20} = (1+x)^{30}$
 which is C_{20}

$$\therefore \sum_{r=0}^{10} A_r B_r = C_{20} = C_{10}$$

$$\text{But } \binom{n}{0}^2 + \binom{n}{1}^2 + \binom{n}{2}^2 + \dots + \binom{n}{n}^2 = \binom{2n}{n}$$

$$\therefore A_0^2 + A_1^2 + A_2^2 + \dots + A_n^2 = \binom{20}{10} = B_{10}$$

$$\sum_{r=0}^{10} A_r^2 = B_{10}. \text{ Now, } \sum_{r=1}^{10} A_r (B_{10} B_r - C_{10} A_r)$$

$$= B_{10} \sum_{r=1}^{10} A_r B_r - C_{10} \sum_{r=1}^{10} A_r^2$$

$$= B_{10} (C_{10} - 1) - C_{10} (B_{10} - 1) = C_{10} - B_{10}.$$

2. If $\begin{bmatrix} 4a^2 & 4a & 1 \\ 4b^2 & 4b & 1 \\ 4c^2 & 4c & 1 \end{bmatrix} \begin{bmatrix} f(-1) \\ f(1) \\ f(2) \end{bmatrix} = \begin{bmatrix} 3a^2 + 3a \\ 3b^2 + 3b \\ 3c^2 + 3c \end{bmatrix}$, $f(x)$ is a

quadratic function and its maximum value occurs at a point V . A is a point of intersection of $y = f(x)$ with x -axis and point B is such that the chord AB subtends a right angle at V . Find the area enclosed by $y = f(x)$ and the chord AB .

(Suresh Prasad, Jharkhand)

Ans. The given equation implies, $(4f(-1) - 3)x^2 + (4f(1) - 3)x + f(2) = 0$ is satisfied by 3 roots a, b, c

$$\Rightarrow \text{It is an identity } \therefore f(-1) = \frac{3}{4}, f(1) = \frac{3}{4}, f(2) = 0$$

$$\text{If } f(x) = \alpha x^2 + \beta x + \gamma, \text{ then } \beta = 0, \alpha = -\frac{1}{4}, \gamma = 1$$

$$\Rightarrow f(x) = -\frac{x^2}{4} + 1 = \frac{4-x^2}{4}$$

The maximum point, $V = (0, 1)$; $A(-2, 0)$. Taking $B = (2t, 1-t^2)$

$$\angle AVB = \frac{\pi}{2} \Rightarrow \frac{1}{2} \cdot \left(-\frac{t}{2}\right) = -1 \Rightarrow t = 4 \therefore B = (8, -15)$$

$$\text{Equation of } AB \text{ is } 3x + 2y + 6 = 0$$

$$\text{The area required is } \int_{-2}^8 \left(\frac{4-x^2}{4} + \frac{3(x+2)}{2} \right) dx = 41.67$$

3. Let ABC be an equilateral triangle inscribed in the circle $x^2 + y^2 = a^2$. Suppose perpendiculars from A ,

$$B, C \text{ to the major axis meet the ellipse } \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

($a > b$) respectively at P, Q, R so that P, Q, R lie on the same side of the major axis as A, B, C respectively. Prove that the normals to the ellipse at P, Q, R are concurrent.

(Priyanshu Sharma, Bihar)

Ans. A, B, C are the vertices of an equilateral triangle.

$$A(a \cos \theta, a \sin \theta),$$

$$B\left(a \cos\left(\theta + \frac{2\pi}{3}\right), a \sin\left(\theta + \frac{2\pi}{3}\right)\right),$$

$$C\left(a \cos\left(\theta + \frac{4\pi}{3}\right), a \sin\left(\theta + \frac{4\pi}{3}\right)\right)$$

$$\text{Hence, } P(a \cos \theta, b \sin \theta),$$

$$Q\left(a \cos\left(\theta + \frac{2\pi}{3}\right), b \sin\left(\theta + \frac{2\pi}{3}\right)\right),$$

$$R\left(a \cos\left(\theta + \frac{4\pi}{3}\right), b \sin\left(\theta + \frac{4\pi}{3}\right)\right)$$

The normals to the ellipse at P, Q, R are respectively,

$$L_1 \equiv ax \sin \theta - by \cos \theta - \frac{a^2 - b^2}{2} \sin 2\theta = 0;$$

$$L_2 \equiv ax \sin\left(\theta + \frac{2\pi}{3}\right) - by \cos\left(\theta + \frac{2\pi}{3}\right) - \frac{a^2 - b^2}{2}$$

$$\sin^2\left(\theta + \frac{2\pi}{3}\right) = 0; L_3 \equiv ax \sin\left(\theta + \frac{4\pi}{3}\right)$$

$$- by \cos\left(\theta + \frac{4\pi}{3}\right) - \frac{a^2 - b^2}{2} \sin 2\left(\theta + \frac{4\pi}{3}\right) = 0$$

$$\text{Since, } \sin \theta + \sin\left(\theta + \frac{2\pi}{3}\right) + \sin\left(\theta + \frac{4\pi}{3}\right) = 0$$

$$\cos \theta + \cos\left(\theta + \frac{2\pi}{3}\right) + \cos\left(\theta + \frac{4\pi}{3}\right) = 0$$

$$\sin 2\theta + \sin 2\left(\theta + \frac{2\pi}{3}\right) + \sin 2\left(\theta + \frac{4\pi}{3}\right) = 0$$

Hence L_1, L_2, L_3 are concurrent.



MATHS MUSING

SOLUTION SET-173

1. (c) : Anil	Balu	Probability
R, R	R, R	1/33
B, B	B, B	1/33
R, B	R, B	10/33

$$\therefore \text{Required probability} = \frac{2}{33} + \frac{10}{33} = \frac{12}{33} = \frac{4}{11}$$

2. (b) : Let (h, k) be the mid point of the chord of circle $x^2 + y^2 = a^2$, then the equation of the chord be $S_1 = T$
 $\Rightarrow xh + yk - a^2 = h^2 + k^2 - a^2$
 $\Rightarrow h^2 + k^2 = xh + yk$
 $\Rightarrow h^2 + k^2 = ah + bk$ (As it passes through (a, b))
 $\Rightarrow x^2 + y^2 = ax + by$ (By changing the locus of $h, k \rightarrow x, y$)

3. (b) : Let $I = \int_0^{\pi/2} \frac{dx}{\cos^6 x + \sin^6 x} \Rightarrow I = \int_0^{\pi/2} \frac{\sec^6 x}{1 + \tan^6 x} dx$

$$\Rightarrow I = \int_0^{\pi/2} \frac{(1 + \tan^2 x)^3}{1 + \tan^6 x} dx$$

$$\Rightarrow I = \int_0^{\pi/2} \frac{1 + \tan^6 x + 3 \tan^2 x (1 + \tan^2 x)}{1 + \tan^6 x} dx$$

$$I = \int_0^{\pi/2} 1 + \frac{3 \tan^2 x \sec^2 x dx}{1 + (\tan^3 x)^2}$$

Put $\tan^3 x = t \Rightarrow dt = 3 \tan^2 x \sec^2 x dx$

$$\Rightarrow I = \int_0^{\pi/2} dx + \int_0^{\infty} \frac{dt}{1+t^2} = \frac{\pi}{2} + \tan^{-1}(t) \Big|_0^{\infty} = \frac{\pi}{2} + \frac{\pi}{2} = \pi$$

4. (d) : Consider $C_r = \binom{24}{r}$

$$(1-x)^{24} = C_0 - C_1x + C_2x^2 - \dots + C_{16}x^{16} + \dots$$

$$(1-x)^{-1} = 1 + x + x^2 + \dots + x^{16} + \dots$$

Considering the coefficient of x^{16} in the product of these two,

$$C_0 - C_1 + C_2 - \dots + C_{16} = \text{coeff. of } x^{16} \text{ in } (1-x)^{23}$$

$$= \binom{23}{16} = \binom{23}{7}$$

5. (a) : Let the A.P.s. be $a, a + \alpha, a + 2\alpha, \dots$ and $b, b + \beta, b + 2\beta, \dots$

$$\Rightarrow ab = a_1b_1 = 120, (a + \alpha)(b + \beta) = a_2b_2 = 143,$$

$$(a + 2\alpha)(b + 2\beta) = a_3b_3 = 154$$

$$\Rightarrow ab = 120, a\beta + b\alpha = 29, \alpha\beta = -6$$

$$\therefore a_8b_8 = (a + 7\alpha)(b + 7\beta) = ab + 7(a\beta + b\alpha) + 49\alpha\beta$$

$$= 120 + 7 \times 29 - 49 \times 6 = 29$$

6. (b, c) : $f(x) = \log_e [x^3 + \sqrt{x^6 + 1}]$

$$\therefore f(-x) = \log_e [-x^3 + \sqrt{x^6 + 1}]$$

$$\therefore f(x) + f(-x) = \log_e 1 = 0$$

$$f(-x) = -f(x)$$

Hence $f(x)$ is an odd function.

\therefore (b) is correct.

$$\text{Again, } f'(x) = \frac{1}{x^3 + \sqrt{x^6 + 1}} \left[3x^2 + \frac{6x^5}{2\sqrt{x^6 + 1}} \right]$$

$$= \frac{1}{x^3 + \sqrt{x^6 + 1}} 3x^2 \left[\frac{\sqrt{x^6 + 1} + x^3}{\sqrt{x^6 + 1}} \right] = \frac{3x^2}{\sqrt{x^6 + 1}} > 0$$

$\therefore f(x)$ is an increasing function, so (c) is correct.

7. (b) : $\angle DBC = \alpha \Rightarrow BD = 20, \angle EBD = \alpha$

By sine rule for $\triangle EBD$, we get

$$\frac{\sin 3\alpha}{20} = \frac{\sin \alpha}{8}$$


$$\Rightarrow 3 - 4 \sin^2 \alpha = \frac{5}{2}$$

$$\Rightarrow \sin^2 \alpha = \frac{1}{8}, \cos 2\alpha = \frac{3}{4}$$

8. (d) : $AE + 28 = AB \cot \alpha$ and $AE + 8 = AB \cot 2\alpha$
 On subtracting, we get $20 = AB (\cot \alpha - \cot 2\alpha)$

$$\therefore AB = 20 \sin 2\alpha = 20 \sqrt{1 - \frac{9}{16}} = 5\sqrt{7}$$

$$\therefore AE + 8 = AB \cot 2\alpha = 5\sqrt{7} \cdot \frac{3}{\sqrt{7}} = 15 \Rightarrow AE = 7.$$



9. (8) : $a + b + c = 0 \Rightarrow a, b, c$ are the roots of $x^3 + qx + r = 0$... (i)

$$\Sigma ab = q, abc = -r, a + b + c = 0$$

$$\Rightarrow \Sigma a^3 = 3abc \therefore abc = 1, r = -1, \Sigma a^2 = -2q$$

$$\text{From (i), we get } \Sigma a^5 + q \Sigma a^3 + r \Sigma a^2 = 0 \Rightarrow 10 + 3q + 2q = 0$$

$$\Rightarrow q = -2$$

$$\text{From (i), } x^3 - 2x - 1 = 0.$$

$$\therefore \Sigma a^3 - 2 \Sigma a - \Sigma 1 = 0 \Rightarrow \Sigma a^4 = 2 \Sigma a^2 + \Sigma a$$

$$\Sigma a^4 = 2(-2q) + 0 = (-4) \times (-2) = 8.$$

10. (b) : P. $23^2 \equiv -1 \pmod{53}, 23^{22} \equiv -1 \pmod{53},$
 $23^{23} \equiv -23 \pmod{53} \equiv 30 \pmod{53}$

Q. $X = x + 2, Y = y + 1, Z = z, U = u - 1$

$$\Rightarrow X + Y + Z + U = 5$$





$$\therefore \text{Number of solutions} = \binom{8}{3} = 56$$

R. Coefficient of x^2y in $(1 + x + 2y)^5$ is $\frac{5!}{2! \cdot 2!} \cdot 2 = 60$

S. $\sum_{r=1}^{10} r \frac{(11-r)}{r} = 1 + 2 + \dots + 10 = 55.$



Now, save up to Rs 2,020*



Subscribe to MTG magazines today.

Our 2017 offers are here. Pick the combo best suited for your needs. Fill-in the Subscription Form at the bottom and mail it to us today. If in a rush, log on to www.mtg.in now to subscribe online.

*On cover price of ₹ 30/- each.

**For JEE
(Main & Advanced),
NEET, AIIMS AND
JIPMER**

About MTG's Magazines

Perfect for students who like to prepare at a steady pace, MTG's magazines-Physics For You, Chemistry Today, Mathematics Today & Biology Today-ensure you practice bit by bit, month by month, to build all-round command over key subjects. Did you know these magazines are the only source for solved test papers of all national and state level engineering and medical college entrance exams?

Trust of over 1 Crore readers since 1982.

- Practice steadily, paced month by month, with very-similar & model test papers
- Self-assessment tests for you to evaluate your readiness and confidence for the big exams
- Content put together by a team

- comprising experts and members from MTG's well-experienced Editorial Board
- Stay up-to-date with important information such as examination dates, trends & changes in syllabi
- All-round skill enhancement –

- confidence-building exercises, new studying techniques, time management, even advice from past JEE/NEET toppers
- **Bonus:** Exposure to competition at a global level, with questions from Intl. Olympiads & Contests

SUBSCRIPTION FORM

Please accept my subscription to:

Note: Magazines are despatched by Book-Post on 4th of every month (each magazine separately).

☒ Tick the appropriate box.

PCMB combo

☐ 1 yr: ₹ 1,000 (save ₹ 440) ☐ 2 yr: ₹ 1,800 (save ₹ 1,080) ☐ 3 yr: ₹ 2,300 (save ₹ 2,020)

PCM combo

☐ 1 yr: ₹ 900 (save ₹ 180) ☐ 2 yr: ₹ 1,500 (save ₹ 660) ☐ 3 yr: ₹ 1,900 (save ₹ 1,340)

PCB combo

☐ 1 yr: ₹ 900 (save ₹ 180) ☐ 2 yr: ₹ 1,500 (save ₹ 660) ☐ 3 yr: ₹ 1,900 (save ₹ 1,340)

Individual magazines

☐ Physics ☐ Chemistry ☐ Mathematics ☐ Biology

☐ 1 yr: ₹ 330 (save ₹ 30) ☐ 2 yr: ₹ 600 (save ₹ 120) ☐ 3 yr: ₹ 775 (save ₹ 305)

Enclose Demand Draft favouring **MTG Learning Media (P) Ltd.**, payable at New Delhi. You can also pay via Money Orders. Mail this Subscription Form to Subscription Dept., **MTG Learning Media (P) Ltd.**, Plot 99, Sector 44, Gurgaon - 122 003 (HR).

Want the magazines by courier; **add** the courier charges given below:

☐ 1 yr: ₹ 240 ☐ 2 yr: ₹ 450 ☐ 3 yr: ₹ 600

☒ Tick the appropriate box.

☐ Student ☒ Class XI ☒ XII ☐ Teacher ☐ Library ☐ Coaching

Name: _____

Complete Postal Address: _____

Pin Code Mobile #

Other Phone #

Email: _____

E-mail subscription@mtg.in. Visit www.mtg.in to subscribe online. Call (0)8800255334/5 for more info.